

LEADER PAIN RELIEF PATCHES- lidocaine patch
Cardinal Health, 110 dba LEADER

Leader PAIN RELIEF PATCHES

Active Ingredients

Lidocaine 4% Topical Anesthetic

Purpose

Temporarily relieves minor pain.

Dosage and Administration

Directions Adults and children over 12 years:

- clean and dry affected area
- remove backing from patch by firmly grasping both ends and gently pulling until backing separates in middle
- carefully remove smaller portion of backing from patch and apply exposed portion of patch to affected area
- once exposed portion of patch is positioned, carefully remove remaining backing to completely apply patch to affected area
- use one patch for up to 12 hours. Children under 12 years of age: consult a physician.

Warnings

For external use only.

Indications and Usage

Uses: Temporarily relieves minor pains.

When using this product

- use only as directed
- read and follow all directions and warnings on this carton
- do not allow contact with the eyes
- do not use at the same time as other topical analgesics
- do not bandage tightly or apply local heat (such as heating pads) to the area of use

- do not microwave

- dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and consult a doctor

- condition worsens

- redness is present

- irritation develops

- symptoms persist for more than 7 days or clear up and occur again within a few days

- you experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

If pregnant or breastfeeding, ask a health professional before use.

Do not Use

- more than one patch on your body at a time

- on cut, irritated or swollen skin

- on puncture wounds

- for more than one week without consulting a doctor

- if you are allergic to any active or inactive ingredients

- if pouch is damaged or opened.

Keep out of reach of children and pets.

If swallowed, get medical help or contact a Poison Control Center right away.

Other Safety Information

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive Ingredients

Aluminum Glycinate, Glycerin, Kaolin, Methylparaben, Polyacrylic Acid, Polysorbate 80, Propylene Glycol, Propylparaben, PVP, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water

Questions

TOLL-FREE

Made in China

Maximum Strength
Lidocaine Pain Relieving Patch
Lidocaine, 4.0% | Topical Anesthetic

Drug Facts

Active ingredient	Purpose
Lidocaine 4.0%	Topical anesthetic

Use: Temporarily relieves minor pain.

Warnings
For external use only

Do not use: More than one patch on your body at a time. On cut, irritated or swollen skin. On puncture wounds. For more than one week without consulting a doctor. If you are allergic to any active or inactive ingredients in this product. If pouch is damaged or opened.

When using this product: Use only as directed. Read and follow all directions and warnings on this carton. Avoid contact with the eyes, mucous membranes or rashes. Do not use at the same time as other topical anesthetics. Do not bandage tightly or apply heat (such as heating pad) to the area of use. Do not microwave. Dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and consult a doctor if: Condition worsens. If Redness is present. If Irritation develops. If Symptoms persist for more than 7 days or clear up and occur again within a few days. If You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center 800-222-1222 right away.

Directions
Adults and children 12 years of age and over: Clean and dry affected area. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Use one patch for up to 8 hours. Apply to affected area not more than 3 to 4 times daily. Discard patch after single use. Children under 12 years of age: consult a physician.

Other Information
Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive ingredients
aluminum glycolate, glycerin, isobutyl, methylparaben, polyacrylic acid, polysorbate 60, propylene glycol, propylparaben, PVP, sodium polyacrylate, tartaric acid, titanium dioxide, water.

Questions or comments? Call 1-888-328-1313

Maximum Strength
Lidocaine Pain Relieving Patch
Lidocaine, 4.0% | Topical Anesthetic

Maximum strength
Lidocaine Pain Relieving Patch
Lidocaine, 4.0%
Topical Anesthetic

Desensitizes Aggravated Nerves and Relieves Pain

No-Mess, Single-Use Application
Stay-Put Flexible Patch
Odor-Free

COMPARE TO SALONPAS®
MAXIMUM STRENGTH LIDOCAINE PATCH
active ingredient!

100% Money Back Guarantee

6 PATCHES
3.93 in. x 5.51 in. (10 cm x 14 cm)

LOT
EXP

*This product is not manufactured or distributed by Heumann Anesthetics, Inc., a distributor of Salonpas® Maximum Strength Lidocaine Patch.

OWNERSHIP BY: CARDINAL HEALTH
CARDINAL HEALTH
CARDINAL HEALTH
www.cardinal.com
1-800-368-4373
©2012 Cardinal Health. All Rights Reserved. LIDOCAINE, MAXIMUM STRENGTH, the Cardinal Health logo, SALONPAS, the SALONPAS logo, and the Leader logo are trademarks of Cardinal Health. All other trademarks are the property of their respective owners.

MADE IN CHINA

0 96295 13972 3

90-79048 06/20 VCE00046

✓ All LEADER® Brand Products Have A 100% Money Back Guarantee

Return to place of purchase if not satisfied.

4 g in 100 g

Inactive Ingredients				
Ingredient Name				Strength
ALUMINUM (UNII: CPD4NFA903)				
GLYCERIN (UNII: PDC6A3C0OX)				
KAOLIN (UNII: 24H4NWX5CO)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
POLYACRYLIC ACID (300000 MW) (UNII: A8371R0U5J)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)				
TARTARIC ACID (UNII: W4888I119H)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
WATER (UNII: 059QF0KO0R)				

Product Characteristics			
Color		Score	
Shape	RECTANGLE	Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0557-1	6 in 1 CARTON	04/01/2020	
1		9 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	04/01/2020	

Labeler - Cardinal Health, 110 dba LEADER (063997360)

Establishment			
Name	Address	ID/FEI	Business Operations
Foshan Aqua Gel Biotech Co.,Ltd.		529128763	manufacture(70000-0557)