

SCARAID SPF- silicone stick with spf stick
Biodermis, LLC

ScarAid SPF

DESCRIPTION:

PROSIL SPF15

SILICONE STICK FOR SCARS

GLYDE-ON THERAPY FOR YOUR SCARS

ACTIVE INGREDIENTS:

Avobenzone - 3% Sunscreen

Octocrylene - 5% Sunscreen agent

Octinoxate - 4% Sunscreen agent

DOSAGE & ADMINISTRATION:

Apply 15 minutes before sun exposure. Clean and dry scar site before each application. Apply ScarAid® liberally over the entire scar. May be covered with a bandage. Repeat several times daily for 8-12 weeks or until scar stops responding. To remove, wipe off with a clean cloth or tissue.

PURPOSE:

SILICONE STICK FOR SCARS

BROAD SPECTRUM UVA/UVB SUNSCREEN

HELPS PREVENT SUNBURN

INDICATIONS & USAGE:

SILICONE STICK FOR SCARS

BROAD SPECTRUM UVA/UVB SUNSCREEN

HELPS PREVENT SUNBURN

DIRECTIONS:

Apply 15 minutes before sun exposure. Clean and dry scar site before each application. Apply ScarAid® liberally over the entire scar. May be covered with a bandage. Repeat several times daily for 8-12 weeks or until scar stops responding. To remove, wipe off with a clean cloth or tissue. Store

between 55°F–95°F (13°C–35°C).

WARNINGS:

FOR EXTERNAL USE ONLY

WARNINGS AND PRECAUTIONS:

EXTERNAL USE ONLY

CAUTIONS: Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.

CAUTIONS:

Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.

KEEP OUT OF REACH OF CHILDREN:

KEEP OUT OF REACH OF CHILDREN:

ASK A DOCTOR:

Ask a doctor about using SCARAID SPF with children under 6 months old.

STORAGE:

Store between 55°F–95°F (13°C–35°C).

CONTACT INFORMATION:

MADE IN THE USA U.S. Patent No. 8,557,287

BIODERMIS LLC

www.biodermis.com

1820 Whitney Mesa Drive, Suite 100

Henderson, Nevada 89014 USA

800.322.3729 (USA)

702.260.4466 (INTL)

EC REP: MDSS GmbH, SCHIFFGRABEN 41 30175 HANNOVER, GERMANY


INACTIVE INGREDIENTS:

Silicone

Primary Package Label

4.25g Tube:

GLIDE-ON THERAPY FOR YOUR SCARS
PRODUCT SEALED FOR YOUR PROTECTION. BREAK SEAL TO OPEN


**STICK**
WITH SPF-15
First Aid for Scars

DIRECTIONS: Apply 15 minutes before sun exposure. Clean and dry scar site before each application. Apply Pro-Sil® liberally over the entire scar. May be covered with a bandage. Repeat several times daily for 8–12 weeks or until scar stops responding. To remove, wipe off with a clean cloth or tissue. Store between 55°F–95°F (13°C–35°C).

CAUTIONS: Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.


HELPS PREVENT SUNBURN BROAD SPECTRUM UVA/UVB SUNSCREEN

Drug Facts	Active Ingredients	Purpose
4.25g / .15 oz	Avobenzone (3%)	Sunscreen
	Octocrylene (5%)	Sunscreen Agent
	Octinoxate (4%)	Sunscreen Agent



EXTERNAL USE ONLY

MADE IN THE USA U.S. Patent No. 8,557,287

 **BIODERMIS LLC**
1820 Whitney Mesa Drive, Suite 100
Henderson, Nevada 89014 USA
800.322.3729 USA 702.260.4466 INTL
www.biodermis.com

EC


REP

MDSS GmbH, SCHIFFGRABEN 41
30175 HANNOVER, GERMANY

REF

XXXX-0000


MD



YYYY-MM-DD

LOT

00000



YYYY-MM-DD

17g Tube:

GLIDE-ON THERAPY FOR YOUR SCARS
PRODUCT SEALED FOR YOUR PROTECTION. BREAK SEAL TO OPEN

SCAR AID[®] STICK

WITH SPF-15
First Aid for Scars

DIRECTIONS: Apply 15 minutes before sun exposure. Clean and dry scar site before each application. Apply ScarAid[®] liberally over the entire scar. May be covered with a bandage. Repeat several times daily for 8–12 weeks or until scar stops responding. To remove, wipe off with a clean cloth or tissue. Store between 55°F–95°F (13°C–35°C).

CAUTIONS: Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.

BROAD SPECTRUM UVA/UVB SUNSCREEN HELPS PREVENT SUNBURN

U.S. Patent No. 8,557,287 MADE IN THE USA EXTERNAL USE ONLY

Drug Facts
17g / .60 oz

Active Ingredients	Purpose
Avobenzone (3%).....	Sunscreen
Octocrylene (5%)...	Sunscreen Agent
Octinoxate (4%)...	Sunscreen Agent



BIODERMIS LLC
1820 Whitney Mesa Drive, Suite 100
Henderson, Nevada 89014 USA
800.322.3729 USA 702.260.4466 INTL
www.blodermls.com

REF XXXX-0000



YYYY-MM-DD



MDSS GmbH, SCHIFFGRABEN 41
30175 HANNOVER, GERMANY

LOT 00000




YYYY-MM-DD

REV: C

DWG: 6195

1g Sample Tube:

tamper seal



SCARAID SPF

silicone stick with spf stick

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	
Route of Administration		TOPICAL	NDC:84440-619	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)			OCTINOXATE	40 mg in 1 g
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)			AVOBENZONE	30 mg in 1 g
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)			OCTOCRYLENE	50 mg in 1 g
Inactive Ingredients				
Ingredient Name			Strength	
SILICON (UNII: Z4152N8IUI)			880 mg in 1 g	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84440-619-01	1 g in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	08/08/2025	
	NDC:84440-	4.25 g in 1 TUBE; Type 4: Device		

2	NDC:84440-619-42	4.25 g in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	08/08/2025	
3	NDC:84440-619-17	17 g in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	08/08/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M020	08/08/2025	

Labeler - Biodermis, LLC (809650005)

Revised: 8/2025

Biodermis, LLC