

SCARAID US SPF- silicone stick with spf stick
Biodermis, LLC

ScarAid US SPF

DESCRIPTION:

PROSIL SPF15

SILICONE STICK FOR SCARS

GLIDE-ON THERAPY FOR YOUR SCARS

ACTIVE INGREDIENTS:

Avobenzone - 3% Sunscreen

Octocrylene - 5% Sunscreen agent

Octinoxate - 4% Sunscreen agent

DOSAGE & ADMINISTRATION:

Apply 15 minutes before sun exposure. Clean and dry scar site before each application. Apply ScarAid® liberally over the entire scar. May be covered with a bandage. Repeat several times daily for 8-12 weeks or until scar stops responding. To remove, wipe off with a clean cloth or tissue.

PURPOSE:

SILICONE STICK FOR SCARS

BROAD SPECTRUM UVA/UVB SUNSCREEN

HELPS PREVENT SUNBURN

INDICATIONS & USAGE:

SILICONE STICK FOR SCARS

BROAD SPECTRUM UVA/UVB SUNSCREEN

HELPS PREVENT SUNBURN

DIRECTIONS:

Apply 15 minutes before sun exposure. Clean and dry scar site before each application. Apply ScarAid® liberally over the entire scar. May be covered with a bandage. Repeat several times daily for 8-12 weeks or until scar stops responding. To remove, wipe off with a clean cloth or tissue. Store

between 55°F–95°F (13°C–35°C).

WARNINGS:

FOR EXTERNAL USE ONLY

WARNINGS AND PRECAUTIONS:

EXTERNAL USE ONLY

CAUTIONS: Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.

CAUTIONS:

Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.

KEEP OUT OF REACH OF CHILDREN:

KEEP OUT OF REACH OF CHILDREN:

ASK A DOCTOR:

Ask a doctor about using SCARAID SPF with children under 6 months old.

STORAGE:

Store between 55°F–95°F (13°C–35°C).

CONTACT INFORMATION:

MADE IN THE USA U.S. Patent No. 8,557,287

BIODERMIS LLC

www.biodermis.com

1820 Whitney Mesa Drive, Suite 100

Henderson, Nevada 89014 USA

800.322.3729 (USA)

702.260.4466 (INTL)


INACTIVE INGREDIENTS:

Silicone

Primary Package Label

4.25g Tube:

GLIDE-ON THERAPY FOR YOUR SCARS
PRODUCT SEALED FOR YOUR PROTECTION. BREAK SEAL TO OPEN



SCAR AID[®] SILICONE STICK SPF-15
First Aid for Scars


DIRECTIONS: Apply 15 minutes before sun exposure. Clean and dry the scar site before each application. Apply ScarAid[®] liberally over the entire scar. Repeat several times daily for 8-12 weeks or until the scar stops responding. To remove, wipe off with a clean cloth or tissue. Recommended storage conditions are 55°F-95°F (13°C-35°C).



CAUTIONS: Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.




HELPS PREVENT SUNBURN BROAD SPECTRUM UVA/UVB SUNSCREEN

Drug Facts	Active Ingredients	Purpose
4.25g / .15 oz	Avobenzone (3%)	Sunscreen
	Octocrylene (5%)	Sunscreen Agent
	Octinoxate (4%)	Sunscreen Agent

LOT 11241
REF SCSTSP-US

**BIODERMIS**
1820 Whitney Mesa Drive, Suite 100
Henderson, Nevada 89014 USA
800.322.3729 USA 702.260.4466 INTL
www.biodermis.com

MD  2023-02-01
 2028-02-01

 **MADE IN THE USA**   **U.S. Patent No. 8,557,287**
EXTERNAL USE ONLY

17g Tube:

GLIDE-ON THERAPY FOR YOUR SCARS
PRODUCT SEALED FOR YOUR PROTECTION. BREAK SEAL TO OPEN

SCAR AID[®] SILICONE STICK

SPF-15

First Aid for Scars

DIRECTIONS: Apply 15 minutes before sun exposure. Clean and dry the scar site before each application. Apply ScarAid[®] liberally over the entire scar. Repeat several times daily for 8-12 weeks or until the scar stops responding. To remove, wipe off with a clean cloth or tissue. Recommended storage conditions are 55°F-95°F (13°C-35°C).

CAUTIONS: Stop use and ask a doctor if rash occurs. Do not use on damaged or broken skin. For children under 5 year of age, use under adult supervision. Children under 6 months of age: Ask a doctor.

HELPS PREVENT SUNBURN BROAD SPECTRUM UVA/UVB SUNSCREEN

Drug Facts	Active Ingredients	Purpose
17g / .60 oz	Avobenzone (3%)	Sunscreen
	Octocrylene (5%)	Sunscreen Agent
	Octinoxate (4%)	Sunscreen Agent

LOT 11242

REF SCSTSP17-US

 **BIODERMIS**
1820 Whitney Mesa Drive, Suite 100
Henderson, Nevada 89014 USA
800.322.3729 USA 702.260.4466 INTL
www.biodermis.com

MD



2023-02-01




MADE IN THE USA



2028-02-01

1g Sample Tube:

tamper seal



SCARAID US SPF

silicone stick with spf stick

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	
Route of Administration		TOPICAL	NDC:84440-626	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)			OCTINOXATE	40 mg in 1 g
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)			AVOBENZONE	30 mg in 1 g
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)			OCTOCRYLENE	50 mg in 1 g
Inactive Ingredients				
Ingredient Name			Strength	
SILICON (UNII: Z4152N8IUI)			880 mg in 1 g	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84440-626-01	1 g in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	08/08/2025	
	NDC:84440-	4.25 g in 1 TUBE; Type 4: Device		

2	NDC:84440-626-42	4.25 g in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	08/08/2025	
3	NDC:84440-626-17	17 g in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	08/08/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M020	08/08/2025	

Labeler - Biodermis, LLC (809650005)