

**MUCUS RELIEF SEVERE SINUS CONGESTION MAXIMUM STRENGTH- acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated**  
**Walgreen Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Walgreens 44-615**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves:
  - nasal congestion
  - headache
  - sinus congestion and pressure
  - minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily promotes nasal and/or sinus drainage

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 4,000 mg in 24 hours, which is the maximum daily amount
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a

doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- thyroid disease
- heart disease
- diabetes
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not take more than directed**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

***Questions or comments?***

**1-800-426-9391**

***Principal display panel***

**Well at  
Walgreens**

**WALGREENS PHARMACIST RECOMMENDED‡**

NDC 0363-0156-09

**MAXIMUM STRENGTH**

**Mucus Relief  
Severe Sinus Congestion**

**Acetaminophen** 325 mg / Pain Reliever  
Guaifenesin 200 mg / Expectorant  
Phenylephrine HCl 5 mg / Nasal Decongestant

- Relieves headache & sinus congestion • Thins & loosens mucus
- For ages 12 & over

**20 CAPLETS**

Actual Size

Compare to Maximum Strength Mucinex<sup>®</sup> SINUS-MAX<sup>®</sup> Severe Congestion Relief active ingredients<sup>‡‡</sup>

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS  
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

<sup>‡‡</sup> Walgreens Pharmacist Survey Study, November 2016.

<sup>‡‡</sup> This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Maximum Strength Mucinex<sup>®</sup> SINUS-MAX<sup>®</sup> Severe Congestion Relief.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

walgreens.com ©2017 Walgreen Co.

50844 REV0117B61509

0906513F\_R02

**Drug Facts (continued)**

**Inactive ingredients:** corn starch, croscollon, FD&C red #40 aluminum lake, FD&C yellow #5 aluminum lake, magnesium stearate, maledonin, microcrystalline cellulose, polyethylene glycol, polyvinyl

**Drug Facts (continued)**

alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**(Questions or comments? 1-800-426-3391)**

**Drug Facts (continued)**

■ cough that occurs with too much phlegm (mucus)

■ bronchitis, or emphysema

■ persistent or chronic cough such as occurs with smoking, asthma, chronic

■ difficulty in urination due to enlargement of the prostate gland

■ high blood pressure

■ liver disease ■ diabetes

■ heart disease ■ thyroid disease

**Ask a doctor before use if you have**

■ if you have ever had an allergic reaction to this product or any of its ingredients

■ if you have ever had a skin reaction to this product

■ pharmacist before taking this product

■ do not know if your prescription drug contains an MAOI, ask a doctor or

(certain drugs for depression, psychiatric or emotional conditions, or

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

■ acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

■ nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

**Do not use**

■ if a skin reaction occurs, stop use and seek medical help right away.

■ if a rash occurs, stop use and seek medical help right away.

■ if you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

■ with any other drug containing acetaminophen (prescription or

■ nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or

■ do not know if your prescription drug contains an MAOI, ask a doctor or

■ pharmacist before taking this product

■ if you have ever had an allergic reaction to this product or any of its

ingredients

**Warnings**

■ liver damage ■ with other drugs containing acetaminophen

■ may occur if you take ■ more than 4,000 mg in 24 hours, which is the maximum daily amount

■ 3 or more alcoholic drinks every day while using this product

■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms

may include: ■ skin redness ■ blisters ■ rash

■ if a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other drug containing acetaminophen (prescription or

■ nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or

■ do not know if your prescription drug contains an MAOI, ask a doctor or

■ pharmacist before taking this product

■ if you have ever had an allergic reaction to this product or any of its

ingredients

**Drug Facts (continued)**

**Other information**

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR

BULSTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

**Directions**

■ do not take more than 2 caplets in any 24-hour period

■ adults and children 12 years and over: take 2 caplets every 4 hours

■ children under 12 years: do not use

**Keep out of reach of children.** In case of accidental overdose, get medical

help or contact a Poison Control Center (1-800-222-1222) right away.

Prompt medical attention is critical for adults as well as for children even if

you do not notice any signs or symptoms.

**Drug Facts (continued)**

■ cough comes back or occurs with rash or headache that lasts

■ new symptoms occur

■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ nervousness, dizziness, or sleeplessness occur

■ Stop use and ask a doctor if

■ pregnant or breast-feeding, ask a health professional before use.

■ If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or

■ do not know if your prescription drug contains an MAOI, ask a doctor or

■ pharmacist before taking this product

■ if you have ever had an allergic reaction to this product or any of its

ingredients

**Warnings**

■ liver damage ■ with other drugs containing acetaminophen

■ may occur if you take ■ more than 4,000 mg in 24 hours, which is the maximum daily amount

■ 3 or more alcoholic drinks every day while using this product

■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms

may include: ■ skin redness ■ blisters ■ rash

■ if a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other drug containing acetaminophen (prescription or

■ nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or

■ do not know if your prescription drug contains an MAOI, ask a doctor or

■ pharmacist before taking this product

■ if you have ever had an allergic reaction to this product or any of its

ingredients

**Drug Facts (continued)**

**Drug Facts (continued)**

**Active ingredients (in each caplet)**

Acetaminophen 325 mg Pain reliever

Guaifenesin 200 mg Expectorant

Phenylephrine HCl 5 mg Nasal decongestant

**Purpose**

■ temporarily relieves: ■ nasal congestion ■ headache

■ sinus congestion and pressure ■ minor aches and pains

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the

■ bronchial passageways of bothersome mucus and make coughs more

productive ■ temporarily promotes nasal and/or sinus drainage

**Uses**

■ temporarily relieves: ■ nasal congestion ■ headache

■ sinus congestion and pressure ■ minor aches and pains

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the

■ bronchial passageways of bothersome mucus and make coughs more

productive ■ temporarily promotes nasal and/or sinus drainage

**Drug Facts (continued)**

**Other information**

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR

BULSTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

**Directions**

■ do not take more than 2 caplets in any 24-hour period

■ adults and children 12 years and over: take 2 caplets every 4 hours

■ children under 12 years: do not use

**Keep out of reach of children.** In case of accidental overdose, get medical

help or contact a Poison Control Center (1-800-222-1222) right away.

Prompt medical attention is critical for adults as well as for children even if

you do not notice any signs or symptoms.

**Drug Facts (continued)**

■ cough comes back or occurs with rash or headache that lasts

■ new symptoms occur

■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ nervousness, dizziness, or sleeplessness occur

■ Stop use and ask a doctor if

■ pregnant or breast-feeding, ask a health professional before use.

■ If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or

■ do not know if your prescription drug contains an MAOI, ask a doctor or

■ pharmacist before taking this product

■ if you have ever had an allergic reaction to this product or any of its

ingredients

**Warnings**

■ liver damage ■ with other drugs containing acetaminophen

■ may occur if you take ■ more than 4,000 mg in 24 hours, which is the maximum daily amount

■ 3 or more alcoholic drinks every day while using this product

■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms

may include: ■ skin redness ■ blisters ■ rash

■ if a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other drug containing acetaminophen (prescription or

■ nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or

■ do not know if your prescription drug contains an MAOI, ask a doctor or

■ pharmacist before taking this product

■ if you have ever had an allergic reaction to this product or any of its

ingredients

**Drug Facts (continued)**

**Drug Facts (continued)**

**Active ingredients (in each caplet)**

Acetaminophen 325 mg Pain reliever

Guaifenesin 200 mg Expectorant

Phenylephrine HCl 5 mg Nasal decongestant

**Purpose**

■ temporarily relieves: ■ nasal congestion ■ headache

■ sinus congestion and pressure ■ minor aches and pains

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the

■ bronchial passageways of bothersome mucus and make coughs more

productive ■ temporarily promotes nasal and/or sinus drainage

**Uses**

■ temporarily relieves: ■ nasal congestion ■ headache

■ sinus congestion and pressure ■ minor aches and pains

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the

■ bronchial passageways of bothersome mucus and make coughs more

productive ■ temporarily promotes nasal and/or sinus drainage

**Drug Facts (continued)**

Well at  
**Walgreens**

WALGREENS PHARMACIST RECOMMENDED<sup>1</sup>

# MAXIMUM STRENGTH Mucus Relief Severe Sinus Congestion

Acetaminophen 325 mg / Pain Reliever  
Guaifenesin 200 mg / Expectorant  
Phenylephrine HCl 5 mg / Nasal Decongestant

- Relieves headache & sinus congestion • Thins & loosens mucus
- For ages 12 & over

**20** CAPLETS 

Compare to Maximum Strength Mucinex<sup>®</sup> SINUS-MAX<sup>®</sup> Severe Congestion Relief active ingredients<sup>2</sup>

NDC 0363-0158-09

Walgreens 44-615

<b>MUCUS RELIEF SEVERE SINUS CONGESTION MAXIMUM STRENGTH</b>			
acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0158

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

### Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;615
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0156-09	2 in 1 CARTON	06/30/2013	08/28/2021
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/30/2013	08/28/2021

**Labeler** - Walgreen Company (008965063)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		038154464	PACK(0363-0156)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	MANUFACTURE(0363-0156)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	PACK(0363-0156)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		967626305	PACK(0363-0156)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		868734088	PACK(0363-0156)