

ACETAMINOPHEN PM EXTRA STRENGTH- acetaminophen and diphenhydramine hydrochloride tablet, coated
WinCo Foods, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

WNC - 1095 - 2015-0203

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 500 mg	Pain reliever
Diphenhydramine HCl 25 mg	Nighttime sleep aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets at bedtime• do not take more than 2 caplets of this product in 24 hours
children under 12 years	<ul style="list-style-type: none">• do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions?

Or to report an adverse event call 800-824-1706 Monday - Friday 9am - 4pm MST

PRINCIPAL DISPLAY PANEL

WinCo™ Foods

*Compare to the active ingredients in TYLENOL® PM

EXTRA STRENGTH

Acetaminophen PM
 Non-Habit Forming
 ACETAMINOPHEN/ PAIN RELIEVER
 DIPHENHYDRAMINE HCl/ NIGHTTIME SLEEP AID
 50 Caplets

COATING FREE AREA

INK AND COATING FREE FOR LOT AND EXPIRATION STAMPING

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

WinCo FOODS

EXTRA STRENGTH Acetaminophen PM

ACETAMINOPHEN/
PAIN RELIEVER
DIPHENHYDRAMINE HCl/
NIGHTTIME SLEEP AID

Non-Habit Forming

50 Caplets

*Compare to the active ingredients in **TYLENOL® PM**

Drug Facts (continued)

- do not drive a motor vehicle or operate machinery
- **Stop use and ask a doctor if**
 - sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
- These could be signs of a serious condition.
- If **pregnant or breast-feeding**, ask a health professional before use.
- **Keep out of reach of children.**
- **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

- adults and children 12 years and over
 - take 2 caplets at bedtime
 - do not take more than 2 caplets of this product in 24 hours
- children under 12 years
 - do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

Other information

- store between 20-25°C (68-77°F) in a dry place. ▼

Drug Facts

Active ingredients (in each caplet) Purpose
 Acetaminophen 500 mg.....Pain reliever
 Diphenhydramine HCl 25 mg.....Nighttime sleep aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ with any other product containing diphenhydramine, even one used on skin
 ■ in children under 12 years of age
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
 ■ liver disease
 ■ glaucoma
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin
 ■ taking sedatives or tranquilizers

When using this product
 ■ drowsiness will occur
 ■ avoid alcoholic drinks

50 Caplets

*Compare to the active ingredients in **TYLENOL® PM**

Non-Habit Forming

50 Caplets

*Compare to the active ingredients in **TYLENOL® PM**

Drug Facts (continued)

- retain carton for complete product information

Inactive ingredients colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions? Or to report an adverse event call 800-824-1706 Monday - Friday 9am - 4pm MST

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704
 *This product is not manufactured or distributed by MCNEIL CONSUMER HEALTHCARE, distributor of **TYLENOL® PM**.

100% SATISFACTION GUARANTEE
 Comments & Questions? 800-824-1706

ACETAMINOPHEN PM EXTRA STRENGTH

acetaminophen and diphenhydramine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-137
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
COPOVIDONE K25-31 (UNII: D9C330MD8B)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1031
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-137-50	1 in 1 CARTON	12/15/2014	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/15/2014	

Labeler - WinCo Foods, LLC (056098817)

Revised: 12/2018

WinCo Foods, LLC