CLOTRIMAZOLE- clotrimazole cream QUALITY CHOICE (Chain Drug Marketing Association)

Clotrimazole Vaginal Cream, USP

Drug Facts

Active ingredient

Clotrimazole, USP 1% (50 mg in each applicatorful)

Clotrimazole, USP 1% (external cream)

Purpose

Vaginal antifungal

Vaginal antifungal

Uses

- treats vaginal yeast infections
- relieves external itching and irritation due to a vaginal yeast infection

Warnings

For external use only.

Do not use

if you have never had a vaginal yeast infection diagnosed by a doctor.

Ask a doctor before use if you have

- vaginal itching and discomfort for the first time
- lower abdominal, back or shoulder pain, fever, chills nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

When using this product,

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur

• if you do not get complete relief, ask a doctor before using another product

Stop use and ask a doctor if

- symptoms do not get better in 3 days
- symptoms last more than 7 days
- you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- before using this product, read the enclosed educational brochure for complete directions and information
- adults and children 12 years of age and over:
 - **vaginal cream**: insert one applicatorful of cream into the vagina at bedtime for 7 days in a row. Wash applicator after each use.
 - external cream: use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.
- children under 12 years of age: ask a doctor

Other information

- to open tube: unscrew cap, lift tab, and pull to remove foil seal prior to use
- do not use if foil seal on tube opening is broken or missing
- do not purchase if carton is open
- store at room temperature 20º to 25ºC (68º to 77ºF)
- before using any medication, read all label directions. Keep carton, it contain important information.

Inactive ingredients

benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polysorbate 60, purified water, sodium phosphate monobasic, sorbitan monostearate

Questions or comments?

Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

Principal display panel

7-Day Treatment

Clotrimazole

Vaginal Cream, USP 1%

VAGINAL ANTIFUNGAL

Clotrimazole Vaginal Cream Can Kill The Yeast That May Cause Vaginal Infection Cures Most Vaginal Yeast Infections

g (OZ) Tube

Tube of Vaginal Cream (1%) and 1 Reusable Applicator

Distributed by: CDMA, Inc.

Novi, MI 48375

Package label

foul-smelling vaginal discharge

- if you do not get complete relief, ask a doctor before using another product
- do not have vaginal intercourse mild increase in vaginal burning, fiching or imtation may occur (S115) S26823610

Condoms and diaphragms may be damaged and fail to prevent pragnancy or sexually transmitted When using this product a do not use tempons, douches, spermicides, or other vaginal products.

- peeu exboseq to the furmen immunodeficiency virus (HW) that causes AIDS
- a serious underlying medical cause for your symptoms, including diabetes or a weakened immune ■ vaginal yeast infections often (such as once a morth or 3 in 6 months). You could be pregnant, or have
- vaginal litching and discomfort for the first time
 lower abdominal, back or shoulder pain, tower, chills, nausea, romiting, or foul-smelling vaginal discharge. You may have a more serious condition. Ask a doctor before use if you have

Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.

For vaginal use only.

■ referves external fiching and imfation due to a vaginal yeast infection Sees Interest length altern Read infections

Continuazole, USP 1% (50 mg in each applicatoriu). Clofrimazole, USP 1% (external cream)...... ječunitnie jeuičev Legnuntine Leri gaV. esoding Active ingredients

Drug Facts

CINESTRONS? CAI 1-800-836-2362 Monday-Friday 9AM-5PM EST

Inactive ingredients beays alcohol, estaclasaryl alcohol, ostyl estara wax, 2-octyldodecanol, polycorbale 60, purified water, sodium phosphate monobasic, aorbitan monostaerate

- before using any medication, read all label directions. Keep carbon, it contains important information. ■ store at morn temperature 20° to 25°C (68° to 77°F)
 - do not purchase if carton is open
 - do not use if foil seal on tube opening is broken or missing to ober tribe: unscrew csp, lift tab, and pull to remove toll seal prior to use

подътот і потыва

children under 12 years of age: ask a doctor


vagine. Use 2 times daily for up to 7 days as needed.

- vagina. Squeeze a small amount of cream onto your fingertlp. Apply to itshy, initiated skin outside the m external creams use the same tube of cream if you have litting and initiation on the skin outside the applicator after each use.
- waginal cream: Insert one applicational of cream into the vagina at bedtime for 7 days in a row. Wash ■ sqrigs and children 12 years of age and over:
 - DIVECTIONS before using this product, read the enciceed educational brochure for complete

(1-800-222-1222) right away.

if pregnant or breast-feeding, sak a health professional below use. Keep out of reach of children, if swallowed, get medical help or contact a Poison Control Center

Dung Facts (continued)



NDC 83324-154-15



Educational Pamphlet Enclosed

7-Day Treatment

otrimazole

Vaginal Cream USP, 1%

Vaginal Antifungal

Clotrimazole Vaginal Cream Can Kill The Yeast That May Cause Vaginal Infection **Cures Most Vaginal Yeast Infections**

One ${f 45}$ g (1.59 OZ) Tube of Vaginal Cream (1%) and 1 Reusable Applicator (7-Day Therapy)





P.D-07138 F0009146

Code: GO/DRUGS/361 AAD3E/04

QUALITY CHOICE Vaginal Antifungal

CLOTRIMAZOLE

clotrimazole cream

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83324-154	
Route of Administration	VAGINAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CLOTRIMAZOLE (UNII: G07GZ97H65) (CLOTRIMAZOLE - UNII:G07GZ97H65)	CLOTRIMAZOLE	10 mg in 1 g	

Inactive Ingredients		
Ingredient Name	Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)		
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)		
CETYL ESTERS WAX (UNII: D072FFP9GU)		
OCTYLDODECANOL (UNII: 461N1O614Y)		
POLYSORBATE 60 (UNII: CAL22UVI4M)		
WATER (UNII: 059QF0KO0R)		
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)		
SORBITAN MONOSTEARATE (UNII: NVZ4I0H58X)		

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
NDC:83324-154- 15	1 in 1 CARTON	07/31/2024			
1	45 g in 1 TUBE; Type 0: Not a Combination Product				

Marketing Information			
Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date

ANDA ANDA074165 07/31/2024

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2024 QUALITY CHOICE (Chain Drug Marketing Association)