# LORATADINE- loratadine tablet Fred's Inc.

-----

**Drug Facts** 

### **ACTIVE INGREDIENT (IN EACH TABLET)**

Loratadine USP, 10 mg

#### **PURPOSE**

Antihistamine

#### **USES**

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

#### **WARNINGS**

#### Do not use

If you have ever had an allergic reaction to this product or any of its ingredients.

#### Ask a doctor before use if you have

Liver or kidney disease. Your doctor should determine if you need a different dose.

#### When using this product

Do not take more than directed. Taking more than directed may cause drowsiness.

#### Stop use and ask a doctor if

An allergic reaction to this product occurs. Seek medical help right away.

#### If pregnant or breast-feeding

Ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **DIRECTIONS**

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours			
children under 6 years of age	ask a doctor			
consumers with liver or kidney disease ask a doctor				

#### OTHER INFORMATION

- store between 20 and 25° C (68 and 77° F)
- protect from excessive moisture
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.

#### INACTIVE INGREDIENTS

Corn starch, lactose monohydrate, magnesium stearate, pregelatinized starch

### **QUESTIONS?**

Call 1-800-406-7984

#### PRINCIPAL DISPLAY PANEL

fred's®

Non-Drowsy\*

\* When taken as directed.

See Drug Facts Panel.

24 Hour

**Allergy Relief** 

Loratadine Tablets USP, 10 mg

**Antihis tamine** 

#### **INDOOR & OUTDOOR ALLERGIES**

#### Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat and Nose

#### 10 TABLETS

Compare to Active Ingredient of Claritin  $^{\mathbb{R}^{\dagger}}$ 

■ store between 20 and 25° C (68 and 77° F) ■ protect from excessive moisture Nother information

consumers with liver or kidney disease	sak a doctor
children under 6 years of age	ask a dodor
adults and children 6 years and over	tablet daily; not more than 1 tablet in 24 hours
SHOUTS HA	

away.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right If pregnant or breast-feeding, ask a health professional before use.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

Muren using this product do not take more than directed. Taking more than directed may cause drowsiness.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ Itchy, watery eyes ■ sneezing ■ Itching of the nose or throat esou /kuun.i

Loratadine USP, 10 mg. Antinistamine Active ingredient (in each tablet) əsodınd

Drug Facts



Non-Drowsy\*

# 24 Hour Allergy Relief

Loratadine Tablets USP, 10 mg



## Non-Drowsy\*

When taken as directed. See Drug Facts Panel.



Shown at Actual Size

'All trademarks are property of their respective Owners. This product is not affiliated with the makers/owners of Claritin.

Non Varnish Area

# 24 Hour Allergy Relief

# Loratadine Tablets USP, 10 mg **Antihistamine**

INDOOR & OUTDOOR ALLERGIES

Relief of:

- Runny Nose Sneezing
- Itchy, Watery Eyes
   Itchy Throat and Nose

10 TABLETS



See end panel for expiration date. Keep the carton. It contains important information.

Questions? call 1-800-406-7984

Inactive ingredients com starch, lactose monohydrate, magnesium stearate, pregelatinized starch







## LORATADINE

loratadine tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55315-528	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg	

Inactive Ingredients		
Ingredient Name	Strength	
STARCH, CORN (UNII: O8232NY3SJ)		
LACTO SE MO NO HYDRATE (UNII: EWQ57Q8I5X)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		

Product Characteristics				
Color	white (White to Off-White)	Score	no score	
Shape	ROUND	Size	6 mm	
Flavor		Imprint Code	RX526	
Contains				

	Packaging				
ı	#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
ı	1	NDC:55315-528-69	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	08/19/2003	
	2	NDC:55315-528-31	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	08/19/2003	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076134	08/19/2003	

## **Labeler** - Fred's Inc. (005866116)

Registrant - Sun Pharmaceutical Industries Inc. (146974886)

Establishment				
Name	Address	ID/FEI	Business Operations	
Ohm Laboratories Inc.		051565745	manufacture(55315-528)	

Revised: 12/2019 Fred's Inc.