

MAX RELIEF JUNIOR GRAPE- children acetaminophen suspension
Akron Pharma Inc.

MAX relief junior
Children's Pain Reliever
Acetaminophen Oral Suspension
Grape Flavor
160mg per 5mL

Dye-Free

Active Ingredient

Active ingredient (in each 5mL)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer

Uses

temporarily: reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:

if sore throat is severe, persist or more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).

- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor or pharmacist before use if the child has

liver disease.

Ask doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. **When using this product do not exceed recommended dose (see overdose warning)**

Stop use and ask a doctor if

- pain gets worse or lasts for more than 5 days
- fever gets worse or last for more than 3 days
- new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Overdose warning:

In case of overdose, get medical help to contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Keep out of Reach of children

Directions

- this product does not contain directions or complete warnings for adult use
- shake well before using
- mL = milliliter
- find right dose on chart below. if possible, use weight to dose, otherwise use age
- use only the enclosed dosing cup designed for use with product.

Weight (lbs)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
25 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	2 1/2 tsp or 12.5 mL

Other Information

- Store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

Inactive Ingredients

Anhydrous Citric Acid, D&C Red #33, FD&C Blue #1, Grape Flavor, Glycerin, Microcrystalline cellulose and Sodium Methyl Cellulose, Potassium Sorbate, Purified Water, Sorbitol, Sucralose, Sucrose, Xanthan gum.

Questions or comments?

call toll-free 1-877-225-6999

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com

NDC 71399-0048-2



Children's
PAIN RELIEVER
and fever Reducer

PAIN + FEVER Acetaminophen
Oral Suspension

160mg per 5mL




Akron Pharma

2 FL.OZ (60 mL)

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Acetaminophen 160 mg	Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warning
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions.
 Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
 ■ this product does not contain directions or complete warnings for adult use, shake well before using
 ■ mL = milliliter
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 ■ use only the enclosed dropper designed for use with product.

Drug Facts (continued)

Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms lasts ■ do not give more than 5 times in 24 hours.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

Other Information
 store at room temperature 20°-25°C (69°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, D&C Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

Tamper Evident: Do Not Use If Printed Safety Seal Around Bottle or Under Cap is Broken or Missing

LOT No.:
EXPDate.:

Akron Pharma
 Manufactured for:
 Akron Pharma, Inc.
 Fairfield, NJ 07004
 www.akronpharma.com

REU0425



N 3 71399 00482 6

PAIN + FEVER

Acetaminophen

Oral Suspension 160mg per 5mL

MAXrelief[™] junior

NDC 71399-0048-2

PAIN + FEVER

Acetaminophen

Oral Suspension 160mg per 5mL

Children's
PAIN RELIEVER
and fever Reducer

PAIN + FEVER
Acetaminophen
Oral Suspension
160mg per 5mL



Akron Pharma

2 FL.OZ (60 mL)



Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily relieves fever relieves minor aches and pains due to: the common cold flu a headache a sore throat a toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

more than 5 doses in 24 hours, which is the maximum daily amount

with other drugs containing acetaminophen.

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include a skin redness or blisters or rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if: pain gets worse or lasts for more than 5 days or fever gets worse or lasts for more than 3 days.

new symptoms occur redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

Use as directed by a doctor.

This product does not contain directions or complete warnings for adult use. Shake well before using.

mL = milliliter

Find right dose on chart below. If possible, use weight to dose; otherwise, use age.

Use only the enclosed dropper designed for use with product.

Do not use any other dosing device.

If needed, repeat dose every 4 hours while symptoms last. Do not give more than 5 times in 24 hours.

NDC 71399-0048-2

PAIN + FEVER

Acetaminophen

Oral Suspension 160mg per 5mL

Children's
PAIN RELIEVER
and fever Reducer

PAIN + FEVER
Acetaminophen
Oral Suspension
160mg per 5mL



Akron Pharma

2 FL.OZ (60 mL)



Drug Facts (continued)

Do not give more than 5 days unless directed by doctor (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 85	11	3 tsp or 15 mL

*as directed by a doctor

Other Information

Store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

Inactive ingredients: anhydrous citric acid, D&C Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol[®] Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

SAVE THIS CARTON FOR FULL LABELING INFORMATION.

Use only with enclosed dropper

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com



NDC 71399-0048-4

PAIN + FEVER

Acetaminophen

Oral Suspension 160mg per 5mL

MAXrelief[™] junior

Children's
PAIN RELIEVER
and fever Reducer

PAIN + FEVER
Acetaminophen
Oral Suspension
160mg per 5mL



Akron Pharma

4 FL.OZ (118 mL)



Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily relieves fever relieves minor aches and pains due to: the common cold flu headache sore throat toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters or rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if: pain gets worse or lasts for more than 5 days or fever gets worse or lasts for more than 3 days. new symptoms occur redness or swelling is present. These could be signs of a serious condition. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions this product does not contain directions or complete warnings for adult use do not give more than directed (see overdose warning) shake well before using mL = milliliter find right dose on chart below. If possible, use weight to dose; otherwise, use age. remove the child protective cap and squeeze your child's dose into the dosing cup repeat dose every 4 hours while symptoms last do not give more than 5 times in 24 hours.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*	Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
24 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

*as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other Information store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light. **Inactive ingredients:** anhydrous citric acid, D&C Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol[®] Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

LOT No.:
EXPIRES:
Un Warnish Area

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com



LBH: 70x45x115mm

15mm

60mm

MAXrelief junior

Children's PAIN RELIEVER

and fever Reducer

Acetaminophen Oral Suspension

160mg per 5mL

NDC 71399-0048-4

MAXrelief junior

Children's PAIN RELIEVER and fever Reducer

Acetaminophen Oral Suspension

160mg per 5mL

PAIN+FEVER

Grape flavor

Akron Pharma

4 FL.OZ (118 mL)



Drug Facts

Active ingredient (in each 5 mL) Purpose

Acetaminophen 160 mg Pain reliever, fever reducer

Uses temporarily relieves fever relieves minor aches and pains due to: common cold flu headache sore throat toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen.

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, skin itching, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if pain gets worse or lasts for more than 5 days fever gets worse or lasts for more than 3 days new symptoms occur redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

This product does not contain directions or complete warnings for adult use

Do not give more than directed (see overdose warning)

Shake well before using mL = milliliter

Find right dose on chart below. If possible, use weight to dose; otherwise, use age.

Remove the child protective cap and squeeze your child's dose into the dosing cup

Repeat dose every 4 hours while symptoms last

NDC 71399-0048-4

MAXrelief junior

Children's PAIN RELIEVER and fever Reducer

Acetaminophen Oral Suspension

160mg per 5mL

PAIN+FEVER

Grape flavor

Akron Pharma

4 FL.OZ (118 mL)



Drug Facts (continued)

Do not give more than 5 times in 24 hours.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

Store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive ingredients: anhydrous citric acid, D&C Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxymethyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by Mohel Consumer Healthcare, distributor of Infant Tylenol Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Akron Pharma

Manufactured for: Akron Pharma, Inc. Fairfield, NJ 07004 www.akronpharma.com

REV 07/25



Size : 50x130x50mm Lock Bottom

NDC 71399-0048-8

MAXrelief™ junior

Children's
PAIN RELIEVER
and fever Reducer

Acetaminophen
Oral Suspension

160mg per 5mL

PAIN + FEVER

Grape
flavor



Akron Pharma

8 FL.OZ (236 mL)

Drug Facts

Active ingredient (in each 5 mL) Purpose

Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use. ■ do not give more than directed (see overdose warning) ■ shake well before using ■ mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last

Drug Facts (continued)

■ do not give more than 5 times in 24 hours.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL) *
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other Information


store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, D&C Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



Akron Pharma

Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004

www.akronpharma.com


REV.07/25

N 3



15mm

58mm



58mm

58mm

58mm

NDC 71399-0048-8

MAX relief junior

Children's PAIN RELIEVER and fever Reducer

Acetaminophen Oral Suspension

160mg per 5mL

PAIN + FEVER

Grape flavor



Alkermes Pharma

8 FL.OZ (236 mL)

Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: reduces fever, relieves minor aches and pains due to: common cold, flu, headache, sore throat, toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: more than 5 doses in 24 hours, which is the maximum daily amount; with other drugs containing acetaminophen.

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if pain gets worse or lasts for more than 5 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- if needed, repeat dose every 4 hours while symptoms lasts

NDC 71399-0048-8

MAX relief junior

Children's PAIN RELIEVER and fever Reducer

Acetaminophen Oral Suspension

160mg per 5mL

PAIN + FEVER

Grape flavor



Alkermes Pharma

8 FL.OZ (236 mL)

Drug Facts (continued)

- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by doctor
- do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other Information

store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

Inactive ingredients anhydrous citric acid, D&G Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999. *This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infant Tylenol Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

REV07025



Alkermes Pharma
Manufactured for:
Alkermes Pharma, Inc.
Fairfield, NJ 07004
www.alkermes.com

N 3

71399 00488 8

Size : 58x160x58mm Lock Bottom

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
 Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to:
 ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warning
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
 ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
When using this product, do not exceed recommended dose (see overdose warning)
Stop use and ask a doctor if
 ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.
Directions
 ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed (see overdose warning)
 ■ shake well before using. ■ mL=milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

NDC 71399-0048-6

MAXrelief™ junior

Children's
PAIN RELIEVER
 and fever Reducer

PAIN+FEVER
Acetaminophen
Oral Suspension
160mg per 5mL



Grape
 flavor

16 FL. OZ (473 mL)

Drug Facts (continued)

- if needed, repeat dose every 4 hours while symptoms lasts
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by doctor.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other Information
 store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, D&C Red #33, FD&C Blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Akron Pharma
 Manufactured for:
 Akron Pharma, Inc.
 Fairfield, NJ 07004
 www.akronpharma.com

REV.07/25

MAX RELIEF JUNIOR GRAPE

children acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71399-0048
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
POTASSIUM SORBATE (UNII: 1VPU26JZ4)	
WATER (UNII: 059QF0KO0R)	

SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399-0048-2	1 in 1 CARTON	04/24/2024	
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:71399-0048-4	1 in 1 CARTON	04/24/2024	
2		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:71399-0048-8	1 in 1 CARTON	04/24/2024	
3		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:71399-0048-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/24/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/24/2024	

Labeler - Akron Pharma Inc. (067878881)