

IBUPROFEN - ibuprofen tablet
MARKSANS PHARMA LIMITED

Ibuprofen Tablets 200mg (OTC; Brown Tablets & Caplets)

Active ingredient(s)

Ibuprofen 200 mg (NSAID)*

* nonstreoidal anti-inflammatory drug

Purpose

Pain reliever / fever reducer

Use(s)

- temporarily relieves minor aches and pains due to :
- backache
- headache
- menstrual cramps
- minor pain of arthritis
- muscular aches
- the common cold
- toothache
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergy reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have bad stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- the more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- You have problems or serious side effects from taking pain relievers or fever reducers
- The stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn
- You have a high blood pressure, heart disease, cirrhosis, kidney disease, asthma or had a stroke
- You are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present in painful area
- any new symptoms appear

Pregnancy/Breastfeeding

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause a problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

- store between 20-25 °C (68-77 °F).
- **do not use if seal under bottle cap imprinted with” SEALED for YOUR PROTECTION” is broken or missing.**

Inactive ingredients

Silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or Comments?

Call 1-877-290-4008

Manufactured for:
Time-Cap Labs, Inc.
 7 Michael Avenue,

Farmingdale,
NY 11735, USA


Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83
Verna Industrial Estate
Verna, Goa-403722, India






Storage

Principal Display Panel

Ibuprofen Tablets USP, 200mg

30 counts
Bottle Label


<p>NDC 25000-114-03</p> <p>Ibuprofen Tablets USP, 200 mg</p> <p>Pain Reliever / Fever Reducer (NSAID)</p> <p>30 Coated Tablets</p> <p> Marksans</p>		<p>Important: Read all product information before using. Keep the box for important information.</p> <p>Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Purpose Pain reliever/fever reducer</p> <p>Uses ■ temporarily relieves minor aches and pains due to: ■ backache ■ headache ■ menstrual cramps ■ minor pain of</p>	<p>LOT : EXP : R804/17</p>
<p>arthritis ■ muscular aches ■ the common cold ■ toothache ■ temporarily reduces fever.</p> <p>Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.</p>	<p>Symptoms may include: ■ asthma (wheezing) ■ blisters ■ facial swelling ■ hives ■ rash ■ shock ■ skin reddening. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product</p>	<p>contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug</p>	
<p>■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than</p>	<p>directed. Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use</p>	<p>if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel</p>	
<p>faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech</p>	<p>■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask</p>	<p>a health professional before use. Keep out of reach of children. Directions : ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older : ■ take 1 tablet every 4 to 6</p>	
<p>hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor. Other information ■ store</p>	<p>between 20-25°C (68-77°F)</p> <p>Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.</p> <p>Mfg. Lic. No. : Go/Drugs/515</p>	<p>Manufactured for: Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735, USA</p> <p>Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, GOA - 403 722, India.</p>	

<p>Description : NDC 25000-114-03 Ibuprofen 200 mg 30-Tablets Label_Open</p>		<p>Labeling Format Information :</p>	
<p>File # : NDC 25000-114-03</p>	<p>Color scheme :</p> <p> PANTONE 484 C</p> <p> PANTONE 300 C</p> <p> PANTONE 273 C</p> <p> PANTONE 7406 C</p>	<p>Fonts : Wnd. SW Condensed, Bold, Bold Italic.</p>	
<p>Version : R804/17</p>	<p>Size :</p> <p>140.00(L) X 80.00(H) mm</p>	<p>Drug Facts : 8.25 pt.</p> <p>Header : 7.00 pt.</p> <p>Subheader : 6.00 pt.</p> <p>Body Text : 6.00 pt.</p> <p>Drug Facts (Continued) : 7.00 pt.</p>	<p>Leading : 6.00 pt.</p> <p>Header : 5.00 pt.</p> <p>Barlines : 2.50 pt.</p> <p>Hairlines : 0.50 pt.</p>
<p>Date : 5/04/2017</p>	<p></p>	<p>Horizontal Scale : 70%</p>	<p>Average Kerning : 0</p>

Ibuprofen Tablets USP, 200mg

30 counts

Carton Label



Drug Facts (Continued)

NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if

■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ the stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
 ■ you are taking a diuretic.

Ask a doctor or pharmacist before use if you are

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ under a doctor's care for any serious condition
 ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Drug Facts (Continued)

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
 ■ you have symptoms of heart problem or stroke:
 ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

■ do not take more than directed
 ■ the smallest effective dose should be used

adults and children 12 years and older ■ take 1 tablet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
 ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years ■ ask a doctor

NDC 25000-114-03

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

30 Coated Tablets



Important : Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)*
Purpose Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

■ temporarily relieves minor aches and pains due to:
 ■ backache ■ headache ■ menstrual cramps
 ■ minor pain of arthritis ■ muscular aches
 ■ the common cold ■ toothache
 ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:
 ■ asthma (wheezing) ■ blisters ■ facial swelling
 ■ hives ■ rash ■ shock ■ skin reddening
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription.

Drug Facts (Continued)

Other Information

■ store between 20-25°C (68-77°F).
 ■ do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing
 ■ see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for:
Time-Cap Labs, Inc.
 7 Michael Avenue,
 Farmingdale,
 NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
 Plot No. L-82, L-83, Verna Indl. Estate,
 Verna, GOA - 403 722, India.
 Mfg. Lic. No. : 60/Drugs/515

R804/17

Description : NDC 25000-114-03 Ibuprofen 200 mg Tablets Carton_Open

File # : NDC 25000-114-03

Version : R804/17

Date : 5/04/2017



Color scheme :

- PANTONE 484 C
- PANTONE 300 C
- PANTONE 273 C
- PANTONE 7406 C

Size :
 60.00(L) X 90.00(H) X 60.00(W) mm

Labeling Format Information :

Fonts : Wnd. SW Condensed, Bold, Bold Italic.


Drug Facts :	8.25 pt.	Leading :	6.00 pt.
Header :	7.00 pt.	Header :	5.00 pt.
Subheader :	6.00 pt.	Barlines :	2.50 pt.
Body Text :	6.00 pt.	Hairlines :	0.50 pt.
Drug Facts (Continued) :	7.00 pt.		






Horizontal Scale : 70%

Average Kerning : 0

Ibuprofen Tablets USP, 200mg (Caplets)

30 counts
Bottle Label


<p>NDC 25000-117-03</p> <p>Ibuprofen Tablets USP, 200 mg</p> <p>Pain Reliever / Fever Reducer (NSAID)</p> <p>30 Coated Caplets* * Capsule-Shaped Tablets</p> <p> Marksans</p>		<p>Important: Read all product information before using. Keep the box for important information.</p> <p>Active ingredient (in each caplet) Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Purpose Pain reliever/fever reducer</p> <p>Uses ■ temporarily relieves minor aches and pains due to: ■ backache ■ headache ■ menstrual cramps ■ minor pain of</p> <p>LOT : EXP : R804/17</p>
<p>arthritis ■ muscular aches ■ the common cold ■ toothache ■ temporarily reduces fever.</p> <p>Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.</p>	<p>Symptoms may include: ■ asthma (wheezing) ■ blisters ■ facial swelling ■ hives ■ rash ■ shock ■ skin reddening. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product</p>	<p>contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug</p>
<p>■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than</p>	<p>directed. Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use</p>	<p>if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel</p>
<p>faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech</p>	<p>■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask</p>	<p>a health professional before use. Keep out of reach of children. Directions : ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older : ■ take 1 caplet every 4 to 6</p>
<p>hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor. Other information ■ store</p>	<p>between 20-25°C (68-77°F)</p> <p>Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.</p> <p>Mfg. Lic. No. : Go/Drugs/515</p>	<p>Manufactured for: Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735, USA</p> <p>Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, GOA - 403 722, India.</p>


<p>Description : NDC 25000-117-03 Ibuprofen 200 mg 30-Caplets Label_Open</p>		<p>Labeling Format Information :</p>	
<p>File # : NDC 25000-117-03</p>	<p>Color scheme :</p> <p> PANTONE 484 C</p> <p> PANTONE 300 C</p> <p> PANTONE 273 C</p> <p> PANTONE 7406 C</p>	<p>Fonts : Wnd. SW Condensed, Bold, Bold Italic.</p>	
<p>Version : R804/17</p>	<p>Size : 140.00(L) X 80.00(H) mm</p>	<p>Drug Facts : 8.25 pt.</p> <p>Header : 7.00 pt.</p> <p>Subheader : 6.00 pt.</p> <p>Body Text : 6.00 pt.</p> <p>Drug Facts (Continued) : 7.00 pt.</p>	<p>Leading : 6.00 pt.</p> <p>Header : 5.00 pt.</p> <p>Barlines : 2.50 pt.</p> <p>Hairlines : 0.50 pt.</p>
<p>Date : 5/04/2017</p>	<p></p>	<p>Horizontal Scale : 70% Average Kerning : 0</p>	

Ibuprofen Tablets USP, 200mg (Caplets)

30 counts

Carton Label



Description : NDC 25000-117-03 Ibuprofen 200 mg Caplets Carton_Open		Labeling Format Information :	
File # : NDC 25000-117-03	Color scheme :	Fonts : Wnd. SW Condensed, Bold, Bold Italic.	
Version : R804/17	<ul style="list-style-type: none"> PANTONE 484 C PANTONE 300 C PANTONE 273 C PANTONE 7406 C 	Drug Facts : 8.25 pt.	Leading : 6.00 pt.
Date : 5/04/2017	Size : 60.00(L) X 90.00(H) X 60.00(W) mm	Header : 7.00 pt.	Header : 5.00 pt.
		Subheader : 6.00 pt.	Barlines : 2.50 pt.
		Body Text : 6.00 pt.	Hairlines : 0.50 pt.
		Drug Facts (Continued) : 7.00 pt.	
		Horizontal Scale : 70%	Average Kerning : 0

Ibuprofen Tablets USP, 200mg

500 counts
Bottle Label

NDC 25000-114-70

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

500 Coated Tablets
Marksans



Drug Facts

Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses


- temporarily relieves minor aches and pains due to:
 - backache ■ headache ■ menstrual cramps
 - minor pain of arthritis ■ muscular aches
 - the common cold ■ toothache
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ asthma (wheezing) ■ hives ■ rash ■ shock ■ skin reddening

Purpose

LOT :
EXP :
R011/19



Drug Facts (Continued)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke.

Drug Facts (Continued)

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Drug Facts (Continued)

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling

Drug Facts (Continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (Continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Drug Facts (Continued)

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for: **Time-Cap Labs, Inc.**
7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by: **Marksans Pharma Ltd.**
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/Drugs/515

Description : Ibuprofen 200mg 500 tablets brown label

File # : NDC 25000-114-70

Version : R011/19

Date : 30/11/2019



Color scheme :

■ PANTONE 484 C

Size :

165.00(L) X 57.00(H) mm

Labeling Format Information :

Fonts : Wnd. SW Condensed, Bold, Bold Italic.

Drug Facts :	8.25 pt.	Leading :	6.00 pt.
Header :	7.00 pt.	Header :	5.00 pt.
Subheader :	6.00 pt.	Barlines :	2.50 pt.
Body Text :	6.00 pt.	Hairlines :	0.50 pt.
Drug Facts (Continued) :	7.00 pt.		

Horizontal Scale : 70%

Average Kerning : 0

Ibuprofen Tablets USP, 200mg

1000 counts
Bottle Label

NDC 25000-114-71

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

**THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN**

1000 Coated Tablets



Marksans

Drug Facts

Active ingredient (in each tablet) *Purpose*
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - backache ■ headache ■ menstrual cramps
 - minor pain of arthritis ■ muscular aches
 - the common cold ■ toothache
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing) ■ blisters ■ facial swelling
- hives ■ rash ■ shock ■ skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen).

Drug Facts (Continued)

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are



Drug Facts (Continued)

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body
 - slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (Continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Drug Facts (Continued)

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue,
Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/Drugs/515

Description : Ibuprofen 200mg 1000 tablets brown label

File # : NDC 25000-114-71

Version : R011/19

Date : 30/11/2019



Color scheme :

■ PANTONE 484 C

Size :

229.00(L) X 86.00(H) mm

Labeling Format Information :

Fonts : Wnd. SW Condensed, Bold, Bold Italic.

Drug Facts :	8.25 pt.	Leading :	6.00 pt.
Header :	7.00 pt.	Header :	5.00 pt.
Subheader :	6.00 pt.	Barlines :	2.50 pt.
Body Text :	6.00 pt.	Hairlines :	0.50 pt.
Drug Facts (Continued) :	7.00 pt.		

Horizontal Scale : 70%

Average Kerning : 0

Ibuprofen Tablets USP, 200mg (Caplets)

300 counts
Bottle Label

NDC 25000-117-69

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)
**THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN**

300 Coated Caplets*
*Capsule-Shaped Tablets
Marksans



Drug Facts

Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - headache
 - menstrual cramps
 - minor pain of arthritis
 - muscular aches
 - the common cold
 - toothache
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

LOT :
EXP :
R011/19



Drug Facts (Continued)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke.

Drug Facts (Continued)

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Drug Facts (Continued)

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

Drug Facts (Continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (Continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Drug Facts (Continued)

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for: **Time-Cap Labs, Inc.**
7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by: **Marksans Pharma Ltd.**
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No.: GO/Drugs/515

Description : Ibuprofen 200mg caplet 300's brown label

File # : NDC 25000-117-69

Version : R011/19

Date : 30/11/2019



Color scheme :

PANTONE 484 C

Size :

165.00(L) X 57.00(H) mm

Labeling Format Information :

Fonts : Wnd. SW Condensed, Bold, Bold Italic.

Drug Facts :	8.25 pt.	Leading :	6.00 pt.
Header :	7.00 pt.	Header :	5.00 pt.
Subheader :	6.00 pt.	Barlines :	2.50 pt.
Body Text :	6.00 pt.	Hairlines :	0.50 pt.
Drug Facts (Continued) :	7.00 pt.		

Horizontal Scale : 70%

Average Kerning : 0

IBUPROFEN

ibuprofen tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:25000-114

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	114
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-114-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
2	NDC:25000-114-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
3	NDC:25000-114-20	4 in 1 BOX	07/25/2011	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-114-30	6 in 1 BOX	07/25/2011	
4		6500 in 1 BAG; Type 0: Not a Combination Product		
5	NDC:25000-114-98	1 in 1 BOX	07/25/2011	
5		34065 in 1 BAG; Type 0: Not a Combination Product		
6	NDC:25000-114-93	1 in 1 BOX	07/25/2011	
6		37470 in 1 BAG; Type 0: Not a Combination Product		
7	NDC:25000-114-70	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
8	NDC:25000-114-71	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	07/25/2011	

IBUPROFEN

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-117
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	BROWN	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	117
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-117-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
2	NDC:25000-117-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
3	NDC:25000-117-20	4 in 1 BOX	07/25/2011	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-117-30	6 in 1 BOX	07/25/2011	
4		6500 in 1 BAG; Type 0: Not a Combination Product		
5	NDC:25000-117-98	1 in 1 BOX	07/25/2011	
5		34065 in 1 BAG; Type 0: Not a Combination Product		
6	NDC:25000-117-93	1 in 1 BOX	07/25/2011	
6		37470 in 1 BAG; Type 0: Not a Combination Product		

7	NDC:25000-117-69	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2011	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA091239	07/25/2011		

Labeler - MARKSANS PHARMA LIMITED (925822975)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	LABEL(25000-117) , MANUFACTURE(25000-114, 25000-117)

Revised: 12/2019

MARKSANS PHARMA LIMITED