

ADVIL LIQUIGEL- ibuprofen capsule, liquid filled
Select Consumer Group

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

Advil Liqui-Gels

FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution


Advil Liqui-Gels Minis

FD&C green no. 3, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

Questions or comments?

call toll free **1-800-88-ADVIL**

PRINCIPAL DISPLAY PANEL



Drug Facts

Active ingredient (in each capsule): Solvated ibuprofen equal to 200 mg ibuprofen (NSAID)¹ (present as the free acid and potassium salt)
Purpose: Pain reliever/fever reducer
 Nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:
 ■ headache ■ toothache ■ backache
 ■ menstrual cramps ■ the common cold
 ■ muscular aches ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin redness ■ rash ■ blistering
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or

Drug Facts (continued)

steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ if you have ever had an allergic reaction to any other pain-reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if

■ stomach bleeding warning applies to you
 ■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic

Do not use if the sealed child resistant pouch is broken or torn.

2511793857



2 Liquid Filled Capsules

READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke:
 ■ chest pain ■ trouble breathing
 ■ weakness in one part or side of body
 ■ slurred speech ■ leg swelling

■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor

Other information

■ each capsule contains **potassium 20 mg**
 ■ read all warnings and directions before use.
 Keep card.
 ■ store at 20-25°C (68-77°F)
 ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients


FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

Questions or comments?

call toll free 1-855-297-3031

Packaged and distributed by Select Corporation with the permission of Haleon, Warren, NJ 07059
 For most recent product information, visit www.Advil.com
 Trademarks owned or licensed by Haleon.
 LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.
 ©2025 Haleon or licensor.
 Pat. info www.productpats.com





Drug Facts

Active ingredient (in each capsule): **Purpose**
Sukilled ibuprofen equal to Pain reliever/
200 mg ibuprofen (NSAID) Fever reducer
(present as the free acid and potassium salt)
Nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backaches
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma/wheezing
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or

Drug Facts (continued)

steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Do not use if the sealed child resistant pouch is broken or torn.

2521793868



4 Liquid Filled Capsules

READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

- you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body

- slurred speech
- leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 capsule, 2 capsules may be used

- do not exceed 6 capsules in 24 hours, unless directed by a doctor

- children under 12 years: ask a doctor

Other information

- each capsule contains potassium 20 mg
- read all warnings and directions before use. Keep card.

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

- FD&C green no. 3, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

Questions or comments?

- call toll free 1-855-267-3031

Packaged and distributed by Select Corporation with the permission of Haleon, Warren, NJ 07059. For most recent product information, visit www.Advil.com. Trademarks owned or licensed by Haleon. LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions. ©2025 Haleon or licensee. Pat. info www.productpats.com





ADVIL LIQUIGEL

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85237-1785(NDC:0573-0149)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
SOYBEAN LECITHIN (UNII: 1DI56QDM62)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green (Transparent green to blue green)	Score	no score
Shape	CAPSULE (oblong softgel)	Size	20mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85237-1785-1	1 in 1 BLISTER PACK	03/28/2025	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:85237-1785-2	2 in 1 BLISTER PACK	12/28/2025	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:85237-1785-4	25 in 1 BOX, UNIT-DOSE	12/28/2025	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:85237-1785-5	30 in 1 BOX, UNIT-DOSE	12/28/2025	
4		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020402	04/10/1995	

Labeler - Select Consumer Group (119185084)

Registrant - Select Consumer Group (119185084)

Establishment

Name	Address	ID/FEI	Business Operations
Select Consumer Group		119185084	repack(85237-1785)

Revised: 12/2025

Select Consumer Group