

BAYER GENUINE ASPIRIN- aspirin tablet
Select Consumer Group

Bayer Genuine Aspirin®

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature

Inactive ingredients

carnauba wax ², corn starch, hypromellose, powdered cellulose, triacetin

2 may contain this ingredient

Questions or comments?

1-800-331-4536

(Mon - Fri 9AM - 5PM EST) or www.bayeraspirin.com

Bayer 1

BAYER  **2 COATED TABLETS SINGLE-PACK**

ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER

GENUINE ASPIRIN
The name that
SAFE PAIN RELIEF +
LIFESAVING BENEFITS

 **325 mg**

 Talk to your doctor or other healthcare provider before using this product for your heart.

The product is packaged by: Seltzer Corporation
402 Newburgh Ave., Conditon, TX 75040
817-344-4428 W - 8 Sun - 5pm,
from a product manufactured by Bayer.
Bld. by Bayer Healthcare LLC, Whisper, IN 47981
©2023 Bayer. Bayer, the Bayer Cross, and
The Wonder Drug are registered trademarks of Bayer.
PRODUCT OF GERMANY

2511828098

Do Not Use if the sealed child resistant pouch is broken or torn.

<p>Drug Facts</p> <p>Active ingredient (in each tablet): Aspirin 325 mg (NSAID) Purpose(s): Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug.</p> <p>Use: temporarily relieves minor aches and pains due to: ● headache ● muscle pain ● backache ● menstrual pain ● colds ● minor pain of arthritis ● temporarily reduces fever</p> <p>Warnings: Roye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior such as nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Roye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include: ● hives ● facial swelling ● asthma (wheezing) ● shock</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed</p> <p>Do not use: ● if you are allergic to aspirin or any other pain reliever/fever reducer ● if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if: ● stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ● you are taking a diuretic ● you have asthma</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for: ● gout ● diabetes ● arthritis</p>	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if: ● an allergic reaction occurs. Seek medical help right away. ● you experience any of the following signs of stomach-bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● redness or swelling is present ● new symptoms occur ● ringing in the ears or a loss of hearing occurs</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions: ● drink a full glass of water with each dose ● adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours ● children under 12 years: consult a doctor</p> <p>Other information: ● save blister card for full directions and warnings ● store at room temperature</p> <p>Inactive ingredients: cane starch, hydroxytoluene, powdered cellulose, titanium</p> <p>Questions or comments? 1-800-331-6338 Mon - Fri 9AM - 5PM EST</p> <p> 6 55708 11828 3</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Bayer2



Bayer 50



BAYER GENUINE ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85237-1828(NDC:0280-2000)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	BAYER
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85237-1828-1	1 in 1 BLISTER PACK	12/24/2025	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:85237-1828-2	2 in 1 BLISTER PACK	12/24/2025	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:85237-1828-4	25 in 1 BOX	12/24/2025	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:85237-1828-6	50 in 1 BOX	12/24/2025	
4		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/31/2014	

Labeler - Select Consumer Group (119185084)

Registrant - Select Consumer Group (119185084)

Establishment

Name	Address	ID/FEI	Business Operations
Select Consumer Group		119185084	repack(85237-1828)

