DG HEALTH NAPROXEN SODIUM- naproxen sodium tablet, film coated Dolgencorp Inc

Dolgencorp, LLC Naproxen Sodium Tablets, 220 mg Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing

- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- · drink a full glass of water with each dose

Adults and children 12 years and older	 take 1 caplet every 8 to 12 hours while symptoms last for the first dose you may take 2 caplets within the first hour do not exceed 2 caplets in any 8- to 12-hour period do not exceed 3 caplets in a 24-hour period
Children under 12 years	ask a doctor

Other information

- each caplet contains: sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-888-309-9030

Principal Display Panel

DG[®]|health

Compare to the active ingredient of Aleve® Caplets

Naproxen Sodium Tablets, 220 mg

Pain Reliever/Fever Reducer (NSAID)


STRENGTH TO LAST 12 HOURS

Actual Caplet Size

200 Caplets†

(†Capsule Shaped Tablets)

220 mg



When using this product

■ take with food or milk if stomach upset occurs Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

■ feel faint

vomit blood
 have bloody or black stools
 have stomach pain that does not get better
I you have symptoms of heart problems or stroke:

■ weakness in one part or side of body
■ slured speech
■ leg swelling
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area

weakness in one part or side of body

■ trouble breathing

any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

do not take more than directed
the smallest effective dose should be used
drink a full glass of water with each dose

Adults and children

- ass or water with each dose

 take 1 caplet every 8 to 12 hours
 while symptoms last
 for the first dose you may take
 2 caplets within the first hour
 do not exceed 2 caplets in any
 8- to 12-hour period
 do not exceed 3 caplets in a
 24-hour period

Children under 12 years

Other information
■ each caplet contains: sodium 22 mg
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?1-888-309-9030

DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE, TN 37072

-100% Satisfaction Guaranteed! (888) 309-9030

DG* health

DG[®] health

Naproxen Sodium Tablets, 220 mg Pain Reliever/ Fever Reducer (NSAID) KEEP PACKAGE FOR REFERENCE

> Compare to the active ingredient of Aleve® Caplets**

DO NOT USE IF

IS BROKEN OR MISSING

Naproxen Sodium

Tablets, 220 mg Pain Reliever/ Fever Reducer (NSAID)





36882 VT C5

Drug Facts

Active ingredient (in each caplet)
Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID)*....*
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- Itemporarily relieves name
 Itemporarily relieves name
 Immor pain of arthritis
 Immor pain of arthritis ■ temporarily relieves minor aches and pains due to:
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- take a blood thinning (anticoagulant) or steroid drug
 take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen,
- naproxen, or others)
 have 3 or more alcoholic drinks every day while using this product
 take more or for a longer time than directed
- Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.





DG HEALTH NAPROXEN SODIUM

naproxen sodium tablet, film coated

Product Information

HUMAN OTC DRUG NDC:55910-500 **Item Code (Source) Product Type**

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ) NAPROXEN SODIUM 220 mg

Inactive Ingredients		
Ingredient Name	Strength	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics					
Color	BLUE (Light Blue)	Score	no score		
Shape	OVAL	Size	12mm		
Flavor		Imprint Code	L368		
Contains					

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:55910-500- 71	1 in 1 CARTON	11/21/2013	10/31/2023		
1		50 in 1 BOTTLE; Type 0: Not a Combination Product				
2	NDC:55910-500- 72	1 in 1 CARTON	11/21/2013	09/30/2021		
2		60 in 1 BOTTLE; Type 0: Not a Combination Product				
3	NDC:55910-500- 78	1 in 1 CARTON	11/21/2013	07/31/2023		
3		100 in 1 BOTTLE; Type 0: Not a Combination Product				
4	NDC:55910-500- 76	1 in 1 CARTON	11/21/2013	10/16/2018		
4		120 in 1 BOTTLE; Type 0: Not a Combination Product				
5	NDC:55910-500- 82	1 in 1 CARTON	09/05/2017			
5		200 in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA074661	11/21/2013			

Revised: 1/2025 Dolgencorp Inc