

**MAXIMUM STRENGTH ANTACID MINT- aluminum hydroxide, magnesium hydroxide, dimethicone liquid**  
**CVS**

-----  
**CVS ANTACID MINT MAX**

**Active ingredients (in each 10 mL dose)**

Aluminum hydroxide 800 mg (equivalent to dried gel, USP)  
Magnesium hydroxide 800 mg  
Simethicone 80 mg

**Purposes**

Antacid

Antigas

**Uses**

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

**Warnings**

**Do not take** more than 60 mL (6 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Directions**

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals, at bedtime, or as directed by a doctor
- children under 12 years of age: ask a doctor

**Other information**

- **each 10 mL dose contains:** magnesium 340 mg, sodium 10 mg
- protect from freezing
- store at room temperature
- keep tightly closed

**Inactive ingredients**

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

**Questions or comments?**

**1-800-540-3765**

**package label**



**MAXIMUM STRENGTH**

# Antacid Liquid

**ALUMINA / MAGNESIA / SIMETHICONE**  
Antacid/Anti-gas

Fast-Acting

**Soothing relief of:**

- Heartburn
- Acid Indigestion
- Pressure and Bloating



Mint Flavor

12 FL OZ (355 mL)



205-06242-3  
REV GC-0621

## Drug Facts

**TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON BOTTLE CAP IS MISSING OR BROKEN.**

### Active ingredients (in each 10 mL dose)

Aluminum hydroxide (equivalent to dried gel, USP) 800 mg .... Antacid  
Magnesium hydroxide 800 mg ..... Antacid  
Simethicone 80 mg ..... Antigas

### Purposes

**Uses** relieves: • heartburn • sour stomach  
• acid indigestion • symptoms referred to as gas

### Warnings

Do not take more than 60 mL (6 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

Ask doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

### Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals, at bedtime, or as directed by a doctor
- children under 12 years of age: ask a doctor

### Other information

- each 10 mL dose contains: magnesium 340 mg, sodium 10 mg
- store at room temperature • protect from freezing
- keep tightly closed

**Inactive ingredients** benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

**Questions or comments?** 1-800-540-3765

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
©2015 CVS/Pharmacy  
CVS.com® 1-800-SHOP CVS  
V-13966  
905-06242-5 REV GC-0821

#696216



## MAXIMUM STRENGTH ANTACID MINT

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-296
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	800 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 10 mL
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-296-73	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2015	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	06/01/2015	

**Labeler** - CVS (062312574)

**Registrant** - GCP Laboratories (965480861)

## Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(69842-296)

