

**IBUPROFEN- ibuprofen capsule, liquid filled
WALGREENS**

610T Walgreen 0363-1610 Ibuprofen Capsules 200 mg

DRUG FACTS

Active ingredient (in each capsule)

Solubilized ibuprofen equal to
200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to :

headache
toothache
backache
menstrual cramps
the common cold
muscular aches
minor pain of arthritis

- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

- you have symptoms of heart problems or stroke:

chest pain

trouble breathing

weakness in one part or side of body

slurred speech

leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 capsule, 2 capsules may be used

do not exceed 6 capsules in 24 hours, unless directed by a doctor

- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? call **1-877-290-4008**

NDC 0363-1610-80



Ibuprofen

IBUPROFEN CAPSULES, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

80 SOFTGELS** (**liquid-filled capsules)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain or arthritis

■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions ■ **do not take more than directed**

■ **the smallest effective dose should be used**

■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used

■ do not exceed 6 capsules in 24 hours, unless directed by a doctor

■ children under 12 years: ask a doctor

Each capsule contains: potassium 20 mg

Questions or comments? call 1-877-290-4008

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2021 Walgreen Co.
MADE IN INDIA




how2recycle.info
PLASTIC BOTTLE

610T REV 0921
ITEM 934532
W00000-0000-0
W3ORG1221-F
REV-0322

Lot No.:
Exp. Date:

Varnish Omit Area

NDC 0363-1610-44



Ibuprofen

IBUPROFEN CAPSULES 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

40 SOFTGELS** (**liquid-filled capsules)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis

■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions ■ **do not take more than directed**

■ **the smallest effective dose should be used**

■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used

■ do not exceed 6 capsules in 24 hours, unless directed by a doctor


■ children under 12 years: ask a doctor

Each capsule contains: potassium 20 mg

Questions or comments? call 1-877-290-4008

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2021 Walgreen Co.
MADE IN INDIA 610T REV 0921



PLASTIC BOTTLE

our934532w04
ITEM 934531
W00000-0000-0
W3ORG1221-F
REV-0322

Lot No.:
Exp. Date:

Varnish Omit Area

Adhesive Area

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease its benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
- sturred speech
- swelling
- your eyes worse or lasts more than 3 days
- fever, rash or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours with symptoms present
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other Information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F) - avoid excessive heat above 40°C (104°F)

Inactive ingredients

ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, stearic, sorbitan monooleate, sorbitol sorbitan

Questions or comments? call 1-877-296-4008

Our pharmacists recommend the Walgreens brand, Wainliu, you to compare to national brands. This product is not manufactured or distributed by JF Consumer Healthcare LLC, owner of the registered trademark, Liqui-Gels®.

VALUE SIZE NDC 0363-1610-31



Ibuprofen

IBUPROFEN CAPSULES, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

300 SOFTGELS** (**liquid-filled capsules)

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Purpose

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt), nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - headache
 - toothache
 - menstrual cramps
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Always alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

 Take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued under label)

ITEM 924534 W00000-0000-0
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com
 ©2021 Walgreen Co. MADE IN INDIA

Lot No.: 6107 REV 0821 W3ORG1221-F REV-1222
 Exp. Date: Varnish Omit Area



Walgreens

Ibuprofen

IBUPROFEN CAPSULES, 200 mg /
PAIN RELIEVER / FEVER REDUCER (NSAID)

• Strength to last 12 hours

20 SOFTGELS**
(**liquid-filled capsules)

NDC 0363-1610-23

**DO NOT REMOVE SEAL UNDER BOTTLE CAP IMPRINTED WITH
"SEALS" FOR YOUR PROTECTION. IF YOU NOTICE ANY
"SEALS" ARE BROKEN OR MISSING**

**READ AND KEEP OUTER CARTON FOR
COMPLETE WARNINGS AND INFORMATION**

Uses

- temporarily relieves minor aches and pains, due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs,

Drug Facts (continued under label)

Lot No. _____
Exp. Date _____
SEE HERE 

Varnish Omit Area

except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Each capsule contains: potassium 20 mg

Questions or comments? call

1-877-290-4008

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.

DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

Walgreens.com ©2021 Walgreen Co.

MADE IN INDIA

ITEM 934529

W000000-0000-0

W3ORGI221-F

610T REV 0921

REV-0723



Adhesive Area

Varnish Omit
Exp Date:
Lot No.:



ITEM 894529 W00000-0000-0
100% SATISFACTION GUARANTEED
DENTON, N.J. 08045
WALGREENS
© 2011 WALGREENS
owner of the registered trademark Liquid-Gels®
Consumer Healthcare LLC, Owner of the registered
trademark Advil® or Advil® Liquid-Gels®
This product is not manufactured or distributed by PF
The invite you to compare to national brands.
Our pharmacists recommend the Walgreens brand.

Drug Facts READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Active ingredient (in each capsule) Purpose
SoluBulized ibuprofen equal to..... Pain reliever/
200 mg ibuprofen (NSAID)*..... Fever reducer
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ● headache ● toothache ● backache ● menstrual cramps ● the common cold ● muscular aches ● minor pain of arthritis ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic

Drug Facts (continued)

drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:

Drug Facts (continued)

● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Other information ■ each capsule contains:

Drug Facts (continued)
potassium 20 mg ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)
Inactive ingredients: ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monobehate, sorbitol sorbitan
Questions or comments? call 1-877-290-0008
DO NOT USE IF SEAL UNDER BOTTLE CAP PROJECTS WITH "CALLED FOR YOUR PROTECTION" IS BROKEN OR MISSING

W3ORG1221-F
610T REV 0921
REV-0723

Varnish Omit



NDC 0363-1610-23



Ibuprofen
IBUPROFEN CAPSULES, 200 mg /
PAIN RELIEVER / FEVER REDUCER (NSAID)

20 SOFTGELS**
(**liquid-filled capsules)

ACTUAL SIZE

C6011R2
Varnish/Print Omit Area

Varnish Omit

Varnish Omit

NDC 0360-1610-66

Walgreens



Compare to the active ingredient in Advil® Liqui-Gels™

Ibuprofen

IBUPROFEN CAPSULES, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

160 SOFTGELS**

(**liquid-filled capsules)



ACTUAL SIZE

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active Ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*. . . . Pain reliever/fever reducer (present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others).

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer; if right before or after heart surgery.



IBUPROFEN CAPSULES, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- takes with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist



Drug Facts (continued)

- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients: ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? call 1-877-290-4008

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® or Catalent Pharma Solutions, Inc., owner of the registered trademark Liqui-Gels®.

DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD., DEERFIELD, IL 60015

Walgreens 100% SATISFACTION GUARANTEED

walgreens.com ©2021 Walgreen Co. MADE IN INDIA

6107 REV 09/21

C6011 CR2

Varnish/Print Omit Area

W3ORG1221-F REV-0522



ITEM 934533 W0000-0000-0
DO NOT USE IF SEAL UNDER BOTTLE CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Varnish Omit

Lot No.:
Exp Date:

Varnish Omit

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule) Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*. Pain reliever/Fever reducer (present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ toothache ■ backache
■ menstrual cramps ■ the common cold
■ muscular aches ■ minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

Drug Facts (continued)
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.



Drug Facts (continued)
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery
Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug
When using this product
■ take with food or milk if stomach upset occurs

Drug Facts (continued)
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING



Varnish/Print Omit Area
Lot No.:
Exp Date:

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used
■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor
Other information
■ each capsule contains: potassium 20 mg
■ read all warnings and directions before use
■ store at 20-25°C (68-77°F)
■ avoid excessive heat above 40°C (104°F)

Drug Facts (continued)
Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan
Questions or comments? call 1-877-290-4008
¹Our pharmacists recommend the Walgreens brand.
²We invite you to compare to national brands.
³This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® or Catalent Pharma Solutions, Inc., owner of the registered trademark Liqui-Gels®.
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
MADE IN INDIA
6107 REV/0921

W00RG1221-F
REV-0522



Ibuprofen

IBUPROFEN CAPSULES, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

40 SOFTGELS** (**liquid-filled capsules)




ACTUAL SIZE

Varnish Omit

Varnish Omit

Varnish/Print Omit Area C6011AR3

NDC 0363-1610-44



IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1610
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZ H3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1610-23	20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/27/2016	
2	NDC:0363-1610-44	40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/27/2016	
3	NDC:0363-1610-80	80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/27/2016	
4	NDC:0363-1610-66	160 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/27/2016	
5	NDC:0363-1610-31	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/27/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	01/27/2016	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(0363-1610)

Revised: 3/2024

WALGREENS