

PINWORM TREATMENT- pyrantel pamoate suspension
BIONPHARMA INC.

pinworm treatment-a+ health

Active ingredient (in each mL)

Pyrantel Pamoate 144mg/mL

(the equivalent of 50mg pyrantel base)

Purpose

Anthelmintic

Use

for the treatment of pinworms

Warnings

Do not exceed recommended dosage

Ask a doctor before use if

■ you are pregnant ■ have liver disease

When using this product

abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these symptoms persist, consult a doctor.

Keep out of reach of children.

In case of overdose get medical help or contact a poison control center right away.

Directions

■ **shake well before use**

■ **read bottle label and package insert carefully before taking this medication**

■ treat the entire household unless otherwise advised. See Warnings

■ do not repeat treatment unless directed by a doctor

■ this product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication

■ if signs of pinworms persist after treatment, consult a doctor

Weight	Dosage (taken as a single dose)
Less than 25 lb. or under 2 yrs.old	Do not use unless directed by a doctor
25-37 lb.	1/2 teaspoonful
38-62 lb.	1 teaspoonful
63-87 lb.	1 1/2 teaspoonfuls
88-112 lb.	2 teaspoonfuls
113-137 lb.	2 1/2 teaspoonfuls
138-162 lb.	3 teaspoonfuls
163-187 lb.	3 1/2 teaspoonfuls
188 lb. and over	4 teaspoonfuls

Other information

■ Store between 15°C-30°C (59°- 86°F)

Inactive ingredients

banana flavor, citric acid, glycerin, lecithin, magnesium aluminum silicate, methylcellulose, methylparaben, povidone, propylene glycol, propylparaben, purified water, simethicone, sodium benzoate, sodium citrate, sodium saccharin and sorbitol.

Questions or comments?

call toll-free **1-888-235-2466**

CAUTION: DO NOT USE IF IMPRINTED SEAL ON BOTTLE IS BROKEN OR MISSING

Distributed by:

BIONPHARMA

Princeton, NJ 08540

MADE IN INDIA

Code: TN/Drugs/TN00002222/2006

949129345

L0001113

R0725

READ THIS INSERT CAREFULLY BEFORE TAKING MEDICATION

PINWORM MEDICINE

Pyrantel Pamoate 144 mg/mL
(the equivalent of 50 mg Pyrantel base per mL)

FOR THE TREATMENT OF PINWORMS

DIRECTIONS

Adults, children 12 years of age and over, and children 2 years to under 12 years of age: oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight not to exceed 1 gram.

Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See warnings. If any worms other than pinworms are present before or after treatment, consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

Shake well before using.

Weight	Dosage (taken as a single dose)
Less than 25 pounds or under 2 years old	consult a doctor
25 to 37 pounds	½ teaspoonful
38 to 62 pounds	1 teaspoonful
63 to 87 pounds	1½ teaspoonfuls
88 to 112 pounds	2 teaspoonfuls
113 to 137 pounds	2½ teaspoonfuls
138 to 162 pounds	3 teaspoonfuls
163 to 187 pounds	3½ teaspoonfuls
188 pounds and over	4 teaspoonfuls

This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during, or after medication.

WARNINGS

Abdominal cramps, nausea, vomiting, diarrhea, headaches, or dizziness, sometimes occur after taking this drug, if any of these conditions persist consult a doctor.

If you are pregnant or have Liver disease, do not take this product unless directed by your doctor.

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

OTHER INFORMATION

This brochure will tell you how to identify and treat pinworm infections effectively with pinworm medicine. Many who have pinworms do not have symptoms, however, many will have itching in and around the rectal opening. The itching may be very annoying and continued scratching can cause an irritation in this area. The itching may be prominent at night when sleeping. That is why restless sleep is sometimes a sign of pinworms, especially in children. Other symptoms include insomnia, gastrointestinal distress,

irritability, enuresis (bed wetting) and secondary infection due to localized scratching. You should treat with pinworm medicine only after you have seen and identified the pinworms.

Pinworms look like tiny white threads. The female pinworm is about 1/2 inch long (the male pinworm is shorter); they live in the bowel. Usually, at night, the female pinworm travels to the rectal opening and lays eggs within the skin folds around the opening. This usually takes place within 1-2 hours after the child has been put to sleep for the night. The sticky gelatin like substance in which the eggs (they are too small to see) are deposited and the movement of the female pinworm may cause annoying itching and possibly restless sleep. By checking the rectal opening at this time, you can attempt to see the pinworm as described.

Scratching will cause pinworm eggs to stick to the fingers. Reinfection will result if the fingers are placed in the mouth. The eggs, which are too small to see, contaminate whatever they come in contact with: bed clothes, underwear, toys, hands, and food touched by contaminated hands. Even eggs floating in the air can be swallowed and cause infection. Eggs deposited can survive for as long as 3 weeks. Pinworms are highly contagious; therefore, after treatment with pinworm medication you should wash hands and fingernails with soap often during the day in order to prevent spreading to others and reinfection. Wear tight underpants both day and night. For several days after treatment clean the bedroom floor by vacuuming or damp mopping. After treatment, wash bed linens and night clothes (don't shake them). Keep toilet seats clean.

Distributed by: **Bionpharma Inc.**

Princeton, NJ 08540

MADE IN INDIA

948026846

L0000896

R0924

carton label 2oz

full prescription strength

a+ health TM

pinworm

treatment

pyrantel pamoate

suspension

- **Single dose effectiveness**
- **Measuring cup included**
- **Doctor recommended**

For the entire family (Ages 2 & up)

2 fl. oz. (60 ml)
contains two (2)-1oz bottles



bottle label 16oz
full prescription strength
a+ health™
pinworm
treatment
pyrantel pamoate
suspension

- Single dose effectiveness
- Doctor recommended

For the entire family (Ages 2 & up)

16 fl. oz. (473 ml)

20 mm x 30 mm
(W x H)

full prescription strength



pinworm treatment

pyrantel pamoate suspension

- Single dose effectiveness
- Doctor recommended



For the entire family (Ages 2 & up)

16 fl. oz. (473 ml)

Drug Facts

Active ingredient (in each mL) **Purpose**
Pyrantel Pamoate 144mg/mL
(the equivalent of 50mg pyrantel base)..... Anthelmintic

Use for the treatment of pinworms

Warnings

Do not exceed recommended dosage

Ask a doctor before use if ■ you are pregnant ■ have liver disease

When using this product abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these symptoms persist, consult a doctor.

Keep out of reach of children. In case of overdose get medical help or contact a poison control center right away.

Directions

- shake well before use
- Carefully read additional product information (located on the inside of label) before using this medication.
- treat the entire household unless otherwise advised
- do not repeat treatment unless directed by a doctor
- this product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication
- if signs of pinworms persist after treatment, consult a doctor

Weight	Dosage (taken as a single dose)
Less than 25 lb. or under 2 yrs. old	Do not use unless directed by a doctor
25-37 lb.	½ teaspoonful
38-62 lb.	1 teaspoonful
63-87 lb.	1½ teaspoonfuls
88-112 lb.	2 teaspoonfuls
113-137 lb.	2½ teaspoonfuls
138-162 lb.	3 teaspoonfuls
163-187 lb.	3½ teaspoonfuls
188 lb. and over	4 teaspoonfuls ▶



**CAUTION: DO NOT USE IF IMPRINTED
SEAL ON BOTTLE IS BROKEN OR MISSING**

UVZ
(Lot and Exp online Printing)

Drug Facts (continued)

Other information

- Store between 15°-30°C (59°-86°F)

Inactive ingredients banana flavour, citric acid, glycerin, magnesium aluminum silicate, methylcellulose, methylparaben, povidone, propylene glycol, propylparaben, purified water, simethicone, sodium benzoate, sodium citrate, sodium saccharin and sorbitol.

Questions or comments?

call toll-free 1-888-235-2466

Distributed by:

BIONPHARMA
Princeton, NJ 08540

MADE IN INDIA

Code: TN/Drugs/TN00002222/2006

948006869

R0725

L0001114



READ THIS PRODUCT INFORMATION CAREFULLY BEFORE TAKING MEDICATION

Pyrantel Pamoate Suspension
(Pyrantel base: 50 mg / mL)

FOR THE TREATMENT OF PINWORMS

DIRECTIONS Adults, children 12 years of age and over, and children 2 years to under 12 years of age: oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight not to exceed 1 gram (4 teaspoonfuls).

Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See warnings. If any worms other than pinworms are present before or after treatment, consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

Shake well before using.

Weight	Dosage (taken as a single dose)
Less than 25 lb. or under 2 yrs. old	Do not use unless directed by a doctor
25-37 lb.	½ teaspoonful
38-62 lb.	1 teaspoonful
63-87 lb.	1½ teaspoonfuls
88-112 lb.	2 teaspoonfuls
113-137 lb.	2½ teaspoonfuls
138-162 lb.	3 teaspoonfuls
163-187 lb.	3½ teaspoonfuls
188 lb. and over	4 teaspoonfuls

This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during, or after medication.

WARNINGS

Abdominal cramps, nausea, vomiting, diarrhea, headaches, or dizziness, sometimes occur after taking this drug, if any of these conditions persist, consult a doctor.

If you are pregnant or have liver disease, do not take this product unless directed by your doctor.

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

OTHER INFORMATION This product information will tell you how to identify and treat pinworm infections effectively with pinworm medicine. Many who have pinworms do not have symptoms, however, many will have itching in and around the rectal opening. The itching may be very annoying and continued scratching can cause an irritation in this area. The itching may be prominent at night when sleeping. That is why restless sleep is sometimes a sign of pinworms, especially in children. Other symptoms include insomnia, gastrointestinal distress, irritability, enuresis (bed wetting) and secondary infection due to localized scratching. You should treat with pinworm medicine only after you have seen and identified the pinworms.

Pinworms look like tiny white threads. The female pinworm is about 1/2 inch long (the male pinworm is shorter); they live in the bowel. Usually, at night, the female pinworm travels to the rectal opening and lays eggs within the skin folds around the opening. This usually takes place within 1-2 hours after the child has been put to sleep for the night. The sticky gelatin like substance in which the eggs (they are too small to see) are deposited and the movement of the female pinworm may cause annoying itching and possibly restless sleep. By checking the rectal opening at this time, you can attempt to see the pinworm as described.

Scratching will cause pinworm eggs to stick to the fingers. Reinfection will result if the fingers are placed in the mouth. The eggs, which are too small to see, contaminate whatever they come in contact with: bed clothes, underwear, toys, hands, and food touched by contaminated hands. Even eggs floating in the air can be swallowed and cause infection. Eggs deposited can survive for as long as 3 weeks. Pinworms are highly contagious; therefore, after treatment with pinworm medication you should wash hands and fingernails with soap often during the day in order to prevent spreading to others and reinfection. Wear tight underpants both day and night. For several days after treatment clean the bedroom floor by vacuuming or damp mopping. After treatment, wash bed linens and night clothes (don't shake them). Keep toilet seats clean.

No Text area
T00247R0725

PINWORM TREATMENT

pyrantel pamoate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69452-515
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PYRANTEL PAMOATE (UNII: 81BK194Z5M) (PYRANTEL - UNII:4QIH0N49E7)	PYRANTEL	144 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
METHYLCELLULOSE (UNII: Z944H5SN0H)	
POVIDONE (UNII: FZ989GH94E)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

SORBITOL (UNII: 506T60A25R)				
WATER (UNII: 059QF0KO0R)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
Product Characteristics				
Color	yellow (pale yellow to yellow)		Score	
Shape			Size	
Flavor	BANANA		Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69452-515-38	2 in 1 CARTON	09/08/2025	
1	NDC:69452-515-36	30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69452-515-88	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/08/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M024	09/08/2025	

Labeler - BIONPHARMA INC. (079637826)

Registrant - BIONPHARMA INC. (079637826)

Establishment

Name	Address	ID/FEI	Business Operations
OrBion Pharmaceuticals Private Limited		854403569	manufacture(69452-515)

Revised: 9/2025

BIONPHARMA INC.