

**MUCUS RELIEF SINUS PRESSURE PAIN AND COUGH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated**  
**CHAIN DRUG MARKETING ASSOCIATION INC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**QCH-1199-2020-1207**

**Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves:
  - headache
  - nasal congestion
  - minor aches and pains
  - sinus congestion and pressure
  - cough
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

- This product contains acetaminophen. Severe liver damage may occur if you take
- more than 12 caplets in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks daily while using this product

**Allergy alert**

- Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
  - blisters
  - rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- diabetes
- liver disease
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not use more than directed**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information**

- store between 20-25°C (68-77°F) in a dry place

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

**PRINCIPAL DISPLAY PANEL**

NDC 63868-587-20

QUALITY CHOICE®

†Compare to the Active Ingredients in MUCINEX® Fast-Max® Pressure, Pain & Cough  
Maximum Strength

Mucus Relief Sinus Pressure, Pain & Cough

Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Relieves Sinus Pressure & Congestion

Relieves Headache

Controls Cough

Thins & Loosens Mucus

Actual Size

20 CAPLETS

100% SATISFACTION GUARANTEED  
 Distributed by C.M.A., Inc.  
 43157 W 9 Mile Rd  
 Novi, MI 48375  
 www.qualitychoice.com  
 Questions: 800-935-2362

**DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN**

This product is not manufactured or distributed by Sinus-Max® Pressure, Pain & Cough.  
 Redit Design is distributor of Mucosin®

**Drug Facts (continued)**

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**  
 When using this product do not use more than directed

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning)  
 ■ do not take more than 12 caplets in any 24-hour period  
 ■ adults and children 12 years of age and over: take 2 caplets every 4 hours  
 ■ children under 12 years of age: do not use

**Other information**  
 ■ store between 20-25°C (68-77°F) in a dry place  
 ■ retain outer package for complete product information and warnings

**Inactive ingredients** colloidal silicon dioxide, corn starch, croscarmellose sodium, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, methylcellulose, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

**Drug Facts**

**Active ingredients (in each caplet) Purpose**  
 Acetaminophen 325 mg Pain reliever  
 Dextromethorphan HBr 10 mg Cough suppressant  
 Guaifenesin 200 mg Expectorant  
 Phenylephrine HCl 5 mg Nasal decongestant

**Uses** ■ temporarily relieves: ■ headache ■ minor aches and pains ■ sinus congestion and pressure ■ cough ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 12 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product  
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
 Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 Ask a doctor before use if you have ■ diabetes ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ▶



F1199020CH\_R0

**QC QUALITY CHOICE**  
**Maximum Strength**  
**Mucus Relief Sinus Pressure, Pain & Cough**  
 Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant  
 For Ages 12+

**QC QUALITY CHOICE**  
**Maximum Strength**  
**Mucus Relief Sinus Pressure, Pain & Cough**  
 Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant  
 For Ages 12+

**QC QUALITY CHOICE**  
 NDC 63868-587-20  
 \*Compare to Active Ingredients in MUCINEX® Sinus-Max® Pressure, Pain & Cough

**Maximum Strength**  
**Mucus Relief Sinus Pressure, Pain & Cough**  
 For Ages 12+

**Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant**

**Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl**  
 Relieves Sinus Pressure & Congestion  
 Relieves Headache  
 Controls Cough  
 Thins & Loosens Mucus

Actual Size



**20 CAPLETS**

NC

NC

NC

# MUCUS RELIEF SINUS PRESSURE PAIN AND COUGH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-587
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	AAA;1165
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:63868-587-20	2 in 1 CARTON	12/01/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/01/2020	

**Labeler** - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Revised: 12/2022

CHAIN DRUG MARKETING ASSOCIATION INC