

MUCUS RELIEF SINUS PRESSURE PAIN AND COUGH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated
CHAIN DRUG MARKETING ASSOCIATION INC

QCH-1199-2020-1207

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - cough
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- liver disease
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F) in a dry place

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

NDC 63868-587-20

QUALITY CHOICE®

†Compare to the Active Ingredients in MUCINEX® Fast-Max® Pressure, Pain & Cough

Maximum Strength

Mucus Relief Sinus Pressure, Pain & Cough

Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Relieves Sinus Pressure & Congestion

Relieves Headache

Controls Cough

Thins & Loosens Mucus

Actual Size

20 CAPLETS

100% SATISFACTION GUARANTEED
 Distributed by C.M.A., Inc.
 43157 W 9 Mile Rd
 Novi, MI 48375
 www.qualitychoice.com
 Questions: 800-935-2362

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

This product is not manufactured or distributed by Sinus-Max® Pressure, Pain & Cough.
 Redit Design is distributor of Mucus Relief Sinus Pressure, Pain & Cough.



Maximum Strength
Mucus Relief Sinus Pressure, Pain & Cough
 Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

For Ages 12+

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin ■ When using this product do not use more than directed <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts <p>These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed (see Overdose warning) ■ do not take more than 12 caplets in any 24-hour period ■ adults and children 12 years of age and over: take 2 caplets every 4 hours ■ children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ store between 20-25°C (68-77°F) in a dry place ■ retain outer package for complete product information and warnings <p>Inactive ingredients</p> <p>colloidal silicon dioxide, corn starch, croscarmellose sodium, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, methylcellulose, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide</p>	<p>Drug Facts</p> <p>Active ingredients (in each caplet): Purpose</p> <p>Acetaminophen 325 mg.....Pain reliever Dextromethorphan HBr 10 mg.....Cough suppressant Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves: <ul style="list-style-type: none"> ■ headache ■ nasal congestion ■ minor aches and pains ■ sinus congestion and pressure ■ sinusitis ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 12 caplets in 24 hours, which is the maximum daily amount.</p> <ul style="list-style-type: none"> ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin redness ■ blisters ■ rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ diabetes ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)
---	--



F1199020CH_R0



Maximum Strength
Mucus Relief Sinus Pressure, Pain & Cough
 Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

For Ages 12+



NDC 63868-587-20

*Compare to Active Ingredients in MUCINEX® Sinus-Max® Pressure, Pain & Cough

For Ages 12+

Maximum Strength
Mucus Relief Sinus Pressure, Pain & Cough

Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl
 Relieves Sinus Pressure & Congestion
 Relieves Headache
 Controls Cough
 Thins & Loosens Mucus

Actual Size



20 CAPLETS

NC

NC

NC

MUCUS RELIEF SINUS PRESSURE PAIN AND COUGH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-587
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	AAA;1165
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:63868-587-20	2 in 1 CARTON	12/01/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2020	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Revised: 12/2024

CHAIN DRUG MARKETING ASSOCIATION INC