ACID GONE ANTACID- aluminum hydroxide and magnesium carbonate liquid Major Pharmaceuticals

MAJOR ACID GONE

Active ingredient (in each 15mL tablespoonful)

Aluminum hydroxide 95mg

Magnesium carbonate 358mg

Purpose

Antacid

Uses

relieves

- heartburn
- · acid indigestion
- sour stomach
- upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

- do not take more than 8 tablespoonfuls in 24 hours
- do not use the maximum dosage for more than 2 weeks
- laxative effect may occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- adults and children 12 years and older: take 1 to 2 tablespoonfuls four times a day or

- as directed by a doctor
- children under 12 years: ask a doctor
- take after meals and at bedtime
- dispense product only by spoon or other measuring device

Other information

- each 15 mL tablespoonful contains: magnesium 105mg, sodium 39mg
- store at up to 25°C (77°F)
- avoid freezing
- keep tightly closed
- **TAMPER-EVIDENT:** Do not use this product if inner foil seal over the mouth of the bottle is broken or missing.

Inactive ingredients (Regular Strength)

benzyl alcohol, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, purified water, saccharin sodium, sodium alginate, sorbitol solution, xanthan gum

Questions or comments?

1-800-540-3765

Principal Display Panel

major

NDC 0904-7727-14

Acid Gone Antacid

Alumina and Magnesium Carbonate Oral Suspension USP

Relieves Heartburn & Acid Indigestion

Compare to the active ingredients in Gaviscon® *

12 FL OZ (355 mL)

NDC 0904-7727-14 **MAJOR** cid Gone

> Alumina and Magnesium Carbonate Oral Suspension USP

Relieves Heartburn & Acid Indigestion



Compare to the active ingredients in Gaviscon®*

12 FL 0Z (355 mL)

226-15212-5 Rev. 03/17

Drug Facts

Active ingredients (in each 15 mL tablespoonful) Aluminum hydroxide 95 mg

Purpose Antacid Antacid

Magnesium carbonate 358 mg

Uses relieves • acid indigestion • heartburn • sour stomach · upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have • kidney disease

a magnesium-restricted diet

· a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

- do not take more than 8 tablespoonfuls in 24 hours
- do not use the maximum dosage for more than 2 weeks
- laxative effect may occur

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • shake well before use

- adults and children 12 years and older: take 1 to 2 tablespoonfuls four times a day or as directed by a doctor
- children under 12 years: ask a doctor
- take after meals and at bedtime
- dispense product only by spoon or other measuring device

Other information

- each 15 mL tablespoonful contains: magnesium 105 mg, sodium 39 mg
- store at up to 25°C (77°F) avoid freezing keep tightly closed TAMPER-EVIDENT: Do not use this product if inner foil seal over the mouth of the bottle is broken or missing.

Inactive ingredients benzyl alcohol, D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavor, glycerin, purified

water, saccharin sodium, sodium alginate, sorbitol solution, xanthan gum

Questions or comments? 1-800-540-3765

This product is not manufactured or distributed by the owner of the registered trademark Gaviscon®.

926-15212-5

Re-Order No. 264978

Distributed by MAJOR® PHARMACEUTICALS

17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152

Rev 03/17 M-99

ACID GONE ANTACID

aluminum hydroxide and magnesium carbonate liquid

Product Information

HUMAN OTC DRUG NDC:0904-7727 **Product Type** Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	95 mg in 15 mL
MAGNESIUM CARBONATE (UNII: 0E53J927NA) (CARBONATE ION - UNII:7UJQ50PE7D)	MAGNESIUM CARBONATE	358 mg in 15 mL

Inactive Ingredients				
Ingredient Name	Strength			
BENZYL ALCOHOL (UNII: LKG8494WBH)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GLYCERIN (UNII: PDC6A3C0OX)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SODIUM ALGINATE (UNII: C269C4G2ZQ)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				
WATER (UNII: 059QF0KO0R)				
XANTHAN GUM (UNII: TTV12P4NEE)				

Product Characteristics			
Color	green	Score	
Shape		Size	
Flavor	SPEARMINT (cool mint)	Imprint Code	
Contains			

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0904-7727- 14	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2017	

Marketing Information			
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	12/30/2004	

Labeler - Major Pharmaceuticals (191427277)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(0904-7727)	

Revised: 11/2023 Major Pharmaceuticals