

TEMCONGYI- salicylic acid liquid liquid
Hubei Redrui Building Materials Co., Ltd.

Active ingredient(s)

Salicylic acid 12%

Purpose(s)

Wart and corn remover

Uses

- For the removal of common warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface.
- For the removal of corns and calluses.

Warnings

- For external use only.
- Flammable. Keep away from fire or flame.

Do not use

- On irritated skin or on any area that is infected or reddened.
- If you are allergic to any of the ingredients in this product.
- On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes.

When using this product

- Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes.
- Avoid inhaling vapors.
- Cap bottle tightly when not in use.

Stop use and ask a doctor if

- Discomfort persists.

Pregnancy or breast feeding section

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash affected area.
- May soak wart in warm water for 5 minutes.
- Dry area thoroughly.
- Using the applicator, apply the liquid one drop at a time to sufficiently cover each wart, corn, or callus.
- Let dry.
- Repeat this procedure once or twice daily as needed (until wart, corn or callus is removed) for up to 12 weeks.

Other information

- Store between 20-25°C(68-77°F). Protect from excessive heat above 40°C(104°F).

Inactive ingredients

ACETIC ACID, ALCOHOL, ANGELICA DAHURICA ROOT, AQUA, ASIATIC ACID, BENZOIC ACID, PROPYLENE GLYCOL

Questions or comments?

934-935-7135 / Yiqinzhi726@gmail.com

Box Front Panel

ADVANCED FORMULA
FAST ACTING LIQUID
TURN TO MASK IN 30 SEC
SAFELY & EFFICIENTLY
PRECISE REMOVED

88 mm

Drug Facts	
Active ingredient(s) (in each dosage unit)	Purpose(s)
Salicylic Acid 12%.....	Wart remover
USES	
To remove warts such as plantar wart, flat wart, common wart and corns, calluses.	
WARNINGS	
• For external use only. • When using this product, avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes. • Cap bottle tightly and store at room temperature away from heat. • Keep out of reach of children. If swallowed, get medical help or contact Poison Control Center right away	
DIRECTIONS	
1. Soak for 5 minutes in warm water. 2. Insert the brush in the bottle. 3. Apply the liquid to the desired area. 4. Apply it 2 times daily to enjoy your clean skin.	
INACTIVE INGREDIENTS	
Aqua, Benzoic Acid, Propylene Glycol, Alcohol Denat, Acetic Acid.	
Other information:	
· store between 20° to 25°C (68° to 77°F). Protect from excessive heat above 40C (104°F).	

Temcongyi

LIQUID WART & CORN REMOVER

ADVANCED FORMULA

FAST ACTING LIQUID



USES:
Corns, Calluses, Flat warts, Plantar warts, Common warts

NET WT
1 FL OZ (30 ML)



Drug Facts

DO NOT USE

1. Do not use on irritated skin or on any area that is infected or reddened.

2. If you are allergic to any of the ingredients in this product.

3. On moles, birthmarks, warts with hairs growing from them, genital warts or warts on the face or mucous membranes.

STOP USE AND SEE A DOCTOR IF
You become pregnant, or are planning to become pregnant, while using the product.

if discomfort persists.



Temcongyi

LIQUID WART & CORN REMOVER

ADVANCED FORMULA

FAST ACTING LIQUID







USES:
Corns, Calluses, Flat warts, Plantar warts, Common warts

NET WT
1 FL OZ (30 ML)



37 mm

43 mm

Drug Facts		Temcongyl		When using this product	
Active ingredient(s) (in each dosage unit)		Purpose(s)		Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes.	
Salicylic Acid 12%.....		Wart remover		Avoid inhaling vapors.	
USES		<div><div>LIQUID</div><div>WART & CORN</div><div>REMOVER</div><div>ADVANCED FORMULA</div><div>FAST ACTING LIQUID</div></div> <div><div></div><div></div><div></div></div>		Pregnancy or breast feeding section If pregnant or breast-feeding, ask a health professional before use.	
For the removal of common warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface. For the removal of corns and calluses.				Keep out of reach of children If swallowed, get medical help or contact a Poison Control Center right away.	
WARNINGS		<div><div>FAST ACTING LIQUID</div></div> <div><div>USES:</div><div>Corns/Calluses, Flat warts, Plantar warts, Common warts</div></div>		Directions 1. Wash affected area. May soak in warm water for 5 minutes dry area thoroughly. 2. Apply a thin layer of the product with the applicator to cover each wart or corn let dry. 3. Repeat this procedure once or twice daily as needed (until wart or corn is removed) for up to 12 weeks.	
For external use only. Flammable. Keep away from fire or flame.				Other information Store between 20-25°C(68-77°F). Protect from excessive heat above 40°C(104°F).	
DO NOT USE		<div><div>NET WT</div><div>1 FL OZ (30 ML)</div><div></div></div>		Inactive ingredients ACETIC ACID, ALCOHOL DENAT, ANGIOLICA DAHURICA EXTRACT, AQUA, ASARIC ACID, BENZOIC ACID, PROPYLENE GLYCOL.	
1. On irritated skin or on any area that is infected or reddened. 2. If you are allergic to any of the ingredients in this product. 3. On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes.					
Stop use and ask a doctor if					
Discomfort persists.					

90 mm

TEMCONGYI

salicylic acid liquid liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:87066-5827
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	3.6 g in 30 g

Inactive Ingredients

Ingredient Name	Strength
AQUA (UNII: 059QF0KO0R)	21.285 g in 30 g
ACETIC ACID (UNII: Q40Q9N063P)	1 g in 30 g
BENZOIC ACID (UNII: 8SKN0B0MIM)	0.015 g in 30 g
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	1.5 g in 30 g
ASIATIC ACID (UNII: 9PA5A687X5)	1 g in 30 g
ALCOHOL (UNII: 3K9958V90M)	0.6 g in 30 g
ANGELICA DAHURICA ROOT (UNII: 1V63N2S972)	1 g in 30 g

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:87066-5827-1	1 g in 1 CARTON; Type 0: Not a Combination Product	08/27/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	08/27/2025	

Labeler - Hubei Redrui Building Materials Co., Ltd. (617973213)

Registrant - Hubei Redrui Building Materials Co., Ltd. (617973213)

Establishment			
Name	Address	ID/FEI	Business Operations
Guangzhou Yishenghe Traditional Chinese Medicine Technology Co., Ltd.		638316827	manufacture(87066-5827)

Establishment			
Name	Address	ID/FEI	Business Operations
Hubei Redrui Building Materials Co., Ltd.		617973213	label(87066-5827)

Revised: 8/2025

Hubei Redrui Building Materials Co., Ltd.