

CVS ACETAMINOPHEN- acetaminophen capsule
CVS Pharmacy

CVS Acetaminophen

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis • toothache • muscular aches • premenstrual and menstrual cramps
- temporarily reduces fever

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	• take 2 caplets every 6 hours while symptoms last • do not take more than 6 caplets in 24 hours, unless directed by a doctor • do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning:In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Other information

- store between 20°–25°C (68°–77°F)

Inactive ingredients

colloidal silicon dioxide, hypromellose, maize starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, sodium starch glycolate, titanium dioxide.

Questions or comments?

Call 1-800-SHOP CVS

Package Labeling:



actual size

ACETAMINOPHEN

Acetaminophen, 500 mg
Pain Reliever, Fever Reducer
For adults

Compare to the active ingredient
in Extra Strength Tylenol®*

Infinitely Recyclable Aluminum

500 CAPLETS

NDC 51316-185-00

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE

Drug Facts

Active ingredients (in each caplet)

Purpose

Acetaminophen 500 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20°–25°C (68°–77°F)

Inactive ingredients

colloidal silicon dioxide, hypromellose, maize starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, sodium starch glycolate, titanium dioxide.

Questions or comments?

Call 1-800-SHOP CVS

[†]This product is not manufactured or distributed by
Johnson & Johnson Consumer Inc., distributor of
Tylenol® Extra Strength.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2021 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

Made in India ITEM# 923789

Lot No. and Exp.
Date: see bottom
of bottle



0 50428 16001 5

Compare to the active ingredient in
Tylenol® Extra Strength†

Acetaminophen 500 mg
Pain Reliever, Fever Reducer

For adults

500 caplets

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE

Drug Facts

Active ingredients (in each caplet)

Purpose

Acetaminophen 500 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.
Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20°–25°C (68°–77°F)

Inactive ingredients

colloidal silicon dioxide, hypromellose, maize starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, sodium starch glycolate, titanium dioxide.

Questions or comments?

Call 1-800-SHOP CVS

[†]This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Tylenol® Extra Strength.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2021 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

Lot No. and Exp. Date:

see bottom of bottle

Made in India

ITEM# 923789



CVS ACETAMINOPHEN

acetaminophen capsule

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-185	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg	
Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
STARCH, CORN (UNII: O8232NY3SJ)				
MAGNESIUM STEARATE (UNII: 70097M6I3O)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	white	Score	no score	
Shape	CAPSULE	Size	18mm	
Flavor		Imprint Code	GL5	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-185-00	500 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		10/01/2025	

Labeler
- CVS Pharmacy (062312574)

Registrant
- Cabinet Health P.B.C. (117102391)

