

**CVS IBUPROFEN- ibuprofen tablet**  
**CVS Pharmacy**

-----  
**CVS Ibuprofen**

***Drug Facts***

***Active ingredient (in each tablet)***

Ibuprofen 200 mg (NSAID)\*\*

\*\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/Fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- headache • toothache • backache • menstrual cramps • the common cold • muscular aches • minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:**Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:**NSAIDs except aspirin increase the risk of

heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 day
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

***Directions***

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- read all warnings and directions before use.
- store between 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

***Inactive ingredients***

colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin.

***Questions or comments?***

Call 1-908-242-6108 Mon-Fri 8:00 AM to 5:00 PM EST

***Package Labeling:***

Infinately Recyclable Aluminum  
500 COATED TABLETS

Compare to the active  
ingredient in Advil®\*

Pain reliever/Fever reducer (NSAID)

**IBUPROFEN**

Ibuprofen, 200 mg

actual size



NDC 51316-187-00

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**  
IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE

### Drug Facts

**Active ingredient (in each tablet)** *Purpose*

Ibuprofen 200 mg (NSAID)\*\* .....Pain reliever/Fever reducer  
\*\*nonsteroidal anti-inflammatory drug

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs except aspirin increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

\*This product is not manufactured or distributed by  
Haleon, owner of the registered trademark Advil®.

(Continued on  
back of the label)

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2021 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS

Made in India  
ITEM#923748



Lot No. and Exp. Date:  
see bottom of bottle



0 50428 16033 6

### Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- read all warnings and directions before use.
- store between 20°–25°C (68°–77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients** colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin.

**Questions or comments?** Call 1-800-SHOP CVS

Compare to the active ingredient  
in Advil®

Ibuprofen  
Tablets 200mg  
Pain reliever/Fever reducer (NSAID)

500 coated tablets

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE

### Drug Facts

**Active ingredient (in each tablet)** **Purpose**

Ibuprofen 200 mg (NSAID)\*\* ..... Pain reliever/Fever reducer  
\*\*nonsteroidal anti-inflammatory drug

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache ■ toothache ■ backache ■ menstrual cramps
  - the common cold ■ muscular aches ■ minor pain of arthritis
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening
- rash ■ blisters

if an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs except aspirin increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

This product is not manufactured or distributed by Wyeth Consumer Healthcare, owner of the registered trademark Advil®

(Continued on back of the label)

Distributed by CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2021 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS

Made in India  
ITEM#923748

Lot No. and Exp. Date: see bottom of bottle



HINGE AREA  
TEXT-FREE ZONE  
NO PRINT/NO VARNISH

BACKGROUND  
OF DRUG FACTS  
SHOULD BE WHITE

### Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs

#### Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain ■ trouble breathing ■ weakness in one part or side of body
  - slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- read all warnings and directions before use.
- store between 20°–25°C (68°–77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients** colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, povidone K-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide, triacetin.

**Questions or comments?** Call 1-800-SHOP CVS

51316-187-01



NDC 51316-187-01

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE**

### Drug Facts

**Active ingredient (in each tablet) Purpose**  
Ibuprofen 200 mg (NSAID)\*\* .....Pain reliever/Fever reducer  
\*\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ headache ■ toothache ■ backache  
■ menstrual cramps ■ the common cold  
■ muscular aches ■ minor pain of arthritis  
■ temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs except aspirin increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Drug Facts (continued)

#### Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery

#### Ask a doctor before use if

■ the stomach bleeding warning applies to you  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

#### When using this product

■ take with food or milk if stomach upset occurs

#### Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

■ **do not take more than directed**  
■ the **smallest effective dose should be used** ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 tablet, 2 tablets may be used  
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor  
■ children under 12 years: ask a doctor

### Other information

■ read all warnings and directions before use.  
■ store between 20–25°C (68–77°F)  
■ avoid excessive heat above 40°C (104°F)

### Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin.

**Questions or comments?** contact 1-908-242-6108 Mon-Fri 8:00 AM to 5:00 PM EST

\*This product is not manufactured or distributed by Haleon, owner of the registered trademark Advil.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2026 CVS/Pharmacy  
CVS.com® 1-800-SHOP-CVS

100% money back guarantee. CVS.com/returnpolicy

Made in India V-41117

Lot No. and Exp. Date: see bottom of bottle

#454604



0 50428 29729 2

# IBUPROFEN

Tablets, 200 mg  
Pain Reliever  
Fever Reducer (NSAID)

Compare to Advil® active ingredient\*

1000 COATED TABLETS

## CVS IBUPROFEN

ibuprofen tablet

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:51316-187 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN         | 200 mg   |

### Inactive Ingredients

| Ingredient Name                                   | Strength |
|---|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                |          |
| STARCH, CORN (UNII: O8232NY3SJ)                   |          |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)      |          |
| FERRIC OXIDE RED (UNII: 1K09F3G675)               |          |
| MAGNESIUM STEARATE (UNII: 70097M6I30)             |          |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)     |          |
| SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D) |          |
| TALC (UNII: 7SEV7J4R1U)                           |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)               |          |
| TRIACETIN (UNII: XHX3C3X673)                      |          |

### Product Characteristics

|              |       |              |          |
|--------------|-------|--------------|----------|
| <b>Color</b> | brown | <b>Score</b> | no score |
| <b>Shape</b> | ROUND | <b>Size</b>  | 10mm     |

| <b>Flavor</b>                |  | <b>Imprint Code</b>                                 | IBU;200              |                    |
|------------------------------|--|---|----------------------|--------------------|
| <b>Contains</b>              |  |   |                      |                    |
| <b>Packaging</b>             |  |   |                      |                    |
| #                            | Item Code                                | Package Description                                 | Marketing Start Date | Marketing End Date |
| 1                            | NDC:51316-187-00                         | 500 in 1 BOTTLE; Type 0: Not a Combination Product  | 10/01/2025           |                    |
| 2                            | NDC:51316-187-01                         | 1000 in 1 BOTTLE; Type 0: Not a Combination Product | 02/01/2026           |                    |
| <b>Marketing Information</b> |  |   |                      |                    |
| Marketing Category           | Application Number or Monograph Citation | Marketing Start Date                                | Marketing End Date   |                    |
| ANDA                         | ANDA079129                               | 10/01/2025  |                      |                    |

**Labeler** - CVS Pharmacy (062312574)

Revised: 2/2026

CVS Pharmacy