NAPROXEN SODIUM- naproxen sodium tablet Wal-Mart Stores, Inc.,

Naproxen Tablets

Active ingredient

(in each caplet) Naproxen sodium 220 mg (NSAID)** **nonsteroidal anti-inflammatory drug

Purpose

Pain Reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause

problems in the unborn child or complications during delivery.

Keep out of the reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

adults and children 12 years and older,

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12- hour period
- do not exceed 3 caplets in a 24- hour period

children under 12 years,

ask a doctor

Other information

- each caplet contains: sodium 20 mg
- Store at 20 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).


Questions and comments?

1-888-287-1915

Inactive ingredients

FD&C blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide.





NAPROXEN	SODIUM
----------	--------

naproxen sodium tablet					
Product Information					
Product T ype	HUMAN OTC DRUG	Item Code (Source	NDC:4903	NDC:49035-762	
Route of Administration	ORAL				
	•				
Active Ingredient/Active M	oiety Ingredient Name				
1	Basis of Strength		Strengtl		
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ) NAPROXEN SODIUM					
T .* T . 1* .					
Inactive Ingredients					
Ingredient Name					Strength
TITANIUM DIO XIDE (UNII: 15FIX9)	/2JP)				
FD&C BLUE NO. 2 (UNII: L06K8R7	DQK)				
POLYETHYLENE GLYCOLS (UNI	: 3WJQ0SDW1A)				
MICRO CRYSTALLINE CELLULO	SE (UNII: OP1R32D61U)				

	: O8232NY3SJ)								
SODIUM STARCH GL	YCOLATE TYP	PE A POTATO (UNII: 5856J3G2A2	2)					
HYPROMELLOSES (U	JNII: 3NXW29V3	WO)							
STEARIC ACID (UNII:	4ELV7Z65AP)								
POVIDONE K30 (UNII	I: U725QWY32X)	1							
Droduct Characte	ristics								
Color	blue Score no score								
Shape	OVA	r				12mm			
-	0.111		Size			220			
Flavor Contains			Imprint Code		_				
Packaging									
# Item Code		Package Desc	ription		Marketing	g Start Date	Marketi	ng End Dat	
1 NDC:49035-762-24	24 in 1 BOTTLE	E; Type 0: Not a	Combination Prod	luct	05/25/2016				
2 NDC:49035-762-10	100 in 1 BOTTI	LE; Type 0: Not a	Combination Pro	duct	05/25/2016				
3 NDC:49035-762-21	225 in 1 BOTTI	LE; Type 0: Not a	Combination Pro	duct	05/25/2016				
4 NDC:49035-762-30	300 in 1 BOTT	LE; Type 0: Not a	a Combination Pro	oduct	05/25/2016				
Marketing Info	ormation								
•		on Number or N	Monograph Cita	ation	Marketing	g Start Date	Market	ing End Date	
Marketing Category			Monograph Cita	ation	Marketinş 0 5/25/20 16	g Start Date	Market	ing End Date	
Marketing Category	y Applicatio		Monograph Cita	ation		g Start Date	Market	ing End Date	
Marketing Category ANDA	y Application ANDA091353		Monograph Cita	ation		g Start Date	Marketi	ing End Date	
Marketing Category ANDA	y Application ANDA091353		Monograph Cita	ition		g Start Date	Market	ing End Date	
Marketing Category ANDA NAPROXEN So aproxen sodium tab	y Application ANDA091353 ODIUM olet		Monograph Cita	ation		g Start Date	Marketi	ing End Date	
Marketing Category ANDA	y Application ANDA091353 ODIUM olet						Marketi NDC:490		
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat	y Application ANDA091353 ODIUM olet tion	3			05/25/2016				
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type	y Application ANDA091353 ODIUM olet tion	HUMAN OTC E			05/25/2016				
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra	y Application ANDA091353 ODIUM olet tion	HUMAN OTC E ORAL			05/25/2016				
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra	y Application ANDA091353 ODIUM olet tion	HUMAN OTC E ORAL	DRUG :		05/25/2016	e)			
Marketing Category ANDA NAPROXEN So naproxen sodium tab Product Informat Product Type	y Application ANDA091353 ODIUM olet tion tion	HUMAN OTC E ORAL ety gredient Nam	PRUG	Ite m C	05/25/2016	e)	NDC:490	35-761	
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra Active Ingredient	y Application ANDA091353	HUMAN OTC E ORAL ety gredient Nam	PRUG	Ite m C	05/25/2016	e) Basis of	NDC:490	35-761 Strengtl	
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra Active Ingredient	y Application ANDA091353	HUMAN OTC E ORAL ety gredient Nam A3C) (NAPROXE	PRUG e EN - UNII:57Y76 R9	Ite m C	05/25/2016	e) Basis of	NDC:490	35-761 Strengtl 220 mg	
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra Active Ingredient NAPROXEN SODIUM Inactive Ingredie	y Application ANDA091353 ODIUM olet tion tion t/Active Moi Ing (UNII: 9TN87S33	HUMAN OTC E ORAL ety gredient Nam A3C) (NAPROXE	PRUG	Ite m C	05/25/2016	e) Basis of	NDC:490	35-761 Strengtl	
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra Active Ingredient NAPROXEN SODIUM Inactive Ingredie	y Application ANDA091353 ODIUM olet tion tion t/Active Moi Ing (UNII: 9TN87S3 nts	HUMAN OTC E ORAL ety gredient Nam A3C) (NAPROXE Ingred 2K)	PRUG e EN - UNII:57Y76 R9	Ite m C	05/25/2016	e) Basis of	NDC:490	35-761 Strengtl 220 mg	
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra Active Ingredient NAPROXEN SODIUM Inactive Ingredie FD&C BLUE NO. 2 (U	y Application ANDA091353	HUMAN OTC E ORAL ety gredient Nam A3C) (NAPROXE Ingred (X) WO)	e EN - UNII:57Y76 R9	Ite m C	05/25/2016	e) Basis of	NDC:490	35-761 Strengtl 220 mg	
Marketing Category ANDA NAPROXEN So haproxen sodium tab Product Informat Product Type Route of Administra Active Ingredient	y Application ANDA091353 ODIUM ole : tion tion t/Active Moi Ins (UNII: 9TN87S3 nts NII: L06K8R7DC JNII: 3NXW29V3 E CELLULOSE	HUMAN OTC E ORAL ety gredient Nam A3C) (NAPROXE Ingred k) WO) (UNII: OP1R32D6	e EN - UNII:57Y76 R9	Ite m C	05/25/2016	e) Basis of	NDC:490	35-761 Strengtl 220 mg	

STEARIC ACID (UNII:	,						
STARCH, CORN (UNI	II: O8232NY3SJ)						
SO DIUM STARCH GI	LYCOLATE TYPE A POTA	ATO (UNII: 5856J3G2A2)					
TITANIUM DIO XIDE	(UNII: 15FIX9V2JP)						
Product Characteristics							
Color	blue	Score	1	no score			
Shape	ROUND	Size		10 mm			
Flavor		Imprint Code	Imprint Code				
Contains							
Packaging							
# Item Code	Package	Package Description		Marketing End Date			
1 NDC:49035-761-21	225 in 1 BOTTLE; Type 0	: Not a Combination Product	05/25/2016				
Marketing Information							
Marketing Categor		er or Monograph Citation	Marketing Start Date	Marketing End Date			
ANDA	ANDA091353		05/25/2016				

Labeler - Wal-Mart Stores, Inc., (051957769)

Revised: 3/2017

Wal-Mart Stores, Inc.,