DG HEALTH INFANTS PAIN AND FEVER- acetaminophen suspension Dolgencorp, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dolgencorp, LLC Infants' Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart					
Weight (lb)	Age (yr)	Dose (mL)*			
under 24	under 2 years	ask a doctor			
24-35	2-3 years	5 mL			

* or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

- store at 20-25°C (68-77°F)
- do not use if printed bottle wrap is broken or missing

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-888-309-9030

Package/Label Principal Display Panel

DG[™] |health

Bonus! 1 OZ FREE

Compare to active ingredient of Infants' Tylenol® Oral Suspension

For Ages 2 to 3 Years

Infants' Pain & Fever

Acetaminophen 160 mg per 5 mL

Pain Reliever / Fever Reducer

Suspension Liquid

- Alcohol free
- Aspirin free
- Ibuprofen free

Pediatrician Preferred Dosing System

Use only with enclosed syringe.

See side panel for more information.

Grape Flavor

1 OZ + 1 OZ = 2 FL OZ (59 mL)

160 mg PER 5 mL



DG HEALTH INFANTS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:55910-946

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 ml			
Inactive Ingredients					
Ingredient Name		Strength			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)					
BUTYLPARABEN (UNII: 3QPI1U3FV8)					
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)					
CARRAGEENAN (UNII: 5C69YCD2YJ)					
D&C RED NO. 33 (UNII: 9DBA0SBB0L)					
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)					
GLYCERIN (UNII: PDC6A3C0OX)					
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)					
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)					
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679	OBS311)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)					
WATER (UNII: 059QF0KO0R)					
SODIUM BENZOATE (UNII: OJ245FE5EU)					
SORBITOL (UNII: 506T60A25R)					
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)					

Product Characteristics

Color	PURPLE	Score			
Shape		Size			
Flavor		Imprint Code			
Contains					

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:55910- 946-10	1 in 1 CARTON	11/19/2011	09/30/2022			
1		30 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product					
2	NDC:55910- 946-16	1 in 1 CARTON	11/19/2011				
2		59 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product					
Marketing Information							
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			

11/19/2011

Labeler - Dolgencorp, LLC (068331990)

Revised: 2/2022

Dolgencorp, LLC