

**STRATUSCARE ADULT GLYCERIN- glycerin suppository**  
**Stratus Pharmaceuticals**

-----  
**Stratuscare Adult Glycerin Suppositories**

***Drug Facts***

**Active ingredient (per suppository)**

Glycerin, USP 2 g

**Purpose**

Laxative

**Use**

- For relief of occasional constipation (irregularity)
- generally produces bowel movement in ¼ to 1 hour.

**Warnings**

**For rectal use only.**

May cause rectal discomfort or burning sensation.

**Ask a doctor before use if you have:**

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than 2 weeks.

**Stop use and ask a doctor:**

- if you have rectal bleeding or fail to have a bowel movement after using a laxative. This may indicate a serious condition.
- **Do not use** laxative products for longer than 1 week unless directed by a doctor
- **If pregnant or breastfeeding**, ask a doctor before use.

**Keep out of the reach of children.** If swallowed, get medical help or contact a Poison Control Center immediately (1-800-222-1212).

**Directions**

- **Adults and children 6 years and over:**
  - Carefully insert one suppository well into rectum
  - It need not melt completely to produce laxative action
  - Generally produces bowel movement in 15 minutes to 1 hour
  - Do not exceed one suppository daily unless directed by a doctor
- **Children under 6 years old:** do not use

**Other information**

- Store glycerin suppositories at room temperature, between 15°C and 30°C (between 59°F and 86°F)
- Keep container tightly closed

**Inactive ingredients**

purified water, sodium stearate

**Questions or comments?**

1-800-442-7882

*Distributed by:*

Stratus Pharmaceuticals Inc  
12379 SW 130th Street  
Miami, Florida 33186

**PRINCIPAL DISPLAY PANEL - 50 Suppository Jar Label**

**NDC 58980-410-50**

**STRATUSCARE**

**ADULT**

**Glycerin  
Suppositories  
Laxative**

*Distributed by:*

**STRATUS**  
PHARMACEUTICALS INC

**DO NOT USE IF INNER SEAL IS BROKEN OR MISSING.**

**50 SUPPOSITORIES**

**NDC 58980-410-50**

**STRATUSCARE**

**ADULT**  
Glycerin Suppositories Laxative

Distributed by:  
**STRATUS**  
PHARMACEUTICALS INC

**50 SUPPOSITORIES**

**Drug Facts**  
**Active ingredient (per suppository)**  
Glycerin, USP 2 g.....Laxative  
**Purpose**  
Laxative

**Use**  
• For relief of occasional constipation (irregularity) • generally produces bowel movement in ¼ to 1 hour.

**Warnings**  
For rectal use only.  
May cause rectal discomfort or burning sensation. Ask a doctor before use if you have: • abdominal pain, nausea or vomiting • a sudden change in bowel habits lasting more than 2 weeks. Stop use and ask a doctor: • if you have rectal bleeding or fail to have a bowel movement after using a laxative. This may indicate a serious condition. • Do not use laxative products for longer than 1 week unless directed by a doctor • If pregnant or breastfeeding, ask a doctor before use. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center immediately (1-800-222-1212).

**Directions** • Adults and children 6 years and over: • Carefully insert one suppository well into rectum • It need not melt completely to produce laxative action • Generally produces bowel movement in 15 minutes to 1 hour • Do not exceed one suppository daily unless directed by a doctor • Children under 6 years old: do not use

**Other information** • Store glycerin suppositories at room temperature, between 15°C and 30°C (between 59°F and 86°F) • Keep container tightly closed

**Drug Facts (continued)**  
**Inactive ingredients** purified water, sodium stearate

Questions or comments?  
1-800-442-7882

Distributed by:  
Stratus Pharmaceuticals Inc  
12379 SW 130th Street  
Miami, Florida 33186

GS25L 2013-70  
3 58980 41050 1

## STRATUSCARE ADULT GLYCERIN

glycerin suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58980-410
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Glycerin</b> (UNII: PDC6A3C00X) (Glycerin - UNII:PDC6A3C00X)	Glycerin	2 g

### Inactive Ingredients

Ingredient Name	Strength
<b>Water</b> (UNII: 059QF0K00R)	
<b>SODIUM STEARATE</b> (UNII: QU7E2XA9TG)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58980-410-12	12 in 1 JAR; Type 0: Not a Combination Product	05/30/2014	
2	NDC:58980-410-25	25 in 1 JAR; Type 0: Not a Combination Product	05/30/2014	
3	NDC:58980-410-50	50 in 1 JAR; Type 0: Not a Combination Product	05/30/2014	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	05/30/2014	

**Labeler** - Stratus Pharmaceuticals (789001641)

**Establishment**

Name	Address	ID/FEI	Business Operations
Tarmac Products		059890491	MANUFACTURE(58980-410)

Revised: 11/2024

Stratus Pharmaceuticals