

IBUPROFEN- ibuprofen tablet, film coated
NuCare Pharmaceuticals, Inc.

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- Temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- Temporarily reduces fever

Warnings

Allergy Alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
- hives
- facial swelling
- asthma (wheezing)
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- fever gets worse or lasts more than 3 days
- any new symptoms occur

If pregnant or breast-feeding,

ask a health professional before use. it is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose , get medical help or contact a Poison Control Center right away.

Directions

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over: take 1 tablet

- every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not take more than 6 tablets in 24 hours unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at controlled room temperature
- avoid excessive heat 40°C (104°F)

Inactive ingredients

carnauba wax, cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, PEG, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?

Call 1-800-540-3765

Package label

NuCare Pharmaceuticals, Inc.

NDC: 66267-115-20
Ibuprofen 200mg #20 Tablets

Each tablet contains: Ibuprofen USP 200mg (NSAID)*... Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing), rash, skin reddening, hives, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning. This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you, are age 60 or older, take more or for a longer time than directed, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Round Brown Tablet Debossed: "44 291" on one side

Product #: P0115020

WARNING: KEEP OUT OF REACH OF CHILDREN **STORE AT CONTROLLED TEMPERATURE 59-86°F.**

Ibuprofen 200mg
 Lot: 000000 NDC: 66267-0115-20
 MFR NDC: 57896-941-01 Exp.: 00-00

Ibuprofen 200mg
 Lot: 000000 NDC: 66267-0115-20
 MFR NDC: 57896-941-01 Exp.: 00-00

GTIN 00366267115209
 Serial# 0000000002
 Exp. Date 00-00
 LOT#: 000000

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Distributed by: Geri-Care Pharmaceuticals Corp., Brooklyn, NY 11204
 Packaged by: NuCare Pharmaceuticals, Inc., Orange, CA 92667

Take _____ every _____ hours
 _____ times a day.

Rev 01/01/19

IBUPROFEN

ibuprofen tablet, film coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|------------------------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:66267-115(NDC:57896-941) |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| LACTOSE (UNII: J2B2A4N98G) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | brown | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | 44291 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:66267-115-20 | 20 in 1 BOTTLE; Type 0: Not a Combination Product | 10/06/2016 | |
| 2 | NDC:66267-115-30 | 30 in 1 BOTTLE; Type 0: Not a Combination Product | 10/06/2016 | |
| 3 | NDC:66267-115-50 | 50 in 1 BOTTLE; Type 0: Not a Combination Product | 10/06/2016 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA075010 | 01/01/2004 | |

Labeler - NuCare Pharmaceuticals, Inc. (010632300)**Establishment**

| Name | Address | ID/FEI | Business Operations |
|------------------------------|----------------|---------------|----------------------------|
| NuCare Pharmaceuticals, Inc. | | 010632300 | repack(66267-115) |

Revised: 6/2024

NuCare Pharmaceuticals, Inc.