

**LORATADINE AND PSEUDOEPHEDRINE- loratadine and pseudoephedrine tablet, film coated, extended release**  
**SUPERVALU INC.**

-----  
**Loratadine and Pseudoephedrine**

***Drug Facts***

<b><i>Active ingredients (in each tablet)</i></b>	<b><i>Purpose</i></b>
Loratadine, USP 10 mg	Antihistamine
Pseudoephedrine sulfate, USP 240 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - itchy, watery eyes
  - runny nose
  - itching of the nose or throat
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product do not take more than directed.** Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

**Directions**

- do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- sodium:** contains 10 mg/tablet
- calcium:** contains 25 mg/tablet
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.**
- store between 20° C to 25° C (68° F to 77° F)
- protect from light and store in a dry place

**Inactive ingredients**

calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide

**Questions?**

call 1-877-932-7948

**DISTRIBUTED BY SUPERVALU INC.  
EDEN PRAIRIE, MN 55344 USA**

**PRINCIPAL DISPLAY PANEL - 10 Tablet Blister Pack Carton**

EQUALINE®  
NDC 41163-165-52

compare to  
Claritin-D®  
24 Hour  
active ingredients†

original prescription strength

allergy &  
congestion relief

loratadine, USP 10 mg (antihistamine)  
pseudoephedrine sulfate, USP 240 mg (nasal decongestant)  
indoor & outdoor allergies

non-drowsy\*

24 hour relief of:

- nasal & sinus congestion due to colds or allergies
- sneezing; runny nose; itchy, watery eyes;  
itchy throat or nose due to allergies

\*WHEN TAKEN AS DIRECTED. SEE DRUG FACTS PANEL.

10 extended-release tablets



5109799



6610915

**Drug Facts (continued)**

**Warnings**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product do not take more than directed.** Taking more than directed may cause drowsiness.

**Other Information**

- sodium: contains 10 mg/tablet
- calcium: contains 25 mg/tablet

**TAMPERING.**

- ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS

protect from light and store in a dry place  
store between 20° C to 25° C (68° F to 77° F)

**Directions**

- do not divide, crush, chew or dissolve the tablet
- adults and children 12 years and over: 1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
- children under 12 years of age: ask a doctor
- consumers with liver or kidney disease: ask a doctor

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Warnings**

- if pregnant or breastfeeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Drug Facts**

**Active ingredients (in each tablet)**

Loratadine, USP 10 mg, Pseudoephedrine sulfate, USP 240 mg, Antihistamine, Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: sneezing; itchy, watery eyes; runny nose; itching of the nose or throat

**EQUALINE**

NDC 41163-165-52

compare to Claritin-D® 24 Hour active ingredients!

# allergy & congestion relief

loratadine, USP 10 mg (antihistamine)  
pseudoephedrine sulfate, USP 240 mg (nasal decongestant)  
indoor & outdoor allergies

24 hour relief of: non-drowsy\*  
• nasal & sinus congestion due to colds or allergies  
• sneezing; runny nose; itchy, watery eyes;  
itchy throat or nose due to allergies

\*WHEN TAKEN AS DIRECTED. SEE DRUG FACTS PANEL.

## 10 extended-release tablets



Expiration Date:

Batch No.

Non Varnish Area

**Inactive ingredients** calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium citrate, sodium chloride, sodium chloride, sodium chloride, titanium dioxide

**Questions?** call 1-877-932-7948

**Keep the carton. It contains important information.**

**See end panel for expiration information.**

† All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Claritin-D®.

DISTRIBUTED BY SUPervalU INC. R0514  
EDEN PRAIRIE, MN 55344 USA  
Contact us at 1-877-932-7948, or  
www.supervalu-ourownbrands.com

0 41163 45510 8

**LORATADINE AND PSEUDOEPHEDRINE**

loratadine and pseudoephedrine tablet, film coated, extended release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41163-165
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE SULFATE	240 mg

**Inactive Ingredients**

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (1200000 MW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	white (white to Off-White)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	RX724
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-165-52	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	11/17/2004	

  

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076557	11/17/2004	

**Labeler** - SUPERVALU INC. (006961411)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

**Establishment**

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		051565745	manufacture(41163-165)

Revised: 8/2018

SUPERVALU INC.