

HEAD CONGESTION PLUS MUCUS PE- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated
HYVEE INC

1173-HYV-2025-1103

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and	▪ take 2 caplets every 4 hours
children	▪ do not take more than 10

12 years and over	caplets in 24 hours
children under 12 years	▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

HyVee®

NDC 42507-173-02

*Compare to the Active Ingredients in Sudafed PE® Head Congestion + Mucus

Non-Drowsy

Head Congestion + Mucus PE

Acetaminophen, Guaifenesin, Phenylephrine HCl

Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

For Relief of:

- Sinus Pressure
- Headache
- Chest Congestion

Actual Size

24 CAPLETS

Drug Facts (continued)

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, croscopolone,

Drug Facts (continued)

Questions or comments? 1-844-705-4384

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

With other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

After alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes ■ thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Directions

adults and children 12 years and over ■ take 2 caplets every 4 hours ■ do not take more than 10 caplets in 24 hours

children under 12 years ■ ask a doctor

Do not take more than directed (see overdose warning)

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use.

These could be signs of a serious condition:

- cough comes back or occurs with rash or headache that lasts
- new symptoms occur
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- nervousness, dizziness, or sleepiness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg Pain reliever/fever reducer

Guaifenesin 200 mg Expectorant

Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves these symptoms associated with

Drug Facts (continued)

may fever or other upper respiratory allergies, and the common cold:

- sinus congestion and pressure
- headache
- nasal congestion
- minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever



DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

SATISFACTION GUARANTEED

OR PURCHASE PRICE REFUNDED FOR INFORMATION CALL 1-800-289-8343

SmartLabel

Scan here for more information or call 1-800-289-8343

DISTRIBUTED BY HY-VEE, INC.

5820 WESTOWN PARKWAY
WEST DES MOINES, IA 50266
AAP40925

This product is not manufactured or distributed by Medial Consumer Healthcare, distributor of Sudafed PE® Head Congestion + Mucus.

Non-Drowsy

Head Congestion + Mucus PE

Acetaminophen, Guaifenesin, Phenylephrine HCl

Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

For Relief of:

- Sinus Pressure
- Headache
- Chest Congestion

24 CAPLETS

Actual Size

NDC 42507-173-02

***Compare to the Active Ingredients in Sudafed PE® Head Congestion + Mucus**

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:42507-173
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN 325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)		GUAIFENESIN 200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE 5 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	AAA;1173
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-173-02	2 in 1 CARTON	11/03/2025	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/03/2025	

Labeler - HYVEE INC (006925671)

Revised: 11/2025

HYVEE INC