

ACETAMINOPHEN 250 MG AND IBUPROFEN 125 MG- acetaminophen and ibuprofen tablet, film coated
Dr. Reddy's Laboratories Inc

Acetaminophen 250 mg and Ibuprofen 125 mg Tablets

Active ingredients (in each caplet)

Acetaminophen USP, 250 mg

Ibuprofen USP, 125 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purposes

Pain Reliever

Pain Reliever

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - muscular aches
 - minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in

people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

adults and children 12 years and over	<ul style="list-style-type: none"> • take 2 caplets every 8 hours while symptoms persist
children under 12 years	<ul style="list-style-type: none"> • ask a doctor

- do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

black iron oxide, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, glyceryl dibehenate, hypromellose, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, sodium lauryl sulfate, titanium dioxide

Questions or comments?

call **1-888-375-3784** Weekdays (8 AM to 8 PM EST)

Distributed By:

Dr. Reddy's Laboratories, Inc.


Princeton, NJ 08540

Made in India


Rev: 10/24



Principal Display Panel- Container

**Acetaminophen 250 mg and
Ibuprofen 125 mg Tablets
Pain Reliever**



**Back of Top Ply
(Page #2)**





Principal Display Panel- Container 250's count

Acetaminophen 250 mg and Ibuprofen 125 mg Tablets

Pain Reliever

Top Ply

R 0.1250"

165 mm

Back of Top Ply (Page #2)

Glue Area (22 mm)

165 mm

Bottom Ply

165 mm

58 mm

58 mm

58 mm

CONTAINS 2 MEDICINES
Acetaminophen+Ibuprofen

Acetaminophen and
Ibuprofen (NSAID)
Tablets, 250 mg/125 mg

PAIN RELIEVER

250 Caplets*
*Capsule-Shaped Tablets

TAMPER EVIDENT: Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage

Drug Facts

Active ingredients (in each caplet) **Purposes**
Acetaminophen USP 250 mg Pain Reliever
Ibuprofen USP 125 mg (NSAID)* Pain Reliever
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ toothache ■ backache
■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings
Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ with other drugs containing acetaminophen ■ more than 6 caplets in 24 hours, which is the maximum daily amount for this product ■ 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters ■ if an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever ■ right before or after heart surgery

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure

Distributed by: Dr. Reddy's Laboratories, Inc. Princeton, NJ 08540

1 5 0 0 X X X X X

Rev. 10/23

LOT
EXP
PULL HERE

Drug Facts (continued)

pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

When using this product
■ take with food or milk, if stomach upset occurs

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed

adults and children 12 years and over	• take 2 caplets every 8 hours while symptoms persist
children under 12 years	• ask a doctor.

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information


■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

black iron oxide, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, glyceryl dibehenate, hypromellose, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, sodium lauryl sulfate, titanium dioxide

Questions or comments?

call 1-888-375-3784 Weekdays (9 AM to 8 PM EST)



ACETAMINOPHEN 250 MG AND IBUPROFEN 125 MG

acetaminophen and ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43598-052
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	125 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XB4)	

CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
GLYCERYL DIBEHENATE (UNII: R8WTH25YS2)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	ACI
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-052-18	1 in 1 CARTON	02/01/2025	
1		18 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:43598-052-25	1 in 1 CARTON	02/01/2025	
2		250 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:43598-052-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	
4	NDC:43598-052-14	144 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	
5	NDC:43598-052-95	250 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA218247	02/01/2025	

Labeler - Dr. Reddy's Laboratories Inc (802315887)

Establishment

Name	Address	ID/FEI	Business Operations
Dr. Reddy's Laboratories Limited		918608162	manufacture(43598-052) , analysis(43598-052)

