HISTANAZ- pseudoephedrine hcl, pyrilamine maleate tablet Capital Pharmaceutical, LLC

Active ingredients (in each tablet) Pseudoephedrine HCl 60 mg Pyrilamine Maleate 30 mg

Uses

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- nasal congestion
- runny nose
- itchy watery eyes
- itching of the nose and throat

Questions? Comments?

Serious side effects may be reported to this number.

Call 1 (614) 638-4622

Decongestant, Antihistamine

Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)
- heart disease
- thyroid disease
- diabetes
- · trouble urinating due to enlargement of the prostate gland

Warnings

Do not exceed recommended dosage.

May cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages while taking this product. Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery.

Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your

prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Stop use and ask a doctor if

- cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Carnauba Wax, Croscarmellose Sodium, FD&C Yellow #6 Aluminum Lake, Iron Oxide Yellow, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Silicon Dioxide, Talc, Titanium Dioxide.


Directions:

Do not exceed recommended dosage.

Adults and Children 12 years of age and over:

Children 6-12 years of age:

1 tablet every 6/8 hours. Do not exceed 4 tablets in 24 hours. 1/2 tablet every 6/8 hours. Do not exceed 2 tablets in



24 hours.

Pseudoephedrine HCL

Pyrilamine Maleate



HISTANAZ

pseudoephedrine hcl, pyrilamine maleate tablet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:29978-129

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F) PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL) PYRILAMINE MALEATE 30 mg

Inactive Ingredients			
Ingredient Name	Strength		
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)			
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)			
POLYVINYL ALCOHOL (UNII: 532B59J990)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
TALC (UNII: 7SEV7J4R1U)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			

Product Characteristics					
Color	yellow	Score	no score		
Shape	OVAL	Size	13mm		
Flavor		Imprint Code	D129		
Contains					

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:29978-129- 10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/08/2025			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M012	09/08/2025			

Labeler - Capital Pharmaceutical, LLC (831545541)

Revised: 9/2025 Capital Pharmaceutical, LLC