

ANTI DIARRHEAL- loperamide hcl capsule
WinCo Foods

248 - Anti Diarrheal

Active ingredient(s)

Loperamide HCl 2mg

Purpose

Anti-diarrheal

Use(s)

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reactions to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

- if you have bloody or black stool

Ask a doctor before use if

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

Pregnancy/Breastfeeding

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose below. If possible, use weight to dose; otherwise, use age.
- adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
- children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
- children 6-8 years (48-59 lbs): 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
- children 2-5 years (34 to 47 lbs): ask a doctor
- children under 2 years (up to 33 lbs): do not use

Other information

store at 20°-25°C (68°-77° F)

avoid excessive heat above 40°C (104°F)

do not use if blister unit is broken or torn

Inactive ingredients

corn starch, D&C yellow #10
aluminum lake, dicalcium phosphate dihydrate, FD&C
blue #1 brilliant blue lake, magnesium stearate,
microcrystalline cellulose, silica gel

Questions

Call 1-800-824-1706 Monday - Friday 9AM - 4PM MST

Principal Display Panel



Anti Diarrheal

ANTI DIARRHEAL

loperamide hcl capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-405
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	CAPSULE	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-405-12	12 in 1 BLISTER PACK; Type 0: Not a Combination Product	03/01/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	03/01/2022	

Labeler - WinCo Foods (056098817)

Revised: 1/2022

WinCo Foods