

**MUCUS RELIEF COUGH COLD SORE THROAT- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
H E B**

Drug Facts

Active ingredients (in each 10 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough due to minor throat and bronchial irritation
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion
 - stuffy nose
 - sore throat
 - minor aches and pains
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product,

do not use more than directed (see Overdose warning).

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever get worse or lasts more than 3 days
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows or as directed by a doctor
- mL = milliliter
- **children 6 to under 12 years of age:** 10 mL in dosing cup provided
- **children under 6 years of age:** do not use

Other information

- **each 10 mL contains:** sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, disodium EDTA,, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call toll free: **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Children's Mucinex® Cold, Cough & Sore Throat active ingredients*

Children's

Mucus Relief

Cough, Cold & Sore Throat

Acetaminophen 325 mg / Pain reliever/Fever reducer

Dextromethorphan HBr 10 mg / Cough suppressant

Guaifenesin 200 mg / Expectorant

Phenylephrine HCl 5 mg / Nasal decongestant

Cold Relief

For Ages 6 to 11 years

Relief of:

- **Cough**
- **Sore Throat**
- **Nasal Congestion**
- **Chest Congestion**

Berry Flavor

FL OZ (mL)

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Cold, Cough & Sore Throat.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

MADE WITH PRIDE AND CARE FOR H-E-B®

SAN ANTONIO, TX 78204

Package Label

Compare to Children's Mucinex® Cold, Cough & Sore Throat active ingredients*

NDC 37808-811-40



Children's Mucus Relief Cough, Cold & Sore Throat

Acetaminophen 325 mg /
Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg /
Cough suppressant
Guaifenesin 200 mg / Expectorant
Phenylephrine HCl 5 mg /
Nasal decongestant

Cold Relief

For Ages 6 to 11 Years

Relief of:

- Cough
- Sore Throat
- Nasal Congestion
- Chest Congestion



Berry Flavor

4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion
 - stuffy nose
 - sore throat
 - minor aches and pains
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not use more than directed (see Overdose warning).

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows, or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call toll free: 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Cold, Cough & Sore Throat.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

PARENTS: Learn about teen medicine abuse. www.StopMedicineAbuse.org

MADE WITH PRIDE AND CARE FOR H-E-B®
SAN ANTONIO, TX 78204



0 41220 47278 9

PLD-A350C FC008427 38895-2304

Lot No.:

Exp. Date:

H-E-B Children's Mucus Relief Cough, Cold & Sore Throat

MUCUS RELIEF COUGH COLD SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:37808-811

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-811-40	1 in 1 BOX	04/30/2023	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	04/30/2023	

Labeler - H E B (007924756)

Revised: 4/2024

H E B