DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMPTOM- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl ASSURED / DOLLAR TREE (Greenbrier International, Inc.)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients in Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients in Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - headache
 - minor aches and pains
 - fever
 - sore throat

NIGHTTIME

- temporarily relieves common cold and fly symptoms
 - o cough due to minor throat and bronchial irritation
 - sore throat

- headache
- minor aches and pain
- fever
- runny nose and sneezing

Warnings

DAYTIME and NIGHTTIME

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

DAYTIME and NIGHTTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occur with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

• taking the blood thinning drug warfarin

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product,

DAYTIME

do not exceed recommended dosage.

NIGHTTIME

- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

DAYTIME and NIGHTTIME

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

DAYTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- children under 12 years : do not use

Nighttime:

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or disolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other information

DAYTIME and NIGHTTIME

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

Daytime butylated hydroxyanisole, butylated hydroxytoluene, carminic acid*, D&C yellow #10*, edible white ink, FD&C red #40*, FD&C yellow #6*, gelatin, glycerin, polyethylene glycol*, povidone, propylene glycol, purified water, sodium metabisulfite*, sorbitan*, sorbitol

*may contain this ingredient

Nighttime D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan*, sorbitol

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL® AND NYQUIL® LIQUICAPS®**

Multi-Symptom Daytime Liquid Capsules

Acetaminophen - Pain Reliever / Fever Reducer

Dextromethorphan HBr - Cough Suppressant

Phenylephrine HCl - Nasal Decongestant

SOFTGELS

Combo Pack

NIGHTTIME

MULTI-SYMPTOM Nite Time

Liquid Capsules

Acetaminophen - Pain Reliever / Fever Reducer

^{*}may contain this ingredient

Dextromethorphan HBr - Cough Suppressant

Doxylamine Succinate - Antihistamine

SOFTGELS

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

**This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquiCap® are registered trademarks of The Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY

GREENBRIER INTERNATIONAL, INC.

500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

Product Label

and Gamble Company "This product is not manufactured or distributed by The Procter & Gamble Company Vicks®, DayQuil®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter

KEEP OUTER CARTON FOR COMPLETE WARNINGS

AND PRODUCT INFORMATI

JAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLIST UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

■ do not take more than 4 doses in 24 hours
■ swallow whole; do not crush, Directions

do not take more than directed (see Overdose warning) adults as well as for children even if you do not notice any signs or symptoms. recommended dose can cause liver damage. In case of overdose, get medical help or contact a Polson Control Center right away. Quick medical attention is critical for Keep out of reach of children. Overdose warning: Taking more than the If pregnant or breast-feeding, ask a health professional before use. ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days

■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. Stop use and ask a doctor if

mervousness, dizziness, or sleeplessness occur When using this product, do not use more than directed.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug ■ bersistent or chronic cough such as occurs with smoking, asthma, or emphysema too much phiegm (mucus) Trouble uninating due to an enlarged prostate gland Ask a doctor before use if you have in liver disease in bear disease in the plood preserve in the disease in the plood preserve in t

The compression of the control of th Do not use with any other drug containing acetaminophen (prescription or

ever, headache, rash, nausea, or vomiting, consult a doctor promptly. f sore throat is severe, persists for more than 2 days, is accompanied or followed by eactions. Symptoms may include: **m** skin reddening **m** blisters **m** rash, if a skin reddening **m** rash, if a skin reddening include; **m** rash, if a skin reddening include; **m** rash, if a skin rash, if a ski occur if you take:

with a nore than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using Jilb product. A libergy alert, beginnophen may cause severe skin will be used to the product of the containing a celebration and the page. Liver warning: This product contains acetaminophen. Severe liver damage may

Drug Facts (continued) Daytime Cold & Flu Softgel (continued)

■ headache ■ minor aches and pains ■ fever ■ sore throat ■ condu que to minor taroat and bronchiai irritation ■ nasai congestion nage = remporanty relieves common cold and thu symptoms

Nasal decongestant Рћепујерћиће НСІ 5 mg. Cough suppressant Dextromethorphan HBr 10 mg. səsodınd Active ingredients (in each softgel)

SSURE

Liquid Capsules

Phenylephrine HCI - Nasal Decongestant

Fever, Headache, Sore Throat, Coughing, Minor Aches and Pains, & Nasal Congestion

Acetaminophen - Pain Reliever / Fever Reducer Dextromethorphan HBr - Cough Suppressant

MULTI-SYMPTOM

Combo Pack

12count

Daytime Cold & Flu Softgel Drug Facts

adults as well as for children even if you do not notice any signs or symptoms. 🔻 or contact a Poison Control Center right away. Quick medical attention is critical for If pregnant or breast-feeding, sak a health professional before use. Keep out of reach of children. Overdose warming: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help

■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ■ fever gets worse or lasts more than 3 days
■ redness or swelling is present Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days

vehicle or operating machinery

and tranquillizers may increase drowsiness

De careful when driving a motor When using this product — excitability may occur, especially in children — marked drowsiness may occur — avoid slcoholic drinks — majcohol, sedatives,

drug warfarin # taking sedatives or tranquilizers Ask a doctor or pharmacist before use if you are at taking the blood thinning

trouble urinating due to an enlarged prostate gland occurs with too much phiegm (mucus) s s breathing problem or chronic cough that last such as occurs with smooting, asthms, chronic bronchilds, or emphysems

Ask a doctor before use if you have liver disease glaucoma cough that

if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking nonprescription). If you are not sure whether a drug contains scetaminophen, sak a doctor or pharmacist.

Inhibitor (MAOI) (leortain drugs for depression, psychiatinc, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your trescription that manager is the parking if your trescription that manager is the parking. Do not use ■ with any other drug containing acetaminophen (prescription or

headache, rash, nausea, or vomiting, consult a doctor promptly. Symptoms may include: ■ skin reddening ■ bilsters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore curs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, Liver warming: This product contains acelaminophen. Severe liver damage may occur if you take:

— more than 4,000 mg of acetaminophen in Z4 hours — with other drugs containing acetaminophen — 3 or more alcoholic drinks every day while using this product. Affergy after Acetaminophen may cause severe askin reactions "Emonthma may include: — a skin reaction — includers — may a higher passing a Semonthma may include: — a skin reaction.

Drug Facts (continued) Nighttime Cold & Flu Softgel (continued)

minor aches and pains ■fever ■runny nose & sneezing ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache naes - remporanly relieves common cold and thu symptoms

Antihistamine Doxylamine succinate 6.25 mg. Cough suppressant Dextromethorphan HBr 15 mg. səsodinə Active ingredients (in each softgel)

> Nighttime Cold & Flu Softgel Drug racts

COMPARE TO ACTIVE INGREDIENTS IN VICKS® DAYQUIL® & VICKS® NYQUIL®

Cold & Flu LiquiCaps®**

NDC 33992-0472-2

MULTI-SYMPTOM

lite i **Liquid Capsules**

- Acetaminophen Pain Reliever / Fever Reducei
 Dextromethorphan HBr Cough Suppressant
- · Doxylamine Succinate Antihistamine

Fever, Headache, Sore Throat, Coughing, Minor Aches and Pains, Runny Nose & Sneezing

Actual Size

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

4 softgels

,way contain this ingredient Inactive ingredients D&C yellow #10, edible white ink FD&C blue #1, gelatin, applyenn, purified water, sorbital glycol, purified glycol, purif

Other information a store between 15-30°C (59-86°F) a avoid excessive heat

water every 6 hours - Children under 12 years; do not use ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ≡ a bulbs and children 12 years and over take 2 softgels with Directions

do not take more than directed (see Overdose warning)

500 VOLVO PARKWAY, CHESAPEAKE, VA 23320 distributed by Greenbrier International, Inc

Lot No.:

-00

contains one or more of these ingredients odium metabisulitie", sorbitan", sorbitol sodium metabisulitie", sorbitan", sorbitol carminic acid*, D&C yellow #10*, edible white ink, FD&C red #40*, FD&C yellow #6*, **jusctive ingredients** butylated hydroxyanisole*, butylated hydroxytoluene*

Other information a store between 15-30°C (59-86°F) avoid excessive heat

vater every 4 hours ■ children under 12 years: do not use shew, or dissolve adults and children 12 years and over: take 2 softgels with

Drug Facts (continued) Daytime Cold & Flu Softgel (continued)

8 softgels

Drug Facts (continued) Nighttime Cold & Flu Softgel (continued)

ASSURED Daytime and Nighttime Liquid Capsules

DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMPTOM

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:33992-0472

Packaging

| # | t Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|-----------------------------|--------------------|
| 1 | NDC:33992-0472-2 | 1 in 1 CARTON; Type 0: Not a Combination Product | 05/31/2017 | 12/31/2021 |

Quantity of Parts

| Quan | Quantity of Farts | | | |
|--------|-------------------|------------------------|--|--|
| Part # | Package Quantity | Total Product Quantity | | |
| Part 1 | 4 BLISTER PACK | 4 | | |
| Part 2 | 8 BLISTER PACK | 8 | | |

Part 1 of 2

NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|----------|
| ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D) | ACETAMINOPHEN | 325 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 15 mg |
| | | |

DOXYLAMINE SUCCINATE (UNII: V9 B19 B5 Y12) (DOXYLAMINE - UNII: 95QB77JKPL) DOXYLAMINE SUCCINATE 6.25 mg

| Ingredient Name | Strength | |
|---|----------|--|
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | | |
| GELATIN (UNII: 2G86QN327L) | | |
| GLYCERIN (UNII: PDC6A3C0OX) | | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | | |
| PO VIDO NE (UNII: FZ989 GH94E) | | |

| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
|-------------------------------------|--|
| WATER (UNII: 059QF0KO0R) | |
| SORBITAN (UNII: 6O92ICV9RU) | |
| SORBITOL (UNII: 506T60A25R) | |

| Product Characteristics | | | |
|-------------------------|---------|--------------|------------------|
| Color | GREEN | Score | no score |
| Shape | CAPSULE | Size | 20 mm |
| Flavor | | Imprint Code | P30;94A;215;P120 |
| Contains | | | |

| F | Packaging | | | | |
|---|-----------|--|-----------------------------|---------------------------|--|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | |
| 1 | | 4 in 1 CARTON | | | |
| 1 | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | | |

| Marketing Information | | | |
|-----------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC MONOGRAPH FINAL | part341 | 05/31/2017 | 12/31/2021 |

Part 2 of 2

DAYTIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule

Product Information

Route of Administration ORAL

| Active Ingredient/Active Moiety | | |
|--|----------------------------------|----------|
| Ingredient Name | Basis of Strength | Strength |
| ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D) | ACETAMINOPHEN | 325 mg |
| DEXTRO METHO RPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH) (DEXTRO METHO RPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6 MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg |

| Inactive Ingredients | | |
|---|----------|--|
| Ingredient Name | Strength | |
| BUTYLATED HYDRO XYANISO LE (UNII: REK4960 K2U) | | |
| BUTYLATED HYDRO XYTO LUENE (UNII: 1P9 D0 Z171K) | | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | | |

| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
|---|--|
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| PO VIDO NE (UNII: FZ989 GH94E) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITAN (UNII: 6O92ICV9RU) | |
| SORBITOL (UNII: 506T60A25R) | |

| Product Characteristics | | | | | | |
|-------------------------|---------|--------------|------------------|--|--|--|
| Color | ORANGE | Score | no score | | | |
| Shape | CAPSULE | Size | 20 mm | | | |
| Flavor | | Imprint Code | P19;95A;512;P119 | | | |
| Contains | | | | | | |

| | Packaging | | | | | | |
|----|-------------|--|-----------------------------|--------------------|--|--|--|
| II | # Item Code | Package Description | Marketing Start Date | Marketing End Date | | | |
| П | 1 | 8 in 1 CARTON | | | | | |
| | 1 | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | | | | |

| Marketing Inform | Marketing Information | | | |
|-------------------------|--|----------------------|--------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC MONOGRAPH FINAL | part341 | 05/31/2017 | 12/31/2021 | |

| Marketing Information | | | | | |
|-----------------------|--|----------------------|--------------------|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | |
| OTC MONOGRAPH FINAL | part341 | 05/31/2017 | 12/31/2021 | | |
| | | | | | |

Labeler - ASSURED / DOLLAR TREE (Greenbrier International, Inc.) (610322518)

Revised: 10/2018 ASSURED / DOLLAR TREE (Greenbrier International, Inc.)