

**Giltuss D**

**Purposes**

## **Drug Facts**

<b>Active ingredients (in each 10 ml)</b>	<b>Purposes</b>
Glorpheniramine maleate, 3.5 mg.....	Antihistamine
Phenylephrine HCl, 10 mg .....	Nasal decongestant

**Uses** Temporarily relieves these Symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose • Sneezing • Itchy, watery eyes • nasal congestion
- itching of the nose or throat • sinus congestion and pressure

## **Warnings**

**Do not use** • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have:**

- heart disease • high blood pressure • thyroid disease • diabetes
- trouble urinating due to an enlarged prostate gland • glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema



**Ask a doctor or pharmacist before use if you are**  
• taking sedatives or tranquilizers

#### Purposes

Antihistamine

Nasal decongestant

#### Uses

Temporary relieves these symptoms due to hay fever (allergy rhinitis) or other respiratory allergies:

runny nose, sneezing, itchy, watery eyes, nasal congestion, itching of nose or throat, sinus congestion and pressure

#### Warnings

Warnings:

Do not use to sedate a child or to make a child sleepy, if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## Drug Facts

**Active ingredients (in each 10 ml)**

**Purposes**

Chlorpheniramine maleate, 3.5 mg..... Antihistamine

Phenylephrine HCl, 10 mg ..... Nasal decongestant

## Uses

Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose • Sneezing • Itchy, watery eyes • nasal congestion
- itching of the nose or throat • sinus congestion and pressure



## Warnings

**Do not use** • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have:

- heart disease • high blood pressure • thyroid disease • diabetes
- trouble urinating due to an enlarged prostate gland • glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers

### Ask a doctor

Ask a doctor before use if you have:

Heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, glaucoma, a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

## Drug Facts

**Active ingredients (in each 10 ml)**

**Purposes**

Chlorpheniramine maleate. 3.5 mg..... Antihistamine



Phenylephrine HCl, 10 mg .....Nasal decongestant

---

**Uses** Temporarily relieves these Symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose • Sneezing • Itchy, watery eyes • nasal congestion
  - itching of the nose or throat • sinus congestion and pressure
- 

## **Warnings**

**Do not use** • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

---

**Ask a doctor before use if you have:**

- heart disease • high blood pressure • thyroid disease • diabetes
  - trouble urinating due to an enlarged prostate gland • glaucoma
  - a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- 

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers

### **Ask a doctor or pharmacist**

Ask a doctor or pharmacist before use if you are, taking sedatives or tranquilizers



# Drug Facts

Active ingredients (in each 10 ml)

Purposes

Chlorpheniramine maleate, 3.5 mg.....	Antihistamine
Phenylephrine HCl, 10 mg .....	Nasal decongestant

**Uses** Temporarily relieves these Symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose • Sneezing • Itchy, watery eyes • nasal congestion
- itching of the nose or throat • sinus congestion and pressure

## Warnings

**Do not use** • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have:**

- heart disease • high blood pressure • thyroid disease • diabetes
- trouble urinating due to an enlarged prostate gland • glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers



## When using

When using this product, do not use more than directed, may cause drowsiness, avoid alcoholic beverages, alcohol, sedatives, and tranquilizers may increase drowsiness, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery, excitability may occur, especially in children.

**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



MANUFACTURED FOR  
**GIL PHARMACEUTICAL CORP.**  
PONCE, PUERTO RICO 00717-1565

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

## **Inactive ingredients**

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

## **Questions or comments?**

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.



## Pregnancy or breast-feeding

If Pregnancy or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away ( 1-800-222-1222)

**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



MANUFACTURED FOR  
GIL PHARMACEUTICAL CORP.  
PONCE, PUERTO RICO 00717-1565

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

## Inactive ingredients

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

## Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.



## Stop use section

Stop use and ask a doctor if, if you get nervous, dizzy or sleepless, symptoms do not get better withi

hin 7 days or are accompanied by fever

**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



MANUFACTURED FOR  
GIL PHARMACEUTICAL CORP.  
PONCE, PUERTO RICO 00717-1565

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

## Inactive ingredients

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

## Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.



# Drug Facts

**Active ingredients (in each 10 ml)**

**Purposes**

Chlorpheniramine maleate, 3.5 mg..... Antihistamine  
Phenylephrine HCl, 10 mg ..... Nasal decongestant

**Uses** Temporarily relieves these Symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose • Sneezing • Itchy, watery eyes • nasal congestion
- itching of the nose or throat • sinus congestion and pressure

## Warnings

**Do not use** • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.


**Ask a doctor before use if you have:**

- heart disease • high blood pressure • thyroid disease • diabetes
- trouble urinating due to an enlarged prostate gland • glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers





Chlorpheniramine maleate 3.5 mg

Phenylephrine HCl, 10 mg

**Directions**

Directions, not take more than 6 doses in any 24 hours period, use enclosed dosing cup, keep dosage cup for use with this product, ml= milliliter

Adults and children 12 years and over 10 ml every 4 hours

Children 6 to under 12 years 5 ml every 4 hours

Children under 6 years Consult a doctor



**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



MANUFACTURED FOR  
**GIL PHARMACEUTICAL CORP.**  
PONCE, PUERTO RICO 00717-1565

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

### **Inactive ingredients**

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

### **Questions or comments?**

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

### **Other information**

Other information store at room temperature 15 - 30 oC (59-86 o F), close cap tightly, save carton for complete drug facts



**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



MANUFACTURED FOR  
**GIL PHARMACEUTICAL CORP.**  
PONCE, PUERTO RICO 00717-1565

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

### **Inactive ingredients**

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

### **Questions or comments?**

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

### **Inactive ingredients**

#### Inactive Ingredients

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified water, Sodium Carboxymethylcellulose, Sucralose.



**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



**MANUFACTURED FOR**  
**GIL PHARMACEUTICAL CORP.**  
**PONCE, PUERTO RICO 00717-1565**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

### ***Inactive ingredients***

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

### ***Questions or comments?***

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

### **Questions or Comments?**

Call 1-787-848-9114 Monday-Friday 9AM-5PM EST, Call your doctor for medical advice in the event of side effects



**When using this product** • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children



MANUFACTURED FOR  
**GIL PHARMACEUTICAL CORP.**  
PONCE, PUERTO RICO 00717-1565

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

### **Inactive ingredients**

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

### **Questions or comments?**

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

## **Indication and usage section**

Indication and usage section

Do not take more than 6 doses in any 24 hours period, use enclosed dosing cup, keep dosage cup for use with this product,



ml= milliliter

Adults and children 12 years and over 10 ml every 4 hours

Children 6 to under 12 years 5 ml every 4 hours

Children under 6 years Consult a doctor

**Drug Facts**

Active Ingredients (in each 10 mL) Purposes  
Chlorpheniramine maleate, 3.5 mg. Antihistamine  
Phenylephrine HCl, 10 mg. Nasal decongestant

**Uses** Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:  
• runny nose • Sneezing • Itchy, watery eyes • nasal congestion  
• itching of the nose or throat • sinus congestion and pressure

**Warnings**  
Do not use • to sedate a child or to make a child sleepy • if you are taking a prescription medication: cardiac inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
Ask a doctor before use if you have:  
• heart disease • high blood pressure • thyroid disease • diabetes  
• trouble urinating due to an enlarged prostate gland • glaucoma  
• a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
Ask a doctor or pharmacist before use if you are  
• taking sedatives or tranquilizers

**NO PSEUDOEPHEDRINE**  
**PSEUDOEPHEDRINE**

**G**

**Giltuss-D**  
**ALLERGY & CONGESTION**  
**ALERGIA Y CONGESTIÓN**

**DYE FREE. PRESERVATIVE FREE. SODIUM FREE. SUGAR FREE**  
**SIN COLORANTES. SIN PRESERVATIVOS. LIBRE DE SODIO. SIN AZÚCAR**

SNEEZING, RUNNY NOSE, ITCHY & WATERY EYES,  
ITCHY THROAT OR NOSE, SINUS PRESSURE, STUFFY NOSE  
ESTORNIDOS, GOTO NASAL, PICOR Y OJOS LLOROSOS, PICOR DE  
GARGANTA Y O NARIZ, PRESIÓN POR SINUSOS, NARIZ TAPADA.

**TUTTI FRUITI FLAVOR**  
**SABOR A TUTTI FRUITI**

**4 FL. OZ. (118 mL)**

When using this product • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children  
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).  
Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever  
**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosing cup for use with this product • mL, = milliliter  
Adults and children 12 years and over 10 mL every 4 hours  
Children 6 to under 12 years 5 mL every 4 hours  
Children under 6 years Consult a doctor  
**Other information** • store at room temperature 15°-30°C (59°-86°F) • close cap tightly • SEE CARTON FOR COMPLETE DRUG FACTS  
**Inactive ingredients**  
Citric Acid, Flaxseed, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.  
**Questions or comments?**  
Call 1-757-548-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

MANUFACTURED FOR  
**GL**  
GIL PHARMACEUTICAL CORP.  
PONCE, PUERTO RICO 00717-1565

Keep out of reach of children.



### Drug Facts

**Active ingredients (in each 10 mL)**

Chlorpheniramine maleate, 3.5 mg	Antihistamine
Phenylephrine HCl, 10 mg	Nasal decongestant

**Uses** Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- nasal congestion
- itching of the nose or throat
- sinus congestion and pressure

**Warnings**

Do not use • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers

**NO PSEUDOEPHEDRINE**  
**PSEUDOEFEEDRINA**

**Giltuss-D**  
**ALLERGY & CONGESTION**  
**ALERGIA Y CONGESTION**

**DYE FREE. PRESERVATIVE FREE. SODIUM FREE. SUGAR FREE**  
**SIN COLORANTES. SIN PRESERVATIVOS. LIBRE DE SODIO. SIN AZÚCAR**

SNEEZING, RUNNY NOSE, ITCHY & WATERY EYES,  
ITCHY THROAT OR NOSE, SINUS PRESSURE, STUFFY NOSE,  
ESTORNIDOS, GOTO NASAL, PICOR Y OJOS LLOROSOS, PICOR DE  
GARGANTA Y O NASAL, PRESIÓN POR SINUSOS, NASAL BARRIDA

**TUTTI FRUITI FLAVOR**  
**SABOR A TUTTI FRUITI**

**4 FL. OZ. (118 mL)**

When using this product • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever

**Directions** • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosing cup for use with this product • mL, = milliliter

Adults and children 12 years and over	10 mL, every 4 hours
Children 6 to under 12 years	5 mL, every 4 hours
Children under 6 years	Consult a doctor

**Other information** • store at room temperature 15°-30°C (59°-86°F) • close cap tightly • SEE CARTON FOR COMPLETE DRUG FACTS

**Inactive ingredients**  
Citric Acid, Flavor, Glycine, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

**Questions or comments?**  
Call 1-787-648-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

**MANUFACTURED FOR**  
**GIL PHARMACEUTICAL CORP.**  
**PONCE, PUERTO RICO 00717-1565**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away ( 1-800-222-1222)







Giltuss D

Allergy & Congestion

Dye Free, preservative free, sodium free, sugar free

Sneezing, Runny Nose, Itchy & Water eyes, Itchy Throat or Nose, Sinus pressure, stuffy nose

Tutti Frutti Flavor

\$ floz 118 ml

## GILTUSS D

giltuss d syrup

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-010
Route of Administration	ORAL		



Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)		CHLORPHENIRAMINE MALEATE	3.5 mg in 10 mL	
Inactive Ingredients				
Ingredient Name			Strength	
CITRIC ACID (UNII: 2968PHW8QP)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
WATER (UNII: 059QF0KO0R)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
Product Characteristics				
Color		Score		
Shape		Size		
Flavor	TUTTI FRUTTI	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65852-010-04	1 in 1 CARTON	04/11/2024	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		04/11/2024	

**Labeler** - Dextrum Laboratories Inc. (007392322)