

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 1- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 2- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 3- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 4- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION ROSE 2- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION ROSE 3- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION ROSE 4- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 2- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 3- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 4- octinoxate and titanium dioxide liquid
SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 5- octinoxate and titanium dioxide liquid
SHISEIDO AMERICAS CORPORATION

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
OCTINOXATE 2.9%	Sunscreen
TITANIUM DIOXIDE 2.4%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure

- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

Inactive Ingredients

DIMETHICONE•WATER•TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE•CAPRYLIC/CAPRIC TRIGLYCERIDE•DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER•PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE•SD ALCOHOL 40-B•MAGNESIUM MYRISTATE•NYLON-12•DIPHENYL DIMETHICONE/VINYL DIPHENYL DIMETHICONE/SILSESQUIOXANE CROSSPOLYMER•DISTEARDIMONIUM HECTORITE•POLYQUATERNIUM-51•THYMUS SERPILLUM EXTRACT•ALUMINUM HYDROXIDE•ALUMINUM DISTEARATE•POLYSILICONE-2•GLYCERIN•BHT•ALCOHOL•BUTYLENE GLYCOL•TOCOPHEROL•METHYL METHACRYLATE CROSSPOLYMER•HYPERICUM ERECTUM FLOWER/LEAF/STEM EXTRACT•TALC•DIMETHICONE/PEG-10/15 CROSSPOLYMER•SILICA•ALUMINA•POLYMETHYLSILSESQUIOXANE•DIISOSTEARYL MALATE•GLYCERYL DIISOSTEARATE•DIPROPYLENE GLYCOL•PHENOXYETHANOL•FRAGRANCE•TITANIUM DIOXIDE•IRON OXIDES•MICA•RED 27•

Other information

- protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Neutral 1

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Neutral 1

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRIFLUOROPROPYLDIMETHYL
 TRIMETHYLSILOXYMETHYL
 TRIMETHYLSILOXYMETHYL
 CAPRYLIC/CAPRIC TRIGLYCERIDE
 DIMETHICONE/VINYL DIMETHICONE
 CROSSPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8
 MAGNESIUM MYRISTATE - NYLON-12
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 CROSSPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYQUATERNIUM-51 -
 THYMUS SERPILLUM EXTRACT -
 ALUMINUM HYDROXIDE -
 ALUMINUM DISTEARATE -
 POLYSILICONE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 CROSSPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 CROSSPOLYMER -
 SILICA - ALUMINA
 POLYMETHYLSILSESQUOXANE -
 DIMETHYLSILOXYMETHYL
 GLYCERYL DIBESTEARATE -
 DIPHENYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
CORPORATION, DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

READ INSERTION



13119-40-5041

AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Neutral 2

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Neutral 2

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active Ingredients Purpose
OCINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - TRIMETHYLSILOXYDIMETHYLTRIMETHYLSILOXYLICATE - CAPRYLIC/CAPRIC TRIGLYCERIDE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - PEG-8 POLYDIMETHYLSILOXYETHYL DIMETHICONE - SD ALCOHOL 40-8 - MAGNESIUM MYRISTATE - NYLON-12 - DIPHENYL DIMETHICONE/VINYL DIPHENYL DIMETHICONE/SILSESQUOXANE CROSSPOLYMER - DISTEARDIMONIUM HECTORITE - POLYOLATERIUM-51 - THYMUS SERPYLLUM EXTRACT - ALUMINIUM HYDROXIDE - ALUMINIUM DISTEARATE - POLYSILICONE-2 - GLYCERIN - BHT - ALCOHOL - BUTYLENE GLYCOL - TOCOPHEROL - METHYL METHACRYLATE CROSSPOLYMER - HYPERICUM ERECTUM FLOWER/LEAF/STEM EXTRACT - TALC - DIMETHICONE/PEG-10/15 CROSSPOLYMER - SILICA - ALUMINA - POLYMETHYLSILSESQUOXANE - DISOCTEARYL MALATE - GLYCERYL DIISOSTEARATE - DIPROPYLENE GLYCOL - PHENOXYETHANOL - FRAGRANCE - TITANIUM DIOXIDE - IRON OXIDES - MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
CORPORATION, DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

READ INSERTION



10-5011
AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Neutral 3

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Neutral 3

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRIFLOROFLUOROMETHYL DIMETHYL
 TRIMETHYLSILOXY SILICATE -
 CAPRYLIC/CAPRIC TRIGLYCERIDE -
 DIMETHICONE/VINYL DIMETHICONE
 CROSSPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8 -
 MAGNESIUM MYRISTATE - NYLON-12 -
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 CROSSPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYQUATERNIUM-51 -
 THYMUS SERPYLLUM EXTRACT -
 ALUMINUM HYDROXIDE -
 ALUMINUM DISTEARATE -
 POLYSILICONE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 CROSSPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 CROSSPOLYMER -
 SILICA - ALUMINA -
 POLYMETHYLSILSESQUOXANE -
 DIMETHACRYL MALATE -
 GLYCERYL DIISOSTEARATE -
 DIPROPYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



SYNCHRO SKIN

Lasting Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

READ INSERTION.



SHISEIDO AMERICAS CORPORATION, DIST. NEW YORK, NY 10022 MADE IN U.S.A. www.shiseido.com

40-5041 AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Neutral 4

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Neutral 4

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - TRIFLUROPROPYLENEDIMETHYL TRIMETHYLSILOXYLICATE - CAPRYLIC/CAPRIC TRIGLYCERIDE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - PEG-8 POLYDIMETHYLSILOXYETHYL DIMETHICONE - SD ALCOHOL 40-8 - MAGNESIUM MYRISTATE - NYLON-12 - DIPHENYL DIMETHICONE/VINYL DIPHENYL DIMETHICONE/SILSESQUOXANE CROSSPOLYMER - DISTEARDIMONIUM HECTORITE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - ALUMINUM HYDROXIDE - ALUMINUM DISTEARATE - POLYSILICONE-2 - GLYCERIN - BHT - ALCOHOL - BUTYLENE GLYCOL - TOCOPHEROL - METHYL METHACRYLATE CROSSPOLYMER - HYPERICUM ERECTUM FLOWER/LEAF/STEM EXTRACT - TALC - DIMETHICONE/PEG-10/15 CROSSPOLYMER - SILICA - ALUMINA - POLYMETHYLSILSESQUOXANE - DIMETHACRYL MALATE - GLYCERYL DIISOSTEARATE - DIPROPYLENE GLYCOL - PHENOXYETHANOL - FRAGRANCE - TITANIUM DIOXIDE - IRON OXIDES - MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
CORPORATION, EAST
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

READ INSERTION



40-5041
AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Rose 2

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Rose 2

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRIMETHYLOLPROPYLEMETHYL
 TRIMETHYLSILOXYLICATE -
 CAPRYLIC/CAPRIC TRIGLYCERIDE -
 DIMETHICONE/VINYL DIMETHICONE
 COPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8 -
 MAGNESIUM MYRISTATE - NYLON-12 -
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 COPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYOLATERANIUM-SI -
 THYMUS SERPILLUM EXTRACT -
 ALUMINIUM HYDROXIDE -
 ALUMINIUM DISTEARATE -
 POLYSILOXANE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 COPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 COPOLYMER -
 SILICA - ALUMINA -
 POLYMETHYLSILSESQUOXANE -
 DIOCTYLARYL MALATE -
 GLYCERYL DIBESTEARATE -
 DIPROPYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
 Lasting
 Liquid Foundation
 Oil Free
 BROAD SPECTRUM
 SPF 20
 Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
 CORPORATION, DIST.
 NY, NY 10022
 MADE IN U.S.A.
 www.shiseido.com

READ INSERTION



40-5041
 AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Rose 3

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Rose 3

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRIMETHYLSILOXYDIMETHYL
 TRIMETHYLSILOXYDIMETHYL
 TRIMETHYLSILOXYDIMETHYL
 CAPRYLIC/CAPRIC TRIGLYCERIDE -
 DIMETHICONE/VINYL DIMETHICONE
 CROSSPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8 -
 MAGNESIUM MYRISTATE - NYLON-12 -
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 CROSSPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYQUATERNIUM-51 -
 THYMUS SERPILLUM EXTRACT -
 ALUMINUM HYDROXIDE -
 ALUMINUM DISTEARATE -
 POLYSILOXANE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 CROSSPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 CROSSPOLYMER -
 SILICA - ALUMINA -
 POLYMETHYLSILSESQUOXANE -
 DIMOSTEARYL MALATE -
 GLYCERYL DIBESTEARATE -
 DIPROPYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 NICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
CORPORATION, DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

READ INSERTION



1-800-40-5041

AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Rose 4

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Rose 4

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun. ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor. ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRIISOPROPYLDIMETHYL
 TRIMETHYLSILOXYLICATE -
 CAPRYLIC/CAPRIC TRIGLYCERIDE -
 DIMETHICONE/VINYL DIMETHICONE
 CROSSPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8 -
 MAGNESIUM MYRISTATE - NYLON-12 -
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 CROSSPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYQUATERNIUM-51 -
 THYMUS SERPILLUM EXTRACT -
 ALUMINUM HYDROXIDE -
 ALUMINUM DISTEARATE -
 POLYSILOXANE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 CROSSPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 CROSSPOLYMER -
 SILICA - ALUMINA -
 POLYMETHYLSILSESQUOXANE -
 DISOSTEARYL MALATE -
 GLYCERYL DISOSTEARATE -
 DIPROPYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
CORPORATION, DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

READ INSERTION.



40-5041
AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Golden 2

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Golden 2

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRIMETHYLOLPROPYLENEDIMETHYL
 TRIMETHYLSILOXYMETHYL
 CAPRYLIC/CAPRIC TRIGLYCERIDE -
 DIMETHICONE/VINYL DIMETHICONE
 CROSSPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8 -
 MAGNESIUM MYRISTATE - NYLON-12
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 CROSSPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYQUATERNIUM-51 -
 THYMUS SERPILLUM EXTRACT -
 ALUMINUM HYDROXIDE -
 ALUMINUM DISTEARATE -
 POLYSILICONE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 CROSSPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 CROSSPOLYMER -
 SILICA - ALUMINA -
 POLYMETHYLSILSESQUOXANE -
 DIMOSTEARYL MALATE -
 GLYCERYL DIBESTEARATE -
 DIPROPYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 NIACA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
CORPORATION, DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

READ INSERTION



1-800-40-5041
AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Golden 3

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Golden 3

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active Ingredients Purpose
OCINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - TRIFLUDROPROPYLDIMETHYL TRIMETHYLSILOXYLICATE - CAPRYLIC/CAPRIC TRIGLYCERIDE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - PEG-8 POLYDIMETHYLSILOXYETHYL DIMETHICONE - SD ALCOHOL 40-8 - MAGNESIUM MYRISTATE - NYLON-12 - DIPHENYL DIMETHICONE/VINYL DIPHENYL DIMETHICONE/SILSESQUOXANE CROSSPOLYMER - DISTEARDIMONIUM HECTORITE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - ALUMINUM HYDROXIDE - ALUMINUM DISTEARATE - POLYSILICONE-2 - GLYCERIN - BHT - ALCOHOL - BUTYLENE GLYCOL - TOCOPHEROL - METHYL METHACRYLATE CROSSPOLYMER - HYPERICUM ERECTUM FLOWER/LEAF/STEM EXTRACT - TALC - DIMETHICONE/PEG-10/15 CROSSPOLYMER - SILICA - ALUMINA - POLYMETHYLSILSESQUOXANE - DIMYRISTEARYL MALATE - GLYCERYL DIBSOSTEARATE - DIPROPYLENE GLYCOL - PHENOXYETHANOL - FRAGRANCE - TITANIUM DIOXIDE - IRON OXIDES - MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
Lasting
Liquid Foundation
Oil Free
BROAD SPECTRUM
SPF 20
Sunscreen

30mL 1 FL. OZ.

READ INSERTION



SHISEIDO AMERICAS CORPORATION, DIST
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

40-5041
AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Golden 4

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Golden 4

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCTINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - TRIBUTYLDIPROPYLDIMETHYLTRIMETHYLSILOXYLICATE - CAPRYLIC/CAPRIC TRIGLYCERIDE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE - SD ALCOHOL 40-8 - MAGNESIUM MYRISTATE - NYLON-12 - DIPHENYL DIMETHICONE/VINYL DIPHENYL DIMETHICONE/SILSESQUOXANE CROSSPOLYMER - DISTEARDIMONIUM HECTORITE - POLYOLATERIUM-51 - THYMUS SERPYLLUM EXTRACT - ALUMINUM HYDROXIDE - ALUMINUM DISTEARATE - POLYSILOXANE-2 - GLYCERIN - BHT - ALCOHOL - BUTYLENE GLYCOL - TOCOPHEROL - METHYL METHACRYLATE CROSSPOLYMER - HYPERICUM ERECTUM FLOWER/LEAF/STEM EXTRACT - TALC - DIMETHICONE/PEG-10/15 CROSSPOLYMER - SILICA - ALUMINA - POLYMETHYLSILSESQUOXANE - DIBENZOYL MALATE - GLYCERYL DIBESTEARATE - DIPROPYLENE GLYCOL - PHENOXYETHANOL - FRAGRANCE - TITANIUM DIOXIDE - IRON OXIDES - MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
Lasting
Liquid Foundation
Oil Free
BROAD SPECTRUM
SPF 20
Sunscreen

30mL 1 FL. OZ.

READ INSERTION.



SHISEIDO AMERICAS CORPORATION, DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
www.shiseido.com

40-5041
AAA

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Golden 5

**SHISEIDO
GINZA TOKYO**

SYNCHRO SKIN

Lasting
Liquid Foundation

Oil Free

BROAD SPECTRUM
SPF 20

Sunscreen

30mL 1 FL. OZ.

Lasting Liquid Foundation



Golden 5

ULTIMATE SKIN AFFINITY. A NEW RESPONSIVE SENSORY TECHNOLOGY.

An intelligent, long-lasting foundation that beautifully enhances your complexion with vibrance. Flawlessly undetectable. Like it belongs to your skin.

Drug Facts

Active ingredients Purpose
OCINOXATE 2.9% Sunscreen
TITANIUM DIOXIDE 2.4% Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun ▶

Drug Facts (continued)

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor ▶

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER
 TRILUOROPROPYLDIMETHYL
 TRIMETHYLSILOXYLICATE -
 CAPRYLIC/CAPRIC TRIGLYCERIDE -
 DIMETHICONE/VINYL DIMETHICONE
 COPOLYMER -
 PEG-9 POLYDIMETHYLSILOXYETHYL
 DIMETHICONE -
 SD ALCOHOL 40-8 -
 MAGNESIUM MYRISTATE - NYLON-12 -
 DIPHENYL DIMETHICONE/VINYL
 DIPHENYL
 DIMETHICONE/SILSESQUOXANE
 COPOLYMER -
 DISTEARDIMONIUM HECTORITE -
 POLYOLATERIUM-51 -
 THYMUS SERPILLUM EXTRACT -
 ALUMINIUM HYDROXIDE -
 ALUMINIUM DISTEARATE -
 POLYSILOXANE-2 - GLYCERIN - BHT -
 ALCOHOL - BUTYLENE GLYCOL -
 TOCOPHEROL -
 METHYL METHACRYLATE
 COPOLYMER -
 HYPERICUM ERECTUM
 FLOWER/LEAF/STEM EXTRACT -
 TALC -
 DIMETHICONE/PEG-10/15
 COPOLYMER -
 SILICA - ALUMINA -
 POLYMETHYLSILSESQUOXANE -
 DIOSTEARYL MALATE -
 GLYCERYL DIISOSTEARATE -
 DIPROPYLENE GLYCOL -
 PHENOXYETHANOL - FRAGRANCE -
 TITANIUM DIOXIDE - IRON OXIDES -
 MICA - RED 27 -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
 Lasting
 Liquid Foundation
 Oil Free
 BROAD SPECTRUM
 SPF 20
 Sunscreen

30mL 1 FL. OZ.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS
 CORPORATION, DIST.
 NY, NY 10022
 MADE IN U.S.A.
 www.shiseido.com

READ INSERTION



40-5041
 AAA

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 1

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-261
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0K0OR)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)	
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-261-10	1 in 1 CARTON	02/01/2016	

1	30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 2

octinoxate and titanium dioxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-262
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL	
Inactive Ingredients			
Ingredient Name	Strength		
DIMETHICONE (UNII: 92RU3N3Y1O)			
WATER (UNII: 059QF0K00R)			
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)			
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)			
MAGNESIUM MYRISTATE (UNII: Z1917F0578)			
NYLON-12 (UNII: 446U8J075B)			
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)			
THYMUS SERPYLLUM (UNII: 86H4S6K51N)			
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)			
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)			
GLYCERIN (UNII: PDC6A3C00X)			
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)			
ALCOHOL (UNII: 3K9958V90M)			
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)			
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)			
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)			
TALC (UNII: 7SEV7J4R1U)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
ALUMINUM OXIDE (UNII: LMI26O6933)			
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)			
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)			
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)			
DIPROPYLENE GLYCOL (UNII: E107L85C40)			
PHENOXYETHANOL (UNII: HIE492ZZ3T)			
FERRIC OXIDE RED (UNII: 1K09F3G675)			
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)			
FERROSFERRIC OXIDE (UNII: XM0M87F357)			
MICA (UNII: V8A1AW0880)			

D&C RED NO. 27 (UNII: 2LRS185U6K)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-262-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 3

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-263
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0K00R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	

DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)
D&C RED NO. 27 (UNII: 2LRS185U6K)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-263-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION NEUTRAL 4

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-264
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
GLYCERIN (UNII: PDC6A3C00X)	

BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALCOHOL (UNII: 3K9958V90M)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)
TALC (UNII: 7SEV7J4R1U)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)
D&C RED NO. 27 (UNII: 2LRS185U6K)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-264-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION ROSE 2

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-265
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)
MAGNESIUM MYRISTATE (UNII: Z1917F0578)
NYLON-12 (UNII: 446U8J075B)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
THYMUS SERPYLLUM (UNII: 86H4S6K51N)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)
GLYCERIN (UNII: PDC6A3C00X)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALCOHOL (UNII: 3K9958V90M)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)
TALC (UNII: 7SEV7J4R1U)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)
D&C RED NO. 27 (UNII: 2LRS185U6K)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-265-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION ROSE 3

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-266
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL

TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) | TITANIUM DIOXIDE | 754 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0K00R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)	
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-266-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION ROSE 4

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-267	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL	
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	DIMETHICONE (UNII: 92RU3N3Y1O)			
	WATER (UNII: 059QF0K00R)			
	MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)			
	PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)			
	MAGNESIUM MYRISTATE (UNII: Z1917F0578)			
	NYLON-12 (UNII: 446U8J075B)			
	DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)			
	THYMUS SERPYLLUM (UNII: 86H4S6K51N)			
	ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)			
	ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)			
	GLYCERIN (UNII: PDC6A3C00X)			
	BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)			
	ALCOHOL (UNII: 3K9958V90M)			
	BUTYLENE GLYCOL (UNII: 3XUS85K0RA)			
	.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)			
	HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)			
	TALC (UNII: 7SEV7J4R1U)			
	SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
	ALUMINUM OXIDE (UNII: LMI26O6933)			
	POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)			
	DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)			
	GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)			
	DIPROPYLENE GLYCOL (UNII: E107L85C40)			
	PHENOXYETHANOL (UNII: HIE492ZZ3T)			
	FERRIC OXIDE RED (UNII: 1K09F3G675)			
	FERRIC OXIDE YELLOW (UNII: EX438O2MRT)			
	FERROSFERRIC OXIDE (UNII: XM0M87F357)			
	MICA (UNII: V8A1AW0880)			
	D&C RED NO. 27 (UNII: 2LRS185U6K)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-267-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing	Application Number or Monograph	Marketing Start	Marketing End	

Category	Citation	Date	Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 2

octinoxate and titanium dioxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-268
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)	
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	

Packaging				
#	Item Code	Package Description	Marketing Start	Marketing End

#	Item Code	Package Description	Date	Date
1	NDC:58411-268-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 3

octinoxate and titanium dioxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-269
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg	in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg	in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0K00R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALCOHOL (UNII: 3K9958V90M)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)	
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)				
FERROSO FERRIC OXIDE (UNII: XM0M87F357)				
MICA (UNII: V8A1AW0880)				
D&C RED NO. 27 (UNII: 2LRS185U6K)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-269-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph drug	M020	02/01/2016		

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 4			
octinoxate and titanium dioxide liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-270
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL	
Inactive Ingredients			
Ingredient Name	Strength		
DIMETHICONE (UNII: 92RU3N3Y1O)			
WATER (UNII: 059QF0K0OR)			
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)			
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)			
MAGNESIUM MYRISTATE (UNII: Z1917F0578)			
NYLON-12 (UNII: 446U8J075B)			
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)			
THYMUS SERPYLLUM (UNII: 86H4S6K51N)			
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)			
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)			
GLYCERIN (UNII: PDC6A3C00X)			
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)			
ALCOHOL (UNII: 3K9958V90M)			
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)			
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)			
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)			
TALC (UNII: 7SEV7J4R1U)			

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)
D&C RED NO. 27 (UNII: 2LRS185U6K)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-270-10	1 in 1 CARTON	02/01/2016	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	

SHISEIDO SYNCHRO SKIN LASTING FOUNDATION GOLDEN 5

octinoxate and titanium dioxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-271
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	911 mg in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	754 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)	
MAGNESIUM MYRISTATE (UNII: Z1917F0578)	
NYLON-12 (UNII: 446U8J075B)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	

ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)
GLYCERIN (UNII: PDC6A3C0OX)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALCOHOL (UNII: 3K9958V90M)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
HYPERICUM ERECTUM FLOWERING TOP (UNII: IT0296HQDC)
TALC (UNII: 7SEV7J4R1U)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
DIISOSTEARYL MALATE (UNII: QBS8A3XZGQ)
GLYCERYL DIISOSTEARATE (UNII: 68BAV42LRC)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)
D&C RED NO. 27 (UNII: 2LRS185U6K)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-271-10	1 in 1 CARTON	02/01/2016	01/08/2021
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M020	02/01/2016	01/08/2021

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Establishment

Name	Address	ID/FEI	Business Operations
SHISEIDO AMERICA INC.		782677132	manufacture(58411-261, 58411-262, 58411-263, 58411-264, 58411-265, 58411-266, 58411-267, 58411-268, 58411-269, 58411-270, 58411-271) , analysis(58411-261, 58411-262, 58411-263, 58411-264, 58411-265, 58411-266, 58411-267, 58411-268, 58411-269, 58411-270, 58411-271)

Revised: 10/2023

SHISEIDO AMERICAS CORPORATION