PATTERSON DENTAL TOPICAL ANESTHETIC- benzocaine gel Patterson Dental

Benzocaine 200mg (in each g)

Oral Anesthetic

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of o product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use


Keep out of reach of children.

Store at 59°-86°F (15°-30°C). Protect from freezing.


flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

800.873.7683









PATTERSON DENTAL TOPICAL ANESTHETIC

Other information Store at 59° - 86°F (15 -30°C).

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color

Questions or comments:

300.873.7683

benzocaine gel

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:50227-1013 DENTAL **Route of Administration**

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients			
Ingredient Name	Strength		
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			

Colorwhite (DYE-FREE)ScoreShapeSizeFlavorMINTImprint Code	Product Characteristics		
	Color	white (DYE-FREE)	Score
Flavor MINT Imprint Code	Shape		Size
	Flavor	MINT	Imprint Code
Contains	Contains		

Pa	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:50227- 1013-3	30 g in 1 JAR; Type 0: Not a Combination Product	10/01/2025	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M022	10/01/2025		

Labeler - Patterson Dental (171843584)

Revised: 9/2025 Patterson Dental