# CHEEKY BONSAI UTI PAIN RELIEF- phenazopyridine hydrochloride tablet IXXA, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

#### **DRUG FACTS**

#### Active ingredient (in each tablet)

Phenazopyridine Hydrochloride 99.5 mg.

#### **Purpose**

**Urinary Analgesic** 

#### Warnings

Do not exceed recommended dosage

Do not use if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician

#### Ask doctor before use if you have

- kidney disease
- allergies to food, preservatives or dyes
- had a hypersensitive reaction to phenazopyridine

#### When using this product

- stomach upset may occur, taking this product with or after meals may reduce stomach upset
- your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

#### Stop use and ask doctor if

- your symptoms last for more than 2 days
- you suspect you are having an adverse reaction to the medication

Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.

#### If pregnant or breast feeding,

Ask a health professional before use.

#### Keep out of reach of children

In case of an overdose, get medical help or contact a Poison Control Center right away.

#### Use

Fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections.

#### **Inactive ingredients**

Corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, talc and triacetin.

#### **Directions**

- adults and children 12 years and over: take 2 tablets 3 times daily with a full glass of water, with or after meals as needed
- children under 12 years: consult a doctor
- Do not use for more than 2 days (12 tablets) without consulting a doctor



#### **CHEEKY BONSAI UTI PAIN RELIEF**

phenazopyridine hydrochloride tablet

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:81860-113

Route of Administration ORAL

#### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
	PHENAZ OPYRIDINE HYDROCHLORIDE	99.5 mg

Inactive Ingredients		
Ingredient Name	Strength	
LACTOSE (UNII: J2B2A4N98G)		
STARCH, CORN (UNII: O8232NY3SJ)		
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWIA)	
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TALC (UNII: 7SEV7J4R1U)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics			
Color	brown	Score	no score
Shape	OVAL	Size	9mm
Flavor		Imprint Code	p99
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:81860- 113-24	1 in 1 CARTON	05/12/2021		
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/12/2021	

### **Labeler -** IXXA, INC (117979157)

## **Registrant -** Reese Pharmaceutical Co (004172052)

Revised: 12/2022 IXXA, INC