

**FEVERALL JR. STRENGTH- acetaminophen suppository**  
**Bryant Ranch Prepack**

-----  
**FeverAll®**  
**Jr. Strength**

***Drug Facts***

**Active ingredient (in each rectal suppository)**

Acetaminophen, USP 325 mg

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily

- reduces fever
- relieves minor aches, pains, and headache

**Warnings**

**Liver warning**

This product contains acetaminophen.

Severe liver damage may occur if

- a child 6 to 12 years takes more than 5 doses in 24 hours
- an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- an adult takes 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**For rectal use only.**

**Do not use**

- in children under 6 years
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease.
- you are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition.

**If pregnant or breast-feeding**,ask a health professional before use. **Keep out of reach of children.**If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Directions**

- **do not use more than directed**
- remove wrapper
- carefully insert suppository well up into the rectum

**Dosing Chart**

<b>Age</b>	<b>Dose</b>
under 6 years	do not use
6 to 12 years	Use 1 suppository every 4 to 6 hours. (maximum of 5 doses in 24 hours)
adults and children 12 years and older	Use 2 suppositories every 4 to 6 hours. (maximum of 6 doses)

**Other information**

- store at 2°-27°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients**

glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

## Questions?

call 1-866-923-4914

Dist. by: **Taro Pharmaceuticals U.S.A., Inc.**  
Hawthorne, NY 10532


## HOW SUPPLIED

Acetaminophen Suppositories 325 mg

- NDC 63629-8240-1: 12 Rectal Suppositories

Repackaged/Relabeled by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504

Acetaminophen Suppositories 325 mg

 GTIN 00363629824014 Lot 208820 Exp 2/18/2027 SN 0123456789	<b>Drug Facts</b>	
	<b>Active ingredient (in each rectal suppository)</b> Acetaminophen, USP 325 mg	<b>Purpose</b> Pain reliever/fever reducer
	<b>Uses</b> -temporarily reduces fever, relieves minor aches, pains, and headache.	
	<b>Warnings</b> Liver warning: This product contains acetaminophen. Severe liver damage may occur. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. For rectal use only. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.	
	<b>Other Information</b> -do not use if imprinted suppository wrapper is opened or damaged. -store at 2°-27°C (35°-80°F). -Scan Package Insert QR code for more information.	
	<b>Directions</b> -do not use more than directed, remove wrapper, carefully insert suppository well up into the rectum. Under 6 years; do not use. 6 to 12 years; Use 1 suppository every 4 to 6 hours. (maximum of 5 doses in 24 hours). Adults and children 12 years and older; Use 2 suppositories every 4 to 6 hours. (maximum of 6 doses).	
<b>Inactive Ingredients</b> glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80.		

**NDC 63629-8240-1**

---

**Acetaminophen  
Suppositories**

**325 mg**

**12 Rectal Suppositories**

**BRP**  
PHARMACEUTICALS

Repackaged by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504 USA

Manufactured by:  
G&W NC Laboratories, Inc.



Package  
Insert

## FEVERALL JR. STRENGTH

acetaminophen suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63629-8240(NDC:51672-2116)
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XE4)	
<b>HYDROGENATED PALM KERNEL OIL</b> (UNII: FM8D1RE2VP)	
<b>PEG-100 STEARATE</b> (UNII: YD01N1999R)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629-8240-1	12 in 1 POUCH	09/10/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018337	12/12/2013	

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

## Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(63629-8240) , RELABEL(63629-8240)

Revised: 1/2025

Bryant Ranch Prepack