

GILTUSS DIABETIC- dextromethorphan hydrobromide, guaifenesin solution
Gil Pharmaceutical Corp

Giltuss Diabetic Cough & Cold

Active Ingredients

Active ingredients

Dextromethorphan HBr, 20 mg

Guaifenesin , 200 mg

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg

Guaifenesin, 200 mg

Purposes

Cough Suppressant

Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Warnings

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or

Purpose & Uses Section

Purposes

Cough suppressant

Expectorant

Uses

temporarily relieves, cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants, the intensity of coughing, the impulse to cough to help you get to sleep, helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg

Guaifenesin, 200 mg

Purposes

Cough Suppressant

Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse

to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Warnings

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or

Warnings

Warnings

Do not use

If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you don't know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg

Guaifenesin, 200 mg

Purposes

Cough Suppressant

Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Warnings

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or

Ask a doctor

Ask a doctor before use if you have, a cough with too much phlegm(mucus) , a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema, when using this product do not use more than directed.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg

Guaifenesin, 200 mg

Purposes

Cough Suppressant

Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Warnings

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or

emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or

Stop use and ask a doctor if

Stop use and ask a doctor if

cough lasts for more than 7 days , come back, or is accompanied by fever, rash or persistent headache. These could be signs of serious condition.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg

Guaifenesin, 200 mg

Purposes

Cough Suppressant

Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Warnings

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or

Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or

If pregnant or breast- feeding

If pregnant or breast- feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poisson Control center immediatly

persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions

- do not exceed recommended dosage
- do not take more than 6 doses in any 24 hour period
- keep dosing cup with product
- mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call **1-787-848-9114**, Monday–Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 06/22



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Directions

persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions

- do not exceed recommended dosage
- do not take more than 6 doses in any 24 hour period
- keep dosing cup with product
- mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call **1-787-848-9114**, Monday–Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 06/22



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Directions

Do not exceed recommended dosage

Do not take more than 6 doses in any 24 hours period

Keep dosing cup with product

ml= milliliters

Adults and children 12 years & over 10 ml every 4 hours

children 6 to under 12 years of age 5 ml every 4 hours

children 4 to under 6 years of age 2.5 ml every 4 hours

children under 4 years of age Consult a doctor

Other information

Other information

- Store at room temperature 15-30 °C (59° -86 ° F)
- Close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions

- do not exceed recommended dosage
- do not take more than 6 doses in any 24 hour period
- keep dosing cup with product
- mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call **1-787-848-9114**, Monday–Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 06/22



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Inactive Ingredients

1

persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions

- do not exceed recommended dosage
- do not take more than 6 doses in any 24 hour period
- keep dosing cup with product
- mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call **1-787-848-9114**, Monday–Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 06/22



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Questions or comments?

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST, Call your doctor for medical advice in the event of side effects.

persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Directions

- do not exceed recommended dosage
- do not take more than 6 doses in any 24 hour period
- keep dosing cup with product
- mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call **1-787-848-9114**, Monday–Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 06/22

Indication and usage section

Indication and usage section

Do not exceed recommended dosage

Do not take more than 6 doses in any 24 hours period

Adults and children 12 years and over 10 ml every 4 hours


Children 6 to under 12 years age 5 ml every 4 hours

Children 4 to under 6 years age 2.5 ml every 4 hours

Children under 4 years of age Consult a doctor



Keep out of reach of children.



Drug Facts

Active ingredients (in each 10 mL)
 Dextromethorphan HBr, 20 mg
 Guaifenesin, 200 mg

Purposes
 Cough Suppressant
 Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes.

Warnings

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or

SUGAR FREE, ALCOHOL FREE
 SORBITOL FREE, SODIUM FREE
 SIN AZÚCAR, SIN ALCOHOL,
 SIN SORBITOL, SIN SODIO



Giltuss
DIABETIC
COUGH & COLD
TOS Y CATARRO

DYE FREE, FRUCTOSE FREE, GLUTEN FREE, NON-DROWSY,
 SIN COLORANTES, SIN FRUCTOSA, SIN GLUTEN, NO CAUSA SUEÑO.

COUGH AND COLD, CHEST CONGESTION,
 LIQUIDS AND LOOSENS PHLEGM AND MUCUS
 TOS Y CATARRO, CONGESTIÓN DE PECHO,
 LÍQUIDA Y SUelta LA FLEMA Y EL MOCO



RASPBERRY FLAVOR,
 SAZÓN A FRAMBUESA

4 FL OZ (118 mL)

persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions

- do not exceed recommended dosage
 - do not take more than 6 doses in any 24 hour period
 - keep dosing cup with product
 - mL = milliliters
- | | |
|-------------------------------------|----------------------|
| adults and children 12 years & over | 10 mL every 4 hours |
| children 6 to under 12 years of age | 5 mL every 4 hours |
| children 4 to under 6 years of age | 2.5 mL every 4 hours |
| children under 4 years of age | Consult a doctor |

Other Information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 06/22



MANUFACTURED FOR
 GIL PHARMACEUTICAL CORP.
 PONCE, PUERTO RICO 00717-1565

giltuss.com

Keep out of reach of children.

In case of accidental overdose, get medical or professional help or contact a Poisson Control center immediatly

**SUGAR FREE. ALCOHOL FREE.
SORBITOL FREE. SODIUM FREE.**

**SIN AZÚCAR. SIN ALCOHOL.
SIN SORBITOL. SIN SODIO.**



Giltuss
DIABETIC
COUGH & COLD
TOS Y CATARRO

DYE FREE. FRUCTOSE FREE. GLUTEN FREE. NON-DROWSY.
SIN COLORANTES. SIN FRUCTOSA. SIN GLUTEN. NO CAUSA SUEÑO.

**COUGH AND COLD, CHEST CONGESTION,
LIQUIFIES AND LOOSENS PHLEGM AND MUCUS**
**TOS Y CATARRO, CONGESTIÓN DE PECHO,
LICUA Y SUELTA LA FLEMA Y EL MOCO**



**SAFE FOR PEOPLE
WITH DIABETES**

SEGURO PARA PERSONAS CON DIABETES

4FL OZ (118 mL)

RASPBERRY FLAVOR.
SABOR A FRAMBUESA.



Giltuss Diabetic

Cough & Cold

Sugar Free, Alcohol free, Sorbitol free, Sodium free

Dye free, Fructose free, Gluten free, Non -Drowsy

Cough & Cold, Chest congestion, liquifies and loosens Phlegm and mucus

Safe for people with diabetes

4 floz 118 ml

Raspberry flavor

GILTUSS DIABETIC

dextromethorphan hydrobromide, guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-009
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
DEXTROMETHORPHAN HYDROCHLORIDE (UNII: 351KFH969D) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROCHLORIDE	20 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
CITRIC ACID (UNII: 2968PHW8QP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	RASPBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:65852-009-04	1 in 1 CARTON	10/17/2018	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/17/2018	

Labeler - Gil Pharmaceutical Corp (007392322)

Revised: 9/2025

Gil Pharmaceutical Corp