

NUMQUICK TOPICAL ANALGESIC- epinephrine hydrochloride, lidocaine hydrochloride gel
Unit Dose, Ltd.

Numquick Topical Analgesic

Drug Facts

Active ingredient

(-) Epinephrine HCL w/w 0.01%

Lidocaine HCl 5% w/w

Purpose

Vasoconstrictor

Local Anesthetic

Uses:

Temporarily relieves local discomfort or pain or burning associated with anorectal disorders.

Warnings:

External use only

certain persons can develop allergic reactions to the ingredients in this product. **Allergy alert:**

Do not use:

this product in the rectum by using fingers or any mechanical device or applicator.

When using this product

do not use more than directed.

Stop use and ask a doctor

if pain worsens or does not improve in 7 days

- if redness, irritation, swelling, pain or other symptoms develop or increase
- if bleeding occurs

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- adults: clean the affected area with mild soap and warm water and rinse thoroughly
- dry gently
- apply 1-2 g. externally to the affected area up to 4 times daily
- children under 12 years: ask a doctor.

Other information:

Store in a cool dark place

Inactive ingredients:

Aqua (DI water), chlorobutanol, methylcellulose, sodium EDTA, sodium metabisulfite

Questions?

Call Toll Free 888-452-4946

Package Labeling:

Drug Facts

Numquick 5% Lidocaine Gel

Active ingredient (in each gm)	Purpose
(-) Epinephrine HCL w/w 0.01%	Vasoconstrictor
Lidocaine HCL 5% w/w	Local Anesthetic

Uses: Temporarily relieves local discomfort or pain or burning associated with anorectal disorders.

Warnings: External use only

Allergy alert: certain persons can develop allergic reactions to the ingredients in this product.

Do not use: this product in the rectum by using fingers or any mechanical device or applicator.

When using this product do not use more than directed.

Stop use and ask a doctor if pain worsens or does not improve in 7 days • if redness, irritation, swelling, pain or other symptoms develop or increase • if bleeding occurs

Keep out of reach of children. *If swallowed, get medical help or contact a Poison Control Center right away.*

Directions: •adults: clean the affected area with mild soap and warm water and rinse thoroughly • dry gently
• apply 1-2 g. externally to the affected area up to 4 times daily
• children under 12 years: ask a doctor.

Other information: Store in a cool dark place

Inactive ingredients: Aqua (DI water), chlorobutanol, methylcellulose, sodium EDTA, sodium metabisulfite

Questions? Call Toll Free 888-452-4946

NUMQUICK TOPICAL ANALGESIC

epinephrine hydrochloride, lidocaine hydrochloride gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67194-005
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EPINEPHRINE HYDROCHLORIDE (UNII: WBB0470038) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE	0.1 mg in 1 mL
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
CHLOROBUTANOL (UNII: HM4YQM8WRC)	
EDETATE SODIUM (UNII: MP1J8420LU)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67194-005-01	59 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/20/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M015	01/20/2016	

Labeler - Unit Dose, Ltd. (119080393)

Revised: 10/2023

Unit Dose, Ltd.