

GILTUSS HBP COUGH AND CHEST CONGESTION- dextromethorphan hydrobromide, guaifenesin solution
Dextrum Laboratories Inc.

Giltuss HBP Cough and Chest Congestion

Active ingredients

Active Ingredients

Dextromethorphan HBr, 20 mg

Guaifenesin, 200 mg

Drug Facts

Active ingredients (in each 10 mL)

Purposes

Dextromethorphan HBr, 20 mg	Cough Suppressant
Guaifenesin, 200 mg	Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Purposes and Uses

Purposes

Cough supresant

Expectorant

Uses

Temporary relieves, cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants , the intensity of coughing, the impulse to cough to help you get to sleep, help loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Drug Facts

Active ingredients (in each 10 mL)

Purposes

Dextromethorphan HBr, 20 mg	Cough Suppresant
Guaifenesin, 200 mg	Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to

increased irritability • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Warnings

Warnings

Do not use

If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you don't know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts

Active ingredients (in each 10 mL)

Purposes

Dextromethorphan HBr, 20 mg	Cough Suppressant
Guaifenesin, 200 mg	Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Ask a doctor

Ask a doctor before use if you have, a cough with too much phlegm(mucus) , a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema, when using this product do not use more than directed.

Drug Facts

Active ingredients (in each 10 mL)

Purposes

Dextromethorphan HBr, 20 mg	Cough Suppressant
Guaifenesin, 200 mg	Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or

emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Stop use and ask a doctor

Stop use and ask a doctor if

cough lasts for more than 7 days , come back, or is accompanied by fever, rash or persistent headache. These could be signs of serious condition.

Drug Facts

Active ingredients (in each 10 mL)

Purposes

Dextromethorphan HBr, 20 mg	Cough Suppressant
Guaifenesin, 200 mg	Expectorant

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease)

psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control center immediately

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions • do not exceed recommended dosage • do not take more than 6 doses in any 24 hour period • keep dosing cup with product • mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- Tamper evident: Do not use if safety seal is broken or missing
- Store at controlled room temperature, between 15°-30°C (59° -86°F)

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Directions

Directions

Do not exceed recommended dosage

Do not take more than 6 doses in any 24 hours period

Keep dosing cup with product

ml= milliliters

Adults and children 12 years & over 10 ml every 4 hours

children 6 to under 12 years of age 5 ml every 4 hours

children 4 to under 6 years of age 2.5 ml every 4 hours

children under 4 years of age Consult a doctor

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions • do not exceed recommended dosage • do not take more than 6 doses in any 24 hour period • keep dosing cup with product • mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- Tamper evident: Do not use if safety seal is broken or missing
- Store at controlled room temperature, between 15°-30°C (59° -86°F)

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Other information

Other information

Tamper evident : Do not use if safety seal is broken or missing

Store at controlled

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions • do not exceed recommended dosage • do not take more than 6 doses in any 24 hour period • keep dosing cup with product • mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- Tamper evident: Do not use if safety seal is broken or missing
- Store at controlled room temperature, between 15°-30°C (59° -86°F)

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday–Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

room temperature 15-30 °C (59° -86 ° F)

Inactive Ingredients

Citric Acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions • do not exceed recommended dosage • do not take more than 6 doses in any 24 hour period • keep dosing cup with product • mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- Tamper evident: Do not use if safety seal is broken or missing
- Store at controlled room temperature, between 15°-30°C (59° -86°F)

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday–Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Questions or comments

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM – 5PM EST

Call your doctor for medical advice in the event of side effects.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions • do not exceed recommended dosage • do not take more than 6 doses in any 24 hour period • keep dosing cup with product • mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- Tamper evident: Do not use if safety seal is broken or missing
- Store at controlled room temperature, between 15°-30°C (59° -86°F)

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

keep out of seach children



keep out of reach of the children.

In case of accidental overdose get medical professional help or contact a Poison Control Center immediately

Indication and usage

Do not exceed recommended dosage

Do not take more than 6 doses in any 24 hours period

Keep dosing cup with product

ml= milliliters

Adults and children 12 years & over 10 ml every 4 hours

children 6 to under 12 years of age 5 ml every 4 hours

children 4 to under 6 years of age 2.5 ml every 4 hours

children under 4 years of age Consult a doctor

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, 20 mg Cough Suppressant
Guaifenesin, 200 mg Expectorant

Purposes

Uses • temporarily relieves: • cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive and drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • when using this product do not use more than directed

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

MAXIMUM STRENGTH • MÁXIMA CONCENTRACIÓN



HIGH BLOOD PRESSURE HBP

Giltuss
COUGH & CHEST
CONGESTION

TOS Y CONGESTIÓN DE PECHO

ALCOHOL FREE. SUGAR FREE. DYE FREE. NON-DROWSY.
SIN ALCOHOL. SIN AZÚCAR. SIN COLORANTES. NO CAUSA SUEÑO.

COUGH AND COLD, CHEST CONGESTION,
LIQUIFIES AND LOOSENS PHEGM AND MUCUS
TOS Y CATARRO, CONGESTIÓN DE PECHO,
LICUA Y SUELTA LA FLEMA Y EL MOCO



RASPBERRY FLAVOR
SABOR A FRAMBUESA

4 FL OZ (118 mL)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical or professional help or contact a Poison Control Center immediately.

Directions • do not exceed recommended dosage • do not take more than 6 doses in any 24 hour period • keep dosing cup with product • mL=milliliters

adults and children 12 years & over	10 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
children 4 to under 6 years of age	2.5 mL every 4 hours
children under 4 years of age	Consult a doctor

Other information

- Tamper evident: Do not use if safety seal is broken or missing
- Store at controlled room temperature, between 15°-30°C (59°-86°F)

Inactive ingredients: Citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water and sucralose.

Questions or Comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1568

giltuss.com

Principal panel

MAXIMUM STRENGTH. MÁXIMA CONCENTRACIÓN.



HIGH BLOOD PRESSURE HBP

Giltuss.

**COUGH & CHEST
CONGESTION**

TOS Y CONGESTIÓN DE PECHO

**ALCOHOL FREE. SUGAR FREE. DYE FREE. NON-DROWSY.
SIN ALCOHOL. SIN AZÚCAR. SIN COLORANTES. NO CAUSA SUEÑO.**

**COUGH AND COLD, CHEST CONGESTION,
LIQUIFIES AND LOOSENS PHLEGM AND MUCUS
TOS Y CATARRO, CONGESTIÓN DE PECHO,
LICUA Y SUELTA LA FLEMA Y EL MOCO**



**RASPBERRY FLAVOR.
SABOR A FRAMBUESA.**

**SEGURO
SAFE**



4FL OZ (118 mL)

Maximum strength

High Blood Pressure

Giltuss HBP

Cough & Chest Congestion

Alcohol free, Sugar Free, Dye free, Non -Drowsy

Cough and cold, chest congestion, liquifies and loosens phlegm and mucus

Raspberry flavor

4 floz 118 ml

GILTUSS HBP COUGH AND CHEST CONGESTION

dextromethorphan hydrobromide, guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-006
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID (UNII: 2968PHW8QP)	
RASPBERRY (UNII: 4N14V5R27W)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	RASPBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65852-006-04	1 in 1 CARTON	11/13/2018	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	11/13/2018	

Labeler - Dextrum Laboratories Inc. (007392322)

Revised: 9/2025

Dextrum Laboratories Inc.