

GILTUSS HONEY DM COUGH- dextromethorphan hbr solution
Dextrum Laboratories, Inc

Giltuss Honey DM Cough

Active Ingredients

Active Ingredients (in each 10 ml)

Dextromethorphan, 30 mg

Drug Facts

<i>Active ingredient (in each 10 mL)</i>	<i>Purpose</i>
Dextromethorphan HBr, 30 mg.....	Cough Suppressant

Uses

• temporarily relieves: • cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Purpose and uses

Purpose

Cough supresant

Uses

Temporary relieves:

Cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants, the intensity of coughing ,the impulse to cough to help you get sleep,

Drug Facts

Active ingredient (in each 10 mL)	Purpose
Dextromethorphan HBr, 30 mg.....	Cough Suppresant

Uses

• temporarily relieves: • cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease)

psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Warning

Warnings

Do not use

If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you don't know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts

Active ingredient (in each 10 mL)	Purpose
Dextromethorphan HBr, 30 mg	Cough Suppressant

Uses

• temporarily relieves a cough due to minor throat and

• temporarily relieves. • cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Ask a Doctor

Ask a doctor before use if you have, a cough with too much phlegm (mucus) , a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema.

Drug Facts

Active ingredient (in each 10 mL)**Purpose**

Dextromethorphan HBr, 30 mg Cough Suppressant

Uses

• temporarily relieves: • cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Stop use and ask a doctor if

cough lasts for more than 7 days , come back, or is accompanied by fever, rash or persistent headache. These could be signs of serious condition.

Drug Facts

Active ingredient (in each 10 mL)

Purpose

Dextromethorphan HBr, 30 mg	Cough Suppressant
-----------------------------------	-------------------

Uses

- temporarily relieves:
- cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding ask a health professional

If pregnant or breast-feeding, ask a health professional before use.

If Pregnant

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Drug Facts

Active ingredient (in each 10 mL)	Purpose
Dextromethorphan HBr, 30 mg.....	Cough Suppressant

Uses

• temporarily relieves: • cough due to minor throat and bronchial irritation associated with the common cold or inhaled irritants • the intensity of coughing • the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

In case of accidental overdose, get medical or professional help or contact a Poison Control center immediately

Indication and usage section

Do not take more than 4 doses in any 24 hours period

Use enclosed dosing cup

Keep dosing cup for use with this product

ml= milliliters

Adults and children 12 years & over 10 ml every 6-8 hours

children 6 to under 12 years of age 5 ml every 6-8 hours

children 4 to under 6 years of age 2.5 ml every 6-8

Drug Facts (continued).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 4 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL = milliliters

adults and children 12 years and over	10 mL every 6-8 hours
children 6 to under 12 years of age	5 mL every 6-8 hours
children 4 to under 6 years of age	2.5 mL every 6-8 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

natural and artificial flavor, natural grade A honey, methylparaben, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose.

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

hours

children under 4 years of age Consult a doctor

Other information

Other information

Store at room temperature 15-30 °C (59° -86 ° F)

Close cap tightly

Drug Facts (continued).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 4 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL = milliliters

adults and children 12 years and over	10 mL every 6-8 hours
children 6 to under 12 years of age	5 mL every 6-8 hours
children 4 to under 6 years of age	2.5 mL every 6-8 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

natural and artificial flavor, natural grade A honey, methylparaben, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose.

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Inactive ingredients

Inactive ingredients

natural and artificial flavor, natural grade A honey, methylparaben, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose

Drug Facts (continued).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 4 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL = milliliters

adults and children 12 years and over	10 mL every 6-8 hours
children 6 to under 12 years of age	5 mL every 6-8 hours
children 4 to under 6 years of age	2.5 mL every 6-8 hours
children under 4 years of age	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

natural and artificial flavor, natural grade A honey, methylparaben, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose.

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com



Dosage and administration

Do not take more than 4 doses in any 24 hours period

Use enclosed dosing cup

Keep dosing cup for use with this product

ml= milliliters

Adults and children 12 years & over 10 ml every 6-8 hours

children 6 to under 12 years of age 5 ml every 6-8 hours

children 4 to under 6 years of age 2.5 ml every 6-8

Keep out of the reach of children.



Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Principal panel

Maximum strength

Alcohol free, dye free, Non-Drowsy

The source Certified Honey

Giltus Honey DM

Cough

Dextromethorphan HBr, 30 mg

Persistent Cough

Up to 8 hours

Honey lemon flavor

4 fl oz 118 mL

MAXIMUM STRENGTH. MÁXIMA CONCENTRACIÓN.

ALCOHOL FREE.

DYE FREE.

NON-DROWSY.

SIN ALCOHOL.

SIN COLORANTES.

NO CAUSA SUEÑO.



Giltuss.

Honey DM



Cough TOS

Dextromethorphan HBr, 30 mg

**PERSISTENT COUGH
UP TO 8 HOURS**

**TOS PERSISTENTE
HASTA POR 8 HORAS**

HONEY LEMON FLAVOR. SABOR A MIEL Y LIMÓN.

4 FL OZ (118 mL)

GILTUSS HONEY DM COUGH

dextromethorphan hbr solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
HONEY (UNII: Y9H1V576FH)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	HONEY (Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65852-004-04	1 in 1 CARTON	06/22/2020	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/22/2020	

Revised: 9/2025

Dextrum Laboratories, Inc