

CHILDRENS GILTUSS EX EXPECTORANT- guaifenesin solution
Dextrum Laboratories Inc.

Children's Giltuss Ex Expectorant

Active Ingredients

Active ingredients (in each 5 ml) Purpose

Guaifenesin, 200 mg -----Expectorant

Drug Facts

<i>Active ingredient (in each 5 mL)</i>	<i>Purpose</i>
Guaifenesin, 200 mg	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Ask a doctor before use if your child has • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with asthma or chronic bronchitis

Stop use and ask a doctor if the child has a cough that lasts for more than 1 week, tends to recur or is accompanied by fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children. In case of overdose, get

medical help or contact a Poison Control Center right away.

Uses

Drug Facts

Active ingredient (in each 5 mL)

Purpose

Guaifenesin, 200 mg	Expectorant
---------------------------	-------------

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
-

Warnings

Ask a doctor before use if your child has • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with asthma or chronic bronchitis

Stop use and ask a doctor if the child has a cough that lasts for more than 1 week, tends to recur or is accompanied by fever, rash or persistent headache. A persistent cough may be a sign of

a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Warnings

Drug Facts

<i>Active ingredient (in each 5 mL)</i>	<i>Purpose</i>
Guaifenesin, 200 mg	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
-

Warnings

Ask a doctor before use if your child has • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with asthma or chronic bronchitis

Stop use and ask a doctor if the child has a cough that lasts

Stop use and ask a doctor if the child has a cough that lasts for more than 1 week, tends to recur or is accompanied by fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Stop use section

Drug Facts

<i>Active ingredient (in each 5 mL)</i>	<i>Purpose</i>
Guaifenesin, 200 mg	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
-

Warnings

Ask a doctor before use if your child has • cough that occurs with too much phlegm (mucus) • persistent or chronic

avoid use if you have too much phlegm (mucus) or persistent or chronic cough such as occurs with asthma or chronic bronchitis

Stop use and ask a doctor if the child has a cough that lasts for more than 1 week, tends to recur or is accompanied by fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children

Drug Facts

<i>Active ingredient (in each 5 mL)</i>	<i>Purpose</i>
Guaifenesin, 200 mg	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
-

Warnings

Ask a doctor before use if your child has • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with asthma or chronic bronchitis

Stop use and ask a doctor if the child has a cough that lasts for more than 1 week, tends to recur or is accompanied by fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Directions

Do not take more than 6 doses in any 24 hours period

use enclosed dosing cup

Keep dosage cup for use with this product

ml= milliliters

children under 4 years of age Consult a doctor

children 4 to under 6 years of age 2.5 ml every 4 hours

children 6 to under 12 years of age 5 ml every 4 hours

Adults and children 12 years & over 10 ml every 4 hours

Drug Facts (continued).

Directions

- do not take more than 6 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL=milliliters

children under 4 years of age	Consult a doctor
children 4 to under 6 years of age	2.5 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
adults and children 12 years and over	10 mL every 4 hours

Other information

- store at room temperature, 15°-30°C (59°- 86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

citric acid, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments?

Call **1-787-848-9114**, Monday–Friday 9AM–5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Other Information

Other information

Store at room temperature 15-30 °C (59° -86 ° F)

Close cap tightly

SAVE CARTON FOR COMPLETE DRUG FACTS

Drug Facts (continued).

Directions

- do not take more than 6 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL=milliliters

children under 4 years of age	Consult a doctor
children 4 to under 6 years of age	2.5 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
adults and children 12 years and over	10 mL every 4 hours

Other information

- store at room temperature, 15°-30°C (59°- 86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

citric acid, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments?

Call **1-787-848-9114**, Monday–Friday 9AM–5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Inactive Ingredients

Inactive Ingredients

Citric Acid, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose

Drug Facts (continued).

Directions

- do not take more than 6 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL=milliliters

children under 4 years of age	Consult a doctor
children 4 to under 6 years of age	2.5 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
adults and children 12 years and over	10 mL every 4 hours

Other information

- store at room temperature, 15°-30°C (59°- 86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

citric acid, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments?

Call **1-787-848-9114**, Monday-Friday 9AM-5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

Questions or comments

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM – 5PM EST

Call your doctor for medical advice in the event of side effects.

Drug Facts (continued).

Directions

- do not take more than 6 doses in any 24-hour period
- use enclosed dosing cup
- **keep dosage cup for use with this product**
- mL=milliliters

children under 4 years of age	Consult a doctor
children 4 to under 6 years of age	2.5 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
adults and children 12 years and over	10 mL every 4 hours

Other information

- store at room temperature, 15°-30°C (59°- 86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

citric acid, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments?

Call **1-787-848-9114**, Monday–Friday 9AM–5PM EST.

Call your doctor for medical advice in the event of side effects.



MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565

giltuss.com

indication for use section

indication an user section

Do not take more than 6 doses in any 24 hours period

use enclosed dosing cup

Keep dosage cup for use with this product

ml= milliliters

children under 4 years of age Consult a doctor

children 4 to under 6 years of age 2.5 ml every 4 hours

children 6 to under 12 years of age 5 ml every 4 hours

Adults and children 12 years & over 10 ml every 4 hours

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Guaifenesin, 200 mg	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Ask a doctor before use if your child has • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with asthma or chronic bronchitis

Stop use and ask a doctor if the child has a cough that lasts for more than 1 week, tends to recur or is accompanied by fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Children's Giltuss EX EXPECTORANT
Guaifenesin, 200 mg

Alcohol Free, Sugar Free, Dye Free, Non-Drowsy.
Sin Alcohol, Sin Azúcar, Sin Colorantes, No Causa Somnolencia.

CHEST CONGESTION, LIQUEFIES AND LOOSENS PHLEGM AND MUCUS
CONGESTION DE PECHO, LÍQUIDA Y SUELTA LA FLEMA Y EL MOCO

TUTTI FRUTTI FLAVOR, SANGRIA DE FRUTAS
4 FL OZ (118 mL)

Drug Facts (continued).

Directions

- do not take more than 6 doses in any 24-hour period
- use enclosed dosing cup
- keep dosage cup for use with this product
- mL=milliliters

children under 4 years of age	Consult a doctor
children 4 to under 6 years of age	2.5 mL every 4 hours
children 6 to under 12 years of age	5 mL every 4 hours
adults and children 12 years and over	10 mL every 4 hours

Other information

- store at room temperature, 15°-30°C (59°- 86°F)
- close cap tightly
- SAVE CARTON FOR COMPLETE DRUG FACTS

Inactive ingredients
citric acid, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments?
Call 1-787-848-9114, Monday-Friday 9AM-5PM EST.
Call your doctor for medical advice in the event of side effects.

MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1566

giltuss.com

Children 4 -12

Children's Giltuss EX

Expectorant

Guaifenesin, 200 mg

Alcohol free, Sugar free, Dye free, Non-Drowsy

Chest congestion liquifies and loosen phlegm and mucus

Tutti Frutti flavor

4 floz 118 ml



Children
Niños
4-12



Children's
Giltuss EX
EXPECTORANT
EXPECTORANTE

Guaifenesin, 200 mg

Alcohol Free. Sugar Free. Dye Free. Non-Drowsy.
Sin Alcohol. Sin Azúcar. Sin Colorantes. No Causa Sueño.

CHEST CONGESTION. LIQUIFIES
AND LOOSEN PHLEGM AND MUCUS
CONGESTIÓN DE PECHO. LICUA
Y SUELTA LA FLEMA Y EL MOCO



TUTTI FRUTTI FLAVOR.
SABOR A TUTTI FRUTTI.

4FL OZ (118 mL)

CHILDRENS GILTUSS EX EXPECTORANT

guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-016
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	
CITRIC ACID (UNII: 2968PHW8QP)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	TUTTI FRUTTI	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65852-016-04	1 in 1 CARTON	06/22/2020	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/22/2020	

Revised: 9/2025

Dextrum Laboratories Inc.