LEVOCETIRIZINE DIHYDROCHLORIDE - levocetirizine dihydrochloride tablet, film coated

A-S Medication Solutions

Levocetirizine Dihydrochloride Tablets USP, 5 mg (OTC)

ACTIVE INGREDIENT(S)

Levocetirizine dihydrochloride USP 5 mg

PURPOSE

Antihistamine

USE(S)

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

DO NOT USE

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

ASK A DOCTOR BEFORE USE IF YOU HAVE

ever had trouble urinating or emptying your bladder

WHEN USING THIS PRODUCT

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

STOP USE AND ASK A DOCTOR IF

• you have trouble urinating or emptying your bladder

• an allergic reaction to this product occurs. Seek medical help right away.

IF PREGNANT OR BREAST-FEEDING

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

adults 65 years of age and older	ask a doctor
adults and children 12 to 64 years of age	 take 1 tablet (5 mg) once daily in the evening do not take more than 1 tablet (5 mg) in 24 hours ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children 6 to 11 years of age	 take ½ tablet (2.5 mg) once daily in the evening do not take more than ½ tablet (2.5 mg) in 24 hours
children under 6 years of age	• do not use
consumers with kidney disease	■ do not use

OTHER INFORMATION

- store between 20° and 25°C (68° and 77°F)
- safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing
- safety sealed: do not use if carton was opened or if individual blister unit is open or torn

INACTIVE INGREDIENTS

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, titanium dioxide.

QUESTIONS or COMMENTS?

call **1-888-588-1418.**

Distributed by:

Camber Consumer Care, Inc.

Piscataway, NJ 08854, USA.

LEVOCETIRIZINE DIHYDROCHLORIDE TABLET, FILM COATED



levocetirizine dihydrochloride tablet, film coated

_			
Droc	liict	Inform	STIAN
FIUU	IULL		ашин

Product Type HUMAN OTC DRUG Item Code (Source) NDC:50090-5341(NDC:692)
--

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength LEVOCETIRIZINE DIHYDROCHLORIDE (UNII: SOD6A38AGA) (LEVOCETIRIZINE - UNII:6U5EA9RT2O) LEVOCETIRIZINE DIHYDROCHLORIDE 5 mg

Inactive Ingredients			
Ingredient Name	Strength		
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)			
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ 8WG20P6)			
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			

Product Characteristics			
Color	WHITE (White to off white)	Score	2 pieces
Shape	OVAL	Size	8mm
Flavor		Imprint Code	H;LL
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50090- 5341-0	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/04/2020	
2	NDC:50090- 5341-1	90 in 1 BOTTLE; Type 0: Not a Combination Product	11/04/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA213513	10/28/2020	

Labeler - A-S Medication Solutions (830016429)

Establishment				
Name	Address	ID/FEI	Business Operations	
A-S Medication Solutions		830016429	RELABEL(50090-5341), REPACK(50090-5341)	

Revised: 4/2021 A-S Medication Solutions