


DOMETUSS- chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

Dextrum Laboratories, Inc

Dometuss



Anthistamine


cough suppresant
nasal descongstant.

Use

For temporary relief of runny nose, sneezing, itching of the nose orthroat and itchywatery eyes due to hay fever.

Temporary relieves cough due to minor throat and bronchial irritation as may occur with the common cold.

temporary restores freer breathing through the nose.



Active Ingredients (in each

5ml-teaspoonful)

Chlorpheniramine Maleate 2mg

Dextromethorphan HBr 10mg

Phenylephrine HCl 5mg

- To sedate a child or to make a child sleepy.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Certain drugs for depression, psychiatric or emotional condition or parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts:

Active ingredients in each 5-ml teaspoonful:
 Chlorpheniramine Maleate 2mg Antihistamine
 Codeine Phosphate 10mg Cough Suppressant
 Phenylephrine HCl 3mg Nasal Decongestant

Uses: Temporary relief of runny nose, sneezing, itching of the nose or throat and dry, itchy watery eyes due to hay fever, allergic rhinitis, or the common cold.

Warnings: Do not use:
 • If you are taking MAO inhibitors (e.g., phenelzamine, tranylcypromine) or have taken them in the last 14 days.
 • If you have taken a monoamine oxidase (MAO) inhibitor in the last 14 days.
 • If you are taking a MAO inhibitor (e.g., phenelzamine, tranylcypromine) or have taken them in the last 14 days.
 • If you are taking a MAO inhibitor (e.g., phenelzamine, tranylcypromine) or have taken them in the last 14 days.

Ask a doctor before use if you have:
 • Glaucoma, narrow-angle glaucoma, or a history of glaucoma.
 • High blood pressure, or a history of high blood pressure.
 • Heart disease, or a history of heart disease.
 • Kidney disease, or a history of kidney disease.
 • Liver disease, or a history of liver disease.
 • Stomach or duodenal ulcer, or a history of stomach or duodenal ulcer.
 • Thyroid disease, or a history of thyroid disease.
 • Urinary tract infection, or a history of urinary tract infection.

Ask a doctor before use if you are:
 • Pregnant, or planning to become pregnant.
 • Breast-feeding, or planning to breast-feed.

Other important information:
 • Do not use for more than 7 days. Continued use for more than 7 days may worsen your condition.
 • Do not use for more than 7 days. Continued use for more than 7 days may worsen your condition.
 • Do not use for more than 7 days. Continued use for more than 7 days may worsen your condition.

Directions: Do not exceed more than 6 doses in any 24 hour period.

Adults and children 12 years of age and over: 2 teaspoonful (10 mL) every 4 hours.

Children 6 to under 12 years of age: 1 teaspoonful (5 mL) every 4 hours.

Children under 6 years of age: Ask a doctor.

Other important information: Do not use for more than 7 days. Continued use for more than 7 days may worsen your condition.

Keep this and all other medicines out of the reach of children. Do not use this product if aluminum foil over bottle opening is torn, broken or missing.

Contains 100 mg of Codeine Phosphate (100 mg) per 5 mL.

Lot#: _____
Exp. Date: _____

DOMETUSS

**Antihistamine/Cough Suppressant/
Nasal Decongestant**

Sugar, Alcohol, And Dye Free

Grape Flavor

4 fl. oz (118 mL)

Drug Facts (continued)

Directions: Do not exceed more than 6 doses in any 24 hour period.

Adults and children 12 years of age and over: 2 teaspoonful (10 mL) every 4 hours.

Children 6 to under 12 years of age: 1 teaspoonful (5 mL) every 4 hours.

Children under 6 years of age: Ask a doctor.

Other important information: Do not use for more than 7 days. Continued use for more than 7 days may worsen your condition.

Keep this and all other medicines out of the reach of children. Do not use this product if aluminum foil over bottle opening is torn, broken or missing.

Contains 100 mg of Codeine Phosphate (100 mg) per 5 mL.

Lot#: _____
Exp. Date: _____

Ask a Doctor Before use if you have heart disease, thyroid disease trouble urinating due to enlarged prostate gland, glaucoma, cough that occurs with too much phlegm (mucus), a breathing problems or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema.

REV-12-2021
Lot#:
Exp.Date

When using this product do not exceed recommended dosage
may cause marked drowsiness, avoid alcoholic beverages
alcohol, sedatives and tranquilizers may cause increased drowsiness
be careful when driving a motor vehicle or operating machinery
excitability may occur, especially with children

Drug Facts (continued)	
Directions: Do not exceed more than 6 doses in any 24 hour period.	
Adults and children 12 years of age and older	2 heparinoid (10 mg) every 6 hours
Children 6 to under 12 years of age	1 heparinoid (5 mg) every 6 hours
Children under 6 years of age	Ask a doctor
Other information: Store at room temperature between 15° and 30°C (59° and 86°F). For more information, do not use the product if aluminum foil over bottle opening is torn, broken or missing.	
Contains inactive ingredients: Citric acid, flavor, methylparaben, propylparaben, propylene glycol, purified water, sodium citrate and sodium benzoate.	
Questions or Comments? Please call (772) 355-3555.	

Stop use and ask a doctor if:

You get nervous, dizzy, or sleepless.

Cough last more than 7 days, comes back, or is accompanied by fever, rash or persistent headaches. These could be signs of a serious condition.

If pregnant

Drug Facts:

Active ingredients in each 5-ml teaspoonful: Antihistamine Chlorpheniramine Maleate 2mg Cough Suppressant Dextromethorphan HBr 10mg Nasal Decongestant Phenylephrine HCl 5mg

Uses: temporary relief of runny nose, sneezing, itching of the nose or throat and dry cough when due to hay fever, colds, or allergies. It also helps relieve nasal congestion.

Warnings: Do not use with other cold or cough medicines. Do not use if you have taken MAO inhibitors (drugs used to treat depression) within the last 14 days. Do not use if you are taking any medicine to treat high blood pressure. Do not use if you are taking any medicine to treat heart disease. Do not use if you are taking any medicine to treat diabetes. Do not use if you are taking any medicine to treat thyroid disease. Do not use if you are taking any medicine to treat seizures. Do not use if you are taking any medicine to treat blood thinning. Do not use if you are taking any medicine to treat urinary problems. Do not use if you are taking any medicine to treat stomach problems. Do not use if you are taking any medicine to treat kidney problems. Do not use if you are taking any medicine to treat liver problems. Do not use if you are taking any medicine to treat prostate problems. Do not use if you are taking any medicine to treat vision problems. Do not use if you are taking any medicine to treat hearing problems. Do not use if you are taking any medicine to treat sleep problems. Do not use if you are taking any medicine to treat anxiety problems. Do not use if you are taking any medicine to treat depression problems. Do not use if you are taking any medicine to treat bipolar problems. Do not use if you are taking any medicine to treat schizophrenia problems. Do not use if you are taking any medicine to treat mental problems. Do not use if you are taking any medicine to treat physical problems. Do not use if you are taking any medicine to treat chronic problems. Do not use if you are taking any medicine to treat acute problems. Do not use if you are taking any medicine to treat any other problems.

Directions: Adults and children 12 years of age and over: 2 teaspoonful (10 mL) every 4 hours. Children 6 to under 12 years: 1 teaspoonful (5 mL) every 4 hours. Children under 6 years of age: Ask a doctor.

Other information: Store at room temperature between 15°-30°C (59°-86°F). Keep out of reach of children. Do not use this product if aluminum foil over bottle opening is torn, broken or missing.

Contains ingredients: Citric acid, flavor, methylparaben, propylparaben, propylene glycol, purified water, sodium citrate and sucrose.

Contains or Contains? Phenol cod (732) 755-1255.

Lot#: _____
Exp. Date: _____

NDC 53809-222-04

DOMETUSS

**Antihistamine/Cough Suppressant/
Nasal Decongestant**



Sugar, Alcohol, And Dye
Free
 Grape Flavor
 4 fl. oz (118 mL)

Dometic, Manufactured for:
 SAN JUAN PHARM 800-838-0100
 0 53809-222-04 3

If pregnant or breast feeding ask a health professional before use.

Keep Out

Directions

Drug Facts:

Active ingredients in each 5-ml teaspoonful:
 Chlorpheniramine Maleate 2mg Antihistamine
 Codeine Phosphate 10mg Cough Suppressant
 Pseudoephedrine HCl 30mg Nasal Decongestant

Uses: temporary relief of runny nose, sneezing, itching of the nose or throat and thin watery eyes due to hay fever, colds, or allergies.
 *Temporarily relieves nasal congestion through the nose.

Warnings: Do not use:
 • if you are allergic to any of the ingredients.
 • if you have taken more than 100 mg of pseudoephedrine in the last 24 hours.
 • if you are taking any medicine that contains pseudoephedrine or phenylephrine.
 • if you are taking any medicine that contains codeine.
 • if you are taking any medicine that contains chlorpheniramine.
 • if you are taking any medicine that contains a monoamine oxidase inhibitor (MAOI).
 • if you are taking any medicine that contains a tricyclic antidepressant (TCA).
 • if you are taking any medicine that contains a sedative or hypnotic.
 • if you are taking any medicine that contains a muscle relaxant.
 • if you are taking any medicine that contains a blood thinner.
 • if you are taking any medicine that contains a diuretic.
 • if you are taking any medicine that contains a heart medicine.
 • if you are taking any medicine that contains a blood pressure medicine.
 • if you are taking any medicine that contains a diabetes medicine.
 • if you are taking any medicine that contains a thyroid medicine.
 • if you are taking any medicine that contains a steroid.
 • if you are taking any medicine that contains a vitamin A or D supplement.
 • if you are taking any medicine that contains a calcium supplement.
 • if you are taking any medicine that contains a potassium supplement.
 • if you are taking any medicine that contains a magnesium supplement.
 • if you are taking any medicine that contains a zinc supplement.
 • if you are taking any medicine that contains a selenium supplement.
 • if you are taking any medicine that contains a chromium supplement.
 • if you are taking any medicine that contains a manganese supplement.
 • if you are taking any medicine that contains a cobalt supplement.
 • if you are taking any medicine that contains a molybdenum supplement.
 • if you are taking any medicine that contains a silicon supplement.
 • if you are taking any medicine that contains a boron supplement.
 • if you are taking any medicine that contains a fluorine supplement.
 • if you are taking any medicine that contains a phosphorus supplement.
 • if you are taking any medicine that contains a sulfur supplement.
 • if you are taking any medicine that contains a chlorine supplement.
 • if you are taking any medicine that contains a bromine supplement.
 • if you are taking any medicine that contains a nitrogen supplement.
 • if you are taking any medicine that contains a carbon supplement.
 • if you are taking any medicine that contains a hydrogen supplement.
 • if you are taking any medicine that contains an oxygen supplement.
 • if you are taking any medicine that contains a fluorine supplement.
 • if you are taking any medicine that contains a phosphorus supplement.
 • if you are taking any medicine that contains a sulfur supplement.
 • if you are taking any medicine that contains a chlorine supplement.
 • if you are taking any medicine that contains a bromine supplement.
 • if you are taking any medicine that contains a nitrogen supplement.
 • if you are taking any medicine that contains a carbon supplement.
 • if you are taking any medicine that contains a hydrogen supplement.
 • if you are taking any medicine that contains an oxygen supplement.

How to use: Adults and children 12 years of age and older: 1 teaspoonful (5 ml) 4 times a day.
 Children 6 to under 12 years: 1 teaspoonful (5 ml) 4 times a day.
 Children under 6 years of age: Ask a doctor.

Other important information: Do not use this product if aluminum foil over bottle opening is torn, broken or missing.

Keep this and all medicines out of the reach of children. Do not use this product if aluminum foil over bottle opening is torn, broken or missing.

Inactive ingredients: Citric acid, flavor, methylparaben, propylparaben, propylene glycol, purified water, sodium citrate and sucrose.

Contains 4.2 fluid ounces (122 mL) of liquid.

Lot#: _____
Exp. Date: _____

DOMETUSS
 Antihistamine/Cough Suppressant/
 Nasal Decongestant

DOMETL Manufactured for:
 SAN JUAN PHARMACEUTICALS
 0153809-2220-04



Sugar, Alcohol, And Dye
Free
 Grape Flavor
 4 fl. oz (118 ml)

Drug Facts (continued)

Directions: Do not exceed more than 6 doses in any 24 hour period.

Adults and children 12 years of age and older	2 teaspoonful (10 mL) every 4 hours
Children 6 to under 12 years of age	1 teaspoonful (5 mL) every 4 hours
Children under 6 years of age	Ask a doctor

Other important information: Do not use this product if aluminum foil over bottle opening is torn, broken or missing.


Keep this and all medicines out of the reach of children. Do not use this product if aluminum foil over bottle opening is torn, broken or missing.

Inactive ingredients: Citric acid, flavor, methylparaben, propylparaben, propylene glycol, purified water, sodium citrate and sucrose.

Contains 4.2 fluid ounces (122 mL) of liquid.

Lot#: _____
Exp. Date: _____

[illegible]



Dometuss

Antihistamine/Cough Suppressant/Nasal Decongestant
Sugar, Alcohol, And Dye
Free
Grape Flavor
4 fl oz (118 ml)

DOMETUSS			
chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-002
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE MONOHYDRATE (UNII: X6JU3FX5UU)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65852-002-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/21/2016	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/21/2016	

Labeler - Dextrum Laboratories, Inc (007392322)

Establishment			
Name	Address	ID/FEI	Business Operations
Dextrum Laboratories, Inc		007392322	manufacture(65852-002)

Revised: 9/2025

Dextrum Laboratories, Inc