

DG HEALTH ACID REDUCER- famotidine tablet, film coated
Dolgenercorp Inc

Dolgenercorp, LLC Acid Reducer Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

Questions or comments?

1-888-309-9030

Package/Label Principal Display Panel

DG™ |health

Compare to the active ingredient of Maximum Strength Zantac 360™

Maximum Strength

Acid Reducer

Famotidine Tablets, 20 mg

Acid Reducer

Just one tablet PREVENTS & RELIEVES HEARTBURN DUE TO ACID INDIGESTION

50 Tablets

Actual Tablet Size

100%
Satisfaction
Guaranteed!
(888) 309-9030

**DO NOT USE IF PRINTED FOIL UNDER
CAP IS BROKEN OR MISSING**
DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN37072



ACTUAL SIZE



Compare to the active ingredient of Maximum Strength Zantac 360™*

**Maximum Strength
Acid Reducer**

Famotidine Tablets, 20 mg
Acid Reducer

Just one tablet
**PREVENTS & RELIEVES
HEARTBURN DUE TO
ACID INDIGESTION**



15471 VT C1



A1499

JUST ONE TABLET prevent and relieve heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.


Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried food, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts		Drug Facts (continued)	
Active ingredient (in each tablet) famotidine 20 mg	Purpose Acid reducer	<ul style="list-style-type: none"> heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease 	<ul style="list-style-type: none"> store at 20°-25° C (68°-77° F) protect from moisture
Uses	<ul style="list-style-type: none"> relieve heartburn associated with acid indigestion and sour stomach prevent heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages 	<p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days <p>If pregnant or breastfeeding, ask health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. (1-800-222-1222)</p>	<p>Directions</p> <ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swallow 1 tablet with a glass of water anytime from 10 to 60 minutes before eating food or drinking beverages that
Warnings	<p>Allergy alert Do not use if you are allergic to famotidine or other acid reducers.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have trouble or pain swallowing food, vomiting with blood or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. 	Drug Facts (continued)	Directions (continued)
		<p>Warnings (continued)</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. 	<p>Directions (continued)</p> <p>to prevent symptoms, swallow 1 tablet with a glass of water anytime from 10 to 60 minutes before eating food or drinking beverages that</p>

**Maximum Strength
Acid Reducer**
Famotidine Tablets, 20 mg
Acid Reducer

Directions	Warnings (continued)
<ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swallow 1 tablet with a glass of water anytime from 10 to 60 minutes before eating food or drinking beverages that 	<ul style="list-style-type: none"> store at 20°-25° C (68°-77° F) protect from moisture <p>In active ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc,</p>



DG HEALTH ACID REDUCER

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-814
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L194
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-814-71	1 in 1 CARTON	06/15/2022	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	06/15/2022	

Labeler - Dolgencorp Inc (068331990)

Revised: 7/2022

Dolgencorp Inc