

**DAYTIME SEVERE COLD AND FLU RELIEF SOFTGELS- acetaminophen,
dextromethorphan hydrobromide, guaifenesin, phenylephrine
hydrochloride capsule, liquid filled
Topco Associates LLC**

TopCare® health DayTime *SEVERE* Cold & Flu Relief SOFTGELS

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sinus congestion & pressure
 - cough due to minor throat & bronchial irritation
 - minor aches & pains
 - headache
 - fever
 - sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin; taking sedatives or tranquilizers.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
 - pain, nasal congestion, or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at controlled room temperature 20-25°C (68-77°F) • protect from light, heat and moisture

Inactive ingredients

FD&C Yellow No.6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

Call **1-888-423-0139**

COMPARE TO VICKS® DAYQUIL™ SEVERE COLD & FLU LIQUICAPS™ ACTIVE INGREDIENTS*

MULTI-SYMPTOM RELIEF

Antihistamine Free


READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION.

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ITASCA, IL 60143 ©TOPCO
STVA0825 topcare@topco.com www.topcarebrand.com
PRODUCT OF INDIA

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks®, DayQuil™ and LiquiCaps™.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

Packaging





Drug Facts Label

Drug Facts	
Active ingredients (in each softgel)	
Acetaminophen 325 mg.....	Pain reliever/fever reducer.
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Drug Facts (continued)
Uses
- temporarily relieves common cold/flu symptoms: - nasal congestion - sinus congestion & pressure - cough due to minor throat & bronchial irritation - minor aches & pains - headache - fever - sore throat - reduces swelling of nasal passages - temporarily restores freer breathing through the nose - promotes nasal and/or sinus drainage - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: - more than 8 softgels in 24 hours, which is the maximum daily amount for this product - with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: - skin reddening - blisters - rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
--

Drug Facts (continued)
Ask a doctor before use if you have - liver disease - heart disease - high blood pressure - thyroid disease - diabetes - glaucoma - trouble urinating due to enlarged prostate gland - cough that occurs with too much phlegm (mucus) - persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin; taking sedatives or tranquilizers. When using this product, do not use more than directed. Stop use and ask a doctor if - you get nervous, dizzy or sleepless - pain, nasal congestion, or cough gets worse or lasts more than 7 days - fever gets worse or lasts more than 3 days - redness or swelling is present - new symptoms occur - cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away.

Directions						
- take only as directed - do not exceed 8 softgels per 24 hrs <table border="1"> <tbody> <tr> <td>adults & children 12 yrs & over</td> <td>2 softgels with water every 4 hours</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </tbody> </table>	adults & children 12 yrs & over	2 softgels with water every 4 hours	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	2 softgels with water every 4 hours					
children 4 to under 12 yrs	ask a doctor					
children under 4 yrs	do not use					

Drug Facts (continued)
Other information ■ store at controlled room temperature 20–25°C (68–77°F) ■ protect from light, heat, and moisture
Inactive ingredients
FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide
Questions or comments? Call 1-888-423-0139

DAYTIME SEVERE COLD AND FLU RELIEF SOFTGELS

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76162-636
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients	
Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	A13
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-636-24	2 in 1 CARTON	12/11/2025	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/11/2025	

Labeler - Topco Associates LLC (006935977)

Revised: 12/2025

Topco Associates LLC