

SODIUM SULFACETAMIDE 9.8% SULFUR 4.8% CLEANSER- sodium sulfacetamide 9.8% sulfur 4.8% liquid
Oncor Pharmaceuticals


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Sodium Sulfacetamide 9.8% Sulfur 4.8% Cleanser

DESCRIPTION

Sodium sulfacetamide is a sulfonamide with antibacterial activity while sulfur acts as a keratolytic agent.

Chemically sodium sulfacetamide is N-[(4-aminophenyl) sulfonyl]-acetamide, monosodium salt, monohydrate. The structural formula is:



Each gram of Sodium Sulfacetamide 9.8% and Sulfur 4.8% Cleanser contains 98 mg of sodium sulfacetamide and 48 mg of sulfur in a cleanser containing Aloe vera leaf extract, Butylated hydroxytoluene, Cetyl alcohol, Citric acid, Cocamidopropyl betaine, Disodium EDTA, Glycerin, Glyceryl stearate SE, PEG-100 stearate, Phenoxyethanol, Purified water, Sodium laureth sulfate, Sodium thiosulfate, Stearyl alcohol, Triacetin, Xanthan gum.

INDICATIONS

Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is indicated in the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis.

DOSAGE AND ADMINISTRATION

Wash affected area once or twice daily, or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin for 10-20 seconds working into a full lather, rinse thoroughly and pat dry. If drying occurs, it may be controlled by rinsing cleanser off sooner or using less often.

FOR EXTERNAL USE ONLY. NOT FOR INTRA VAGINAL OR OPHTHALMIC USE.
(KEEP AWAY FROM EYES).

KEEP OUT OF REACH OF CHILDREN.

Shake well before use

CONTRAINDICATIONS

Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is contraindicated for use by patients having known hypersensitivity to sulfonamides, sulfur or any other component of this preparation. Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is not to be used by patients with kidney disease.

CAUTION

If redness or irritation occurs, discontinue use.

STORAGE

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized.

In case of accidental ingestion contact a Poison Control Center immediately. Keep container tightly closed.

You may report side effects by calling Oncor Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.

CLINICAL PHARMACOLOGY

Sodium sulfacetamide exerts a bacteriostatic effect against sulfonamide sensitive gram-positive and gram-negative microorganisms commonly isolated from secondary cutaneous pyogenic infections. It acts by restricting the synthesis of folic acid required by bacteria for growth, by its competition with para-aminobenzoic acid. There is no clinical data available on the degree and rate of systemic absorption of this product when applied to the skin or scalp. However, significant absorption of sodium sulfacetamide through the skin has been reported.

The following in vitro data is available but the clinical significance is unknown. Organisms that show susceptibility to sodium sulfacetamide are: Streptococci, Staphylococci, E. coli, Klebsiella pneumoniae, Pseudomonas pyocyanea, Salmonella species, Proteus vulgaris, Nocardia and Actinomyces.

The exact mode of action of sulfur in the treatment of acne is unknown, but it has been reported that it inhibits the growth of Propionibacterium acnes and the formation of free fatty acids.

WARNINGS

Although it is rare, sensitivity to sodium sulfacetamide may occur. Therefore, caution and careful supervision should be observed when prescribing this drug for patients who may be prone to hypersensitivity to topical sulfonamides. Systemic toxic reactions such as agranulocytosis, acute hemolytic anemia, purpura hemorrhagica, drug fever, jaundice, and contact dermatitis indicate hypersensitivity to sulfonamides. Particular caution should be employed if areas of denuded or abraded skin are involved.

PRECAUTIONS

GENERAL

If irritation develops, use of the product should be discontinued and appropriate therapy instituted. Patients should be carefully observed for possible local irritation or sensitization during long-term therapy. The object of this therapy is to achieve desquamation without irritation, but sodium sulfacetamide and sulfur can cause reddening and scaling of the epidermis. These side effects are not unusual in the treatment of acne vulgaris, but patients should be cautioned about the possibility.

INFORMATION FOR PATIENTS

Avoid contact with eyes, eyelids, lips and mucous membranes. If accidental contact occurs, rinse with water. If excessive irritation develops, discontinue use and consult your physician.

CARCINOGENESIS, MUTAGENESIS AND IMPAIRMENT OF FERTILITY

Long-term studies in animals have not been performed to evaluate carcinogenic potential.

PREGNANCY

Category C

Animal reproduction studies have not been conducted with Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser. It is also not known whether Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser should be given to a pregnant woman only if clearly needed.

NURSING MOTHERS

It is not known whether sodium sulfacetamide is excreted in the human milk following topical use of Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser. However, small amounts of orally administered sulfonamides have milk. In view of this and because many drugs are excreted in human milk, caution should be exercised when Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is administered to a nursing woman.

PEDIATRIC USE

Safety and effectiveness in children under the age of 12 have not been established.

ADVERSE REACTIONS

Although rare, sodium sulfacetamide may cause local irritation. Call your doctor for medical advice about side effects.

HOW SUPPLIED

Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is available:
16 oz (454 g) bottle, NDC 83720-546-16.

NOTICE

Protect from freezing and excessive heat. The product may tend to darken slightly on storage. Slight discoloration does not impair the efficacy or safety of the product. Keep bottle tightly closed.

Occasionally, a slight discoloration of fabric may occur when an excessive amount of the product is used and comes in contact with white fabrics. This discoloration, however, presents no problem, as it is readily removed by ordinary laundering without bleaches.

Manufactured In USA For :
Oncor Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States.
Rev. 08/25

NCOR PHARMACEUTICALS
NDC 83720-546-16
Rx Only

Sodium Sulfacetamide & Sulfur
Sodium Sulfacetamide 9.8%
Sulfur 4.8%

Cleanser

For External Use Only
NET WT. 16 OZ. (454 g)

INGREDIENTS
Each gram of Sodium Sulfacetamide 9.8% and Sulfur 4.8% Cleanser contains 98 mg of sodium sulfacetamide and 48 mg of sulfur in a cleanser containing Aloe vera leaf extract, Butylated hydroxytoluene, Cetyl alcohol, Citric acid, Cocamidopropyl betaine, Disodium EDTA, Glycerin, Glyceryl stearate SE, PEG-100 stearate, Phenoxethanol, Purified water, Sodium lauryl sulfate, Sodium thiosulfate, Stearyl alcohol, Tricetin, Xanthan gum.

INDICATIONS
Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is indicated in the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis.

DOSEAGE AND ADMINISTRATION
Wash affected area once or twice daily, or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleaned, massage gently into skin for 10-20 seconds working into a full lather, rinse thoroughly and pat dry. If drying occurs, it may be controlled by rinsing cleanser off sooner or using less often.

FOR EXTERNAL USE ONLY NOT FOR INTRA VAGINAL OR OPHTHALMIC USE. (KEEP AWAY FROM EYES).
KEEP OUT OF REACH OF CHILDREN.
Shake well before use

CONTRAINDICATIONS
Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is contraindicated for use by patients having known hypersensitivity to sulfonamides, sulfur or any other component of this preparation. Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is not to be used by patients with kidney disease.

STORAGE
Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized.

In case of accidental ingestion contact a Poison Control Center immediately. Keep container tightly closed.

You may report side effects by calling Oncor Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.

This bottle is not filled to the top but does contain 16 oz of product as identified on the front panel of the bottle.

Manufactured in USA For :
Oncor Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States.
Rev. 08/25

DESCRIPTION
Sodium sulfacetamide is a sulfonamide with antibacterial activity while sulfur acts as a keratolytic agent.

Chemically sodium sulfacetamide is N-(4-aminophenyl) sulfonylacetamide, monosodium salt, monohydrate. The structural formula is:

$$\text{NH}_2 - \text{C}_6\text{H}_4 - \text{SO}_2\text{N}(\text{COOCH}_2\text{H}_5)_2$$

CLINICAL PHARMACOLOGY
Sodium sulfacetamide exerts a bacteriostatic effect against sulfonamide sensitive gram-positive and gram-negative microorganisms commonly isolated from secondary cutaneous pyogenic infections. It acts by restricting the synthesis of folic acid required by bacteria for growth, by its competition with para-aminobenzoic acid. There is no clinical data available on the degree and rate of systemic absorption of this product when applied to the skin or scalp. However, significant absorption of sodium sulfacetamide through the skin has been reported.

The following in vitro data is available but the clinical significance is unknown. Organisms that show susceptibility to sodium sulfacetamide are: Streptococci, Staphylococci, E. coli, Klebsiella pneumoniae, Pseudomonas pyocyanus, Salmonella species, Proteus vulgaris, Neocardia and Actinomyces.

The exact mode of action of sulfur in the treatment of acne is unknown, but it has been reported that it inhibits the growth of Propionibacterium acnes and the formation of free fatty acids.

WARNINGS
Although it is rare, sensitivity to sodium sulfacetamide may occur. Therefore, caution and careful supervision should be observed when prescribing this drug for patients who may be prone to hypersensitivity to topical sulfonamides. Systemic toxic reactions such as agranulocytosis, acute hemolytic anemia, purpura hemorrhagica, drug fever, jaundice, and contact dermatitis indicate hypersensitivity to sulfonamides. Particular caution should be employed if areas of denuded or abraded skin are involved.

PRECAUTIONS
GENERAL
If irritation develops, use of the product should be discontinued and appropriate therapy instituted. Patients should be carefully observed for possible local irritation or sensitization during long-term therapy. The object of this therapy is to achieve desquamation without irritation, but sodium sulfacetamide and sulfur can cause redness and scaling of the epidermis. These side effects are not unusual in the treatment of acne vulgaris, but patients should be cautioned about the possibility.

INFORMATION FOR PATIENTS
Avoid contact with eyes, eyelids, eyelids, lips and mucous membranes. If accidental contact occurs, rinse with water. If excessive irritation develops, discontinue use and consult your physician.

CARCINOGENESIS, MUTAGENESIS AND IMPAIRMENT OF FERTILITY
Long-term studies in animals have not been performed to evaluate carcinogenic potential.

PREGNANCY
Category C
Animal reproduction studies have not been conducted with Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser. It is also not known whether Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser should be given to a pregnant woman only if clearly needed.

NURSING MOTHERS
It is not known whether sodium sulfacetamide is excreted in the human milk following topical use of Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser. However, small amounts of orally administered sulfonamides have milk. In view of this and because many drugs are excreted in human milk, caution should be exercised when Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is administered to a nursing woman.

PEDIATRIC USE
Safety and effectiveness in children under the age of 12 have not been established.

ADVERSE REACTIONS
Although rare, sodium sulfacetamide may cause local irritation. Call your doctor for medical advice about side effects.

CAUTION
If redness or irritation occurs, discontinue use.

HOW SUPPLIED
Sodium Sulfacetamide 9.8% & Sulfur 4.8% Cleanser is available: 16 oz (454 g) bottle, NDC 83720-546-16.

NOTICE
Protect from freezing and excessive heat. The product may tend to darken slightly on storage. Slight discoloration does not impair the efficacy or safety of the product. Keep bottle tightly closed.

Occasionally, a slight discoloration of fabric may occur when an excessive amount of the product is used and comes in contact with white fabrics. This discoloration, however, presents no problem, as it is readily removed by ordinary laundering without bleaches.

Manufactured in USA For :
Oncor Pharmaceuticals
6755 Business Pkwy Suite 202
Elkridge, MD 21075
United States.
Rev. 08/25

SODIUM SULFACETAMIDE 9.8% SULFUR 4.8% CLEANSER

sodium sulfacetamide 9.8% sulfur 4.8% liquid

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83720-546
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SULFACETAMIDE SODIUM (UNII: 4NRT660KJQ) (SULFACETAMIDE - UNII:4965G3J0F5)	SULFACETAMIDE SODIUM	98 mg in 1 g

SULFUR (UNII: 70FD1KFU70) (SULFUR - UNII:70FD1KFU70)	SULFUR	48 mg in 1 g
---	--------	-----------------

Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
CITRIC ACID (UNII: 2968PHW8QP)	
COCAMIDOPROPYL BETAINE (UNII: 5OCF3O11KX)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	
GLYCERYL STEARATE SE (UNII: FCZ5MH785I)	
PEG-100 STEARATE (UNII: YD01N1999R)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
WATER (UNII: 059QF0KO0R)	
SODIUM LAURETH SULFATE (UNII: BPV390UAP0)	
SODIUM THIOSULFATE (UNII: HX1032V43M)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83720-546-16	454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/14/2025	
2	NDC:83720-546-10	285 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/14/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/14/2025	

Labeler - Oncor Pharmaceuticals (119032580)

Registrant - Oncor Pharmaceuticals (119032580)