

BACK AND MUSCLE PAIN- naproxen sodium tablet, film coated
CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED

CVS 44-417BackandMuscle

Active ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - the common cold
 - toothache
 - headache
 - menstrual cramps
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- skin reddening
- shock
- blisters
- rash
- facial swelling
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - leg swelling
 - chest pain
 - slurred speech
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- adults and children 12 years and older
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal display panel

NDC 51316-817-52

CVS™

STRENGTH

TO LAST

12

HOURS

actual size

**Back &
Muscle
Pain**

**Naproxen Sodium Tablets
220 mg (NSAID)
Pain Reliever / Fever Reducer**

Compare to Aleve®
Back & Muscle Pain Tablets
active ingredient†

90 TABLETS

100% money back
guaranteed.
[CVS.com/returnpolicy](https://www.cvs.com/returnpolicy)

†This product is not manufactured or distributed
by Bayer HealthCare LLC, owner of the registered
trademark Aleve® Back & Muscle Pain Tablets.
50844 ORG122141752

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2025 CVS/pharmacy
[CVS.com](https://www.cvs.com)® 1-800-SHOP CVS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

100% money back guarantee
 CVS.com/returnpolicy
 100% money back guarantee
 CVS.com/returnpolicy
 V-11112
 © 2025 CVS/pharmacy
 CVS.com 1-800-SHOP CVS
 Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 50844 086122141752
 Muscle Pain Tablets
 of the registered trademark Aleve® Back &
 distributed by Bayer HealthCare LLC, owner
 This product is not manufactured or

Drug Facts KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each tablet) Purpose
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ backache ■ muscular aches ■ the common cold
 ■ toothache ■ menstrual cramps
 ■ headache ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
 Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ asthma (wheezing) ■ skin reddening ■ shock
 ■ blisters ■ rash ■ facial swelling ■ hives
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ take more or for a longer time than directed
 ■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery
Ask a doctor before use if
 ■ the stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
 ■ taking any other drug
When using this product
 ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ you have symptoms of heart problems or stroke:
 ■ leg swelling ■ chest pain
 ■ slurred speech ■ trouble breathing
 ■ weakness in one part or side of body
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ you have difficulty swallowing
 ■ it feels like the pill is stuck in your throat
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and older
 ■ take 1 tablet every 8 to 12 hours while symptoms last
 ■ for the first dose you may take 2 tablets within the first hour
 ■ do not exceed 2 tablets in a 24-hour period
 ■ do not exceed 3 tablets in a 24-hour period
 ■ children under 12 years: ask a doctor

Other information
 ■ each tablet contains: sodium 20 mg
 ■ store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
 ■ see end flap for expiration date and lot number

Inactive ingredients
 croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?
 Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday



NDC 51316-817-52



actual size

BACK & MUSCLE PAIN

Naproxen Sodium Tablets
 220 mg (NSAID)
 Pain Reliever / Fever Reducer

Compare to Aleve®
 Back & Muscle Pain Tablets
 active ingredient!

90 TABLETS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

#340693

FPO 100%
 UPC# 050428165805

B-0231-417BM-52-J
 ORG122141752

X XXXXXX XXXXXX X

No Print/No Varnish
 Lot & EXP

CVS 44-417

BACK AND MUSCLE PAIN

naproxen sodium tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-817
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;417
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-817-08	1 in 1 CARTON	04/18/2022	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:51316-817-52	1 in 1 CARTON	04/18/2022	
2		90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:51316-817-15	1 in 1 CARTON	04/18/2022	08/08/2024
3		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204872	04/18/2022	

Labeler - CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(51316-817)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(51316-817)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(51316-817)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(51316-817)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(51316-817)

Revised: 7/2025

CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED