

TERRAMYCIN 200- oxytetracycline powder
Phibro Animal Health

Terramycin® 200
(oxytetracycline)
TYPE A MEDICATED ARTICLE

Active Drug Ingredient:

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to
oxytetracycline hydrochloride 200 g/lb

CAUTION:

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

CAUTION: For use in manufacturing medicated animal feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE AT OR BELOW 25°C (77°F), EXCURSIONS PERMITTED UP TO 40°C (104°F)

FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

Mixing and Use Directions

Thoroughly mix the amount of this Type A medicated article according to the directions below with at least an equal amount by weight of feed formula ingredients prior to blending into a complete feed.

Indications for Use	Oxytetracycline Amount	lb of Terramycin 200/ton
CHICKENS		
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	100-200 g/ton Feed continuously for 7-14 days	0.5-1
Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline	400 g/ton Feed continuously for 7-14 days	2
Reduction of mortality due to air	500 g/ton	2.5

sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to oxytetracycline	Feed continuously for 5 days	
WARNING: At 500 g/ton level, withdraw 24 hours before slaughter. Zero-day withdrawal period for lower use levels. Do not use in low-calcium feed containing less than 0.55 dietary calcium. Use in such may result in violative residues. Do not administer to chickens producing eggs for human consumption.		

TURKEYS

Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to oxytetracycline	100 g/ton Feed continuously for 7-14 days	0.5
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to oxytetracycline	200 g/ton Feed continuously for 7-14 days	1
Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline	25 mg/lb of body weight daily Feed continuously for 7-14 days	4.15 ¹

WARNING: At 200 g/ton use level or higher, withdraw 5 days before slaughter. Zero-day withdrawal period for lower use levels. Do not administer to turkeys producing eggs for human consumption.

SWINE

Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	2.5 ²
For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for not more than 14 days	2.5 ²

CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE

For growing cattle (over 400 lb) for reduction of incidence of liver abscesses	75 mg/head/day Feed continuously	0.375 ³
Prevention and treatment of the early stages of shipping fever complex. (Feed 3-5 days before and after arrival in feedlots.)	0.5-2.0 g/head/day	2.5-10 ³
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible	10 mg/lb of body weight daily Feed continuously for 7-14 days	25 ⁴

to oxytetracycline		
SHEEP		
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	6 ⁵

WARNING: 5-day withdrawal before slaughter.

- ¹If bird weighs 10 lb, consuming 0.6 lb of complete feed per day
²Pig consuming 4 lb of complete feed per 100 lb body weight per day
³Include in feed supplement based on consumption of 2 lb of supplement per head per day
⁴If animal weighs 500 lb, consuming 2 lb of supplement per head per day
⁵If lamb weighs 60 lb, consuming 1 lb of supplement per head per day

RESTRICTED DRUG (CALIFORNIA) – USE ONALY AS DIRECTED

NOT FOR HUMAN USE

**SEE BACK PANEL FOR COMPLETE MIXING DIRECTIONS,
USE DIRECTIONS, AND WARNINGS**

**Terramycin is a registered trademark of Pfizer, Inc.,
licensed to Phibro Animal Health Corporation, for Oxytetracycline
PHIBRO and PHIBRO logo design are trademarks owned by
Phibro Animal Health Corporation or its affiliates.**

Phibro Animal Health Corporation, Teaneck, NJ 07666

Net Weight 50 lb (22.7 kg)

NADA #95-143, Approved by FDA

8845000

101-9010-09

Principal Display Panel



Terramycin[®] 200 (oxytetracycline)



FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

Mixing and Use Directions

Thoroughly mix this amount of dry feed per 1/2 metric ton or 1/2 ton of feed for each ingredient per 100 lbs. of dry feed for a complete ton.

Indications for Use	Oxytetracycline Amount	lb of Terramycin 200 ton
CHICKENS		
Control of infectious organisms caused by <i>Mycoplasma</i> or <i>Chlamydia</i>	500 grams Feed continuously for 14 days	0.5
Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> (M.G.) or <i>Mycoplasma meleagridis</i> (M.Me.)	600 grams Feed continuously for 14 days	2
Reduction of mortality due to air sacculitis (or non-infectious) caused by <i>Ascaridia</i> or other coccidia	120 grams Feed continuously for 5 days	2.5
Warnings: Do not use on broilers. Do not use on turkeys. Do not use on ducks. Do not use on geese. Do not use on waterfowl. Do not use on game birds. Do not use on wild birds. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey.		
TURKEYS		
Control of respiratory disease caused by <i>Mycoplasma</i> or <i>Chlamydia</i>	500 grams Feed continuously for 14 days	0.5
Control of infectious organisms caused by <i>Mycoplasma</i> or <i>Chlamydia</i>	600 grams Feed continuously for 14 days	1
Control of parasitic (bacterial) organisms associated with blackhead	25 mg/lb of body weight daily Feed continuously for 14 days	4.15*
Warnings: Do not use on broilers. Do not use on turkeys. Do not use on ducks. Do not use on geese. Do not use on waterfowl. Do not use on game birds. Do not use on wild birds. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey.		
DUCKS		
Prevention of bacterial infections caused by <i>Chlamydia</i> or <i>Mycoplasma</i>	10 mg/lb of body weight daily Feed continuously for 14 days	2.2*
Prevention of bacterial infections caused by <i>Chlamydia</i> or <i>Mycoplasma</i>	10 mg/lb of body weight daily Feed continuously for 14 days	2.2*
Prevention of bacterial infections caused by <i>Chlamydia</i> or <i>Mycoplasma</i>	10 mg/lb of body weight daily Feed continuously for 14 days	2.2*
Warnings: Do not use on broilers. Do not use on turkeys. Do not use on ducks. Do not use on geese. Do not use on waterfowl. Do not use on game birds. Do not use on wild birds. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey.		
CATTLE, BUFF CATTLE, AND ROMANOVSKY GOAT CATTLE		
For young cattle (up to 600 lb) for the reduction of the incidence of bacterial pneumonia	25 mg/head/day Feed continuously	0.531*
Prevention and treatment of the early stages of shipping fever in calves (over 200 lb) before and after arrival at feedlot	0.5-2 g/head/day	2.5-10*
Prevention of bacterial infections caused by <i>Chlamydia</i> or <i>Mycoplasma</i>	10 mg/lb of body weight daily Feed continuously for 14 days	2.2*
Warnings: Do not use on broilers. Do not use on turkeys. Do not use on ducks. Do not use on geese. Do not use on waterfowl. Do not use on game birds. Do not use on wild birds. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey. Do not use on birds of prey.		
SHEEP		
Prevention of bacterial infections caused by <i>Chlamydia</i> or <i>Mycoplasma</i>	10 mg/lb of body weight daily Feed continuously for 14 days	0*

* For 100 lb weight, 10 lb, containing 0.5 lb of complete feed per day.
 * For 200 lb weight, 20 lb, containing 1 lb of complete feed per day.
 * For 300 lb weight, 30 lb, containing 1.5 lb of complete feed per day.
 * For 400 lb weight, 40 lb, containing 2 lb of complete feed per day.
 * For 500 lb weight, 50 lb, containing 2.5 lb of complete feed per day.
 * For 600 lb weight, 60 lb, containing 3 lb of complete feed per day.
 * For 700 lb weight, 70 lb, containing 3.5 lb of complete feed per day.
 * For 800 lb weight, 80 lb, containing 4 lb of complete feed per day.
 * For 900 lb weight, 90 lb, containing 4.5 lb of complete feed per day.
 * For 1000 lb weight, 100 lb, containing 5 lb of complete feed per day.

100-3010-01
 Approved by Phibro Animal Health Corporation, Teaneck, NJ 07666

(oxytetracycline)
Terramycin[®] 200
Phibro
 ANIMAL HEALTH CORPORATION



Terramycin[®] 200 (oxytetracycline)

TYPE A
MEDICATED ARTICLE

Active Drug Ingredient:
 Oxytetracycline (from oxytetracycline dihydrate base) equivalent to
 oxytetracycline hydrochloride 200 g/lb

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.
CAUTION: For use in manufacturing medicated animal feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-attesting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE AT OR BELOW 25°C (77°F), EXCURSIONS PERMITTED UP TO 40°C (104°F)

SEE BACK PANEL FOR COMPLETE MIXING DIRECTIONS, USE DIRECTIONS, AND WARNINGS

TERRAMYCIN is a registered trademark of Pfizer, Inc licensed to Phibro Animal Health Corporation for Oxytetracycline PHIBRO and PHIBRO logo design are trademarks owned by Phibro Animal Health Corporation or its affiliates.
 Phibro Animal Health Corporation, Teaneck, NJ 07666

NADA #95-143, Approved by FDA
 8845000 101-9010-09

Net Weight 50 lb (22.7 kg)



TERRAMYCIN 200

oxytetracycline powder

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-7868
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYTETRACYCLINE HYDROCHLORIDE (UNII: 4U7K4N52ZM) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	200 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
RICE BRAN (UNII: R60QEP13IC)	
SODIUM ALUMINIUM SILICATE (UNII: 058TS43PSM)	
MINERAL OIL (UNII: T5L8T28FGP)	

SUCROSE (UNII: C151H8M554)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-7868-0	22.7 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA095143	03/31/2010	

Labeler - Phibro Animal Health (006989008)

Registrant - Phibro Animal Health (006989008)

Revised: 1/2019

Phibro Animal Health