

PHENYLEPHRINE HYDROCHLORIDE- phenylephrine hydrochloride injection

Medical Purchasing Solutions, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PHENYLEPHRINE HYDROCHLORIDE INJECTION safely and effectively. See full prescribing information for PHENYLEPHRINE HYDROCHLORIDE INJECTION.

PHENYLEPHRINE HYDROCHLORIDE injection, for intravenous use

Initial U.S. Approval: 2012

----- RECENT MAJOR CHANGES -----

Indications and Usage (1)

03/2023

Dosage and Administration (2.1, 2.2)

03/2023

----- INDICATIONS AND USAGE -----

Phenylephrine Hydrochloride Injection 10 mg per mL is alpha-1 adrenergic receptor agonist indicated for increasing blood pressure in adults with clinically important hypotension resulting primarily from vasodilation in the settings of anesthesia and septic shock. (1)

----- DOSAGE AND ADMINISTRATION -----

Phenylephrine Hydrochloride Injection 10 mg per mL:

- MUST BE DILUTED before administration. (2.1)

Dosing for Perioperative Hypotension

- Intravenous bolus administration: 50 mcg to 250 mcg (2.4)
- Intravenous continuous infusion: 0.5 mcg/kg/minute to 1.4 mcg/kg/minute titrated to effect (2.4)

Dosing for Patients with Vasodilatory Shock

- Intravenous continuous infusion: 0.5 mcg/kg/minute to 6 mcg/kg/minute titrated to effect (2.5)

----- DOSAGE FORMS AND STRENGTHS -----

Phenylephrine Hydrochloride Injection:

- 10 mg in 1 mL (10 mg per mL) of phenylephrine hydrochloride in a single-dose vial (3)

----- CONTRAINDICATIONS -----

- Hypersensitivity to the products or any of their components (4)

----- WARNINGS AND PRECAUTIONS -----

- Severe bradycardia and decreased cardiac output: (5.2)
- Extravasation: during intravenous administration may cause necrosis or sloughing of tissue (5.4)
- Concomitant use with oxytocic drugs: pressor effect of sympathomimetic pressor amines is potentiated (5.5)
- Allergic-type reactions with phenylephrine hydrochloride injection 10 mg per mL: Sulfite (5.6)

----- ADVERSE REACTIONS -----

Most common adverse reactions: nausea and vomiting, headache, nervousness (6)

To report SUSPECTED ADVERSE REACTIONS, contact Meitheal Pharmaceuticals Inc. at 1-844-824-8426 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- DRUG INTERACTIONS -----

- Agonistic effects with monoamine oxidase inhibitors (MAOI), β -adrenergic blocking agents, α -2 adrenergic agonists, steroids, tricyclic antidepressants, norepinephrine transport inhibitors, ergot alkaloids, centrally-acting sympatholytic agents and atropine sulfate (7.1)
- Antagonistic effects on and by α -adrenergic blocking agents (7.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 5/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 General Administration Instructions

2.2 Preparation of Phenylephrine Hydrochloride Injection

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Exacerbation of Angina, Heart Failure, or Pulmonary Arterial Hypertension

5.2 Bradycardia

5.3 Risk in Patients with Autonomic Dysfunction

5.4 Skin and Subcutaneous Necrosis

5.5 Pressor Effect with Concomitant Oxytocic Drugs

5.6 Allergic Reactions

5.7 Peripheral and Visceral Ischemia

5.8 Renal Toxicity

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

7.1 Agonists

7.2 Antagonists

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Phenylephrine Hydrochloride Injection 10 mg per mL is indicated for increasing blood

pressure in adults with clinically important hypotension resulting primarily from vasodilation in the settings of anesthesia and septic shock.

2 DOSAGE AND ADMINISTRATION

2.1 General Administration Instructions

- Phenylephrine hydrochloride injection 10 mg per mL formulation (10 mg per mL) MUST BE DILUTED before administration as an intravenous bolus or for continuous intravenous infusion. The diluted solution should not be held for more than 4 hours at room temperature or for more than 24 hours under refrigerated conditions (2° to 8°C).

Parenteral drug products should be inspected for particulate matter and discoloration prior to administration. Discard any unused portion.

During phenylephrine hydrochloride injection 10 mg per mL administration:

- Correct intravascular volume depletion.
- Correct acidosis. Acidosis may reduce the effectiveness of phenylephrine.

2.2 Preparation of Phenylephrine Hydrochloride Injection

Preparing a 100 mcg per mL Solution for Intravenous Bolus Administration

For intravenous bolus administration, withdraw 10 mg (1 mL of a 10 mg per mL concentration) of phenylephrine hydrochloride injection 10 mg per mL and dilute with 99 mL of 5% Dextrose Injection, USP or 0.9% Sodium Chloride Injection, USP. This will yield a final concentration of 100 mcg per mL. Withdraw an appropriate dose from the 100 mcg per mL solution prior to intravenous bolus administration.

Preparing a 20 mcg per mL Solution for Continuous Intravenous Infusion

For continuous intravenous infusion, withdraw 10 mg (1 mL of 10 mg per mL concentration) of phenylephrine hydrochloride injection 10 mg per mL and add to 500 mL of 5% Dextrose Injection, USP or 0.9% Sodium Chloride Injection, USP (providing a final concentration of 20 mcg per mL).

Dosing for Perioperative Setting

In adult patients undergoing surgical procedures with either neuraxial anesthesia or general anesthesia:

- Phenylephrine hydrochloride injection 10 mg per mL (diluted to 100 mcg per mL): 50 mcg to 250 mcg by intravenous bolus administration. The most frequently reported initial bolus dose is 50 mcg or 100 mcg.
- Phenylephrine hydrochloride injection 10 mg per mL (diluted to 20 mcg per mL): 0.5 mcg/kg/min to 1.4 mcg/kg/min by intravenous continuous infusion, titrated to blood pressure goal.

Dosing for Septic or Other Vasodilatory Shock (Phenylephrine Hydrochloride Injection 10 mg per mL only)

In adult patients with septic or other vasodilatory shock:

- No bolus.
- 0.5 mcg/kg/min to 6 mcg/kg/min by intravenous continuous infusion, titrated to blood

pressure goal. Doses above 6 mcg/kg/min do not show significant incremental increase in blood pressure.

3 DOSAGE FORMS AND STRENGTHS

Phenylephrine Hydrochloride Injection, USP: 10 mg per mL phenylephrine hydrochloride is a clear, colorless, essentially free of visible foreign matter sterile parenteral solution, available in:

- 10 mg in 1 mL (10 mg per mL) of phenylephrine hydrochloride in a single-dose vial

4 CONTRAINDICATIONS

The use of phenylephrine hydrochloride injection 10 mg per mL is contraindicated in patients with:

- Hypersensitivity to the products or any of their components

5 WARNINGS AND PRECAUTIONS

5.1 Exacerbation of Angina, Heart Failure, or Pulmonary Arterial Hypertension

Because of its pressor effects, phenylephrine hydrochloride can precipitate angina in patients with severe arteriosclerosis or history of angina, exacerbate underlying heart failure, and increase pulmonary arterial pressure.

5.2 Bradycardia

Phenylephrine hydrochloride injection 10 mg per mL can cause severe bradycardia and decreased cardiac output.

5.3 Risk in Patients with Autonomic Dysfunction

The pressor response to adrenergic drugs, including phenylephrine, can be increased in patients with autonomic dysfunction, as may occur with spinal cord injuries.

5.4 Skin and Subcutaneous Necrosis

Extravasation of phenylephrine can cause necrosis or sloughing of tissue.

5.5 Pressor Effect with Concomitant Oxytocic Drugs

Oxytocic drugs potentiate the pressor effect of sympathomimetic pressor amines including phenylephrine hydrochloride injection 10 mg per mL [see *Drug Interactions* (7.1)] , with the potential for hemorrhagic stroke.

5.6 Allergic Reactions

Phenylephrine hydrochloride injection 10 mg per mL contains sodium metabisulfite, a sulfite that may cause allergic-type reactions, including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.

5.7 Peripheral and Visceral Ischemia

Phenylephrine hydrochloride injection 10 mg per mL can cause excessive peripheral and visceral vasoconstriction and ischemia to vital organs, particularly in patients with extensive peripheral vascular disease.

5.8 Renal Toxicity

Phenylephrine hydrochloride injection 10 mg per mL can increase the need for renal replacement therapy in patients with septic shock. Monitor renal function.

6 ADVERSE REACTIONS

The following adverse reactions associated with the use of phenylephrine hydrochloride were identified in the literature. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency reliably or to establish a causal relationship to drug exposure.

Cardiac disorders: Bradycardia, AV block, ventricular extrasystoles, myocardial ischemia

Gastrointestinal disorders: Nausea, vomiting

General disorders and administrative site conditions: Chest pain, extravasation

Immune system disorders: Sulfite sensitivity

Nervous system disorders: Headache, nervousness, paresthesia, tremor

Psychiatric disorders: Excitability

Respiratory: Pulmonary edema, rales

Skin and subcutaneous tissue disorders: Diaphoresis, pallor, piloerection, skin blanching, skin necrosis with extravasation

Vascular disorders: Hypertensive crisis

7 DRUG INTERACTIONS

7.1 Agonists

The pressor effect of phenylephrine hydrochloride is *increased* in patients receiving:

- Monoamine oxidase inhibitors (MAOI), such as selegiline.
- β -adrenergic blockers
- α -2 adrenergic agonists, such as clonidine
- Steroids
- Tricyclic antidepressants
- Norepinephrine transport inhibitors, such as atomoxetine
- Ergot alkaloids, such as methylergonovine maleate
- Centrally-acting sympatholytic agents, such as guanfacine or reserpine
- Atropine sulfate

7.2 Antagonists

α -adrenergic blocking agents, including phenothiazines (e.g., chlorpromazine) and amiodarone block phenylephrine and are in turn blocked by phenylephrine.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

In animal reproductive and developmental studies, decreased fetal body weights were noted at 0.4 times the human daily dose (HDD) of 10 mg. No malformations were reported, however, an increased incidence of agenesis of the intermediate lobe of the lung, a visceral variation, was reported at levels as low as 0.08 times the HDD.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

No malformations were noted when normotensive pregnant rats were treated with a single daily intravenous bolus dose of 50 mcg, 150 mcg, or 300/200 mcg/kg phenylephrine hydrochloride from Gestation Day 6 to 17 (high dose is 0.3/0.2 times the human daily dose (HDD) of 10 mg/day based on body surface area). Evidence of maternal toxicity, including mortality, was noted at the highest tested dose of 300/200 mcg/kg.

Decreased fetal body weights but no clear treatment-related malformations were reported when normotensive pregnant rabbits were treated with a single daily intravenous bolus dose of 40 mcg, 100 mcg and 200 mcg/kg (0.08, 0.2, and 0.4 times the HDD based on body surface area) phenylephrine hydrochloride from Gestation Day 7 to 19. Maternal toxicity, as manifested by decreased food consumption and body weight gain at all doses. An increased incidence of agenesis of the intermediate lobe of the lung, a visceral variation, was noted in all treatment groups compared to controls.

No adverse effects on the offspring were reported when pregnant rats were treated via a single daily intravenous bolus dose of up to 200 mcg/day phenylephrine hydrochloride (0.2 times the HDD based on body surface area) from Gestation Day 6 to Lactation Day 20.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Clinical studies of phenylephrine did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be

cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Hepatic Impairment

In patients with liver cirrhosis [Child Pugh Class A (n=3), Class B (n=5) and Class C (n=1)], dose-response data indicate decreased responsiveness to phenylephrine. Consider using larger doses than usual in hepatic impaired subjects.

8.7 Renal Impairment


In patients with end stage renal disease (ESRD) undergoing hemodialysis, dose-response data indicates increased responsiveness to phenylephrine. Consider using lower doses of phenylephrine hydrochloride in ESRD patients.

10 OVERDOSAGE

Overdose of phenylephrine hydrochloride injection 10 mg per mL can cause a rapid rise in blood pressure. Symptoms of overdose include headache, vomiting, hypertension, reflex bradycardia, and cardiac arrhythmias including ventricular extrasystoles and ventricular tachycardia, and may cause a sensation of fullness in the head and tingling of the extremities. Consider using an α -adrenergic antagonist.

11 DESCRIPTION

Phenylephrine Hydrochloride Injection, USP contains active pharmaceutical ingredient phenylephrine in the form of hydrochloride salt. Phenylephrine is a synthetic sympathomimetic agent in sterile form for parenteral injection. Phenylephrine hydrochloride chemical name is (-)- *m*-Hydroxy- α [(methylamino)methyl]benzyl alcohol hydrochloride and has the following structural formula:



Phenylephrine hydrochloride is very soluble in water, freely soluble in ethanol, and insoluble in chloroform and ethyl ether. Phenylephrine hydrochloride is sensitive to light.

Phenylephrine Hydrochloride Injection, USP, 10 mg per mL

Phenylephrine Hydrochloride Injection, USP is a clear, colorless, aqueous solution that is essentially free of visible foreign matter. It MUST BE DILUTED before administration as bolus intravenous infusion or continuous intravenous infusion.

Each mL contains: 10 mg of Phenylephrine Hydrochloride (equivalent to 8.2 mg of phenylephrine base); 3.5 mg of Sodium Chloride USP as tonicity agent; 1 mg of Citric Acid Monohydrate USP and 4 mg of Sodium Citrate Dihydrate USP, as buffering agents; 2 mg of Sodium Metabisulfite NF, as antioxidant, and Sodium Hydroxide NF and Hydrochloric Acid NF, as pH adjusters in Water for Injection. Phenylephrine

Hydrochloride Injection, USP pH range is 3.0 to 6.5.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Phenylephrine hydrochloride is an α -1 adrenergic receptor agonist.

12.2 Pharmacodynamics

Phenylephrine is the active moiety. Metabolites are inactive at both the α -1 and α -2 adrenergic receptors. Following parenteral administration of phenylephrine hydrochloride, increases in systolic blood pressure, diastolic blood pressure, mean arterial blood pressure, and total peripheral vascular resistance are observed. The onset of blood pressure increase following an intravenous bolus phenylephrine hydrochloride administration is rapid and the effect may persist for up to 20 minutes. As mean arterial pressure increases following parenteral doses, vagal activity also increases, resulting in reflex bradycardia.

Most vascular beds are constricted, including renal, splanchnic, and hepatic.

12.3 Pharmacokinetics

Following an intravenous infusion of phenylephrine hydrochloride, the effective half-life was approximately 5 minutes. The steady-state volume of distribution (340 L) exceeded the body volume by a factor of 5, suggesting a high distribution into certain organ compartments. The average total serum clearance (2095 mL/min) was close to one-third of the cardiac output.

A mass balance study showed that phenylephrine is extensively metabolized by the liver with only 12% of the dose excreted unchanged in the urine. Deamination by monoamino oxidase is the primary metabolic pathway resulting in the formation of the major metabolite (m-hydroxymandelic acid) which accounts for 57% of the total administered dose.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term animal studies that evaluated the carcinogenic potential of orally administered phenylephrine hydrochloride in F344/N rats and B6C3F1 mice were completed by the National Toxicology Program using the dietary route of administration. There was no evidence of carcinogenicity in mice administered approximately 270 mg/kg/day (131-times the human daily dose (HDD) of 10 mg/day based on body surface area) or rats administered approximately 50 mg/kg/day (48-times the HDD based on body surface area comparisons).

Mutagenesis

Phenylephrine hydrochloride tested negative in the *in vitro* bacterial reverse mutation assay (*S. typhimurium* strains TA98, TA100, TA1535 and TA1537), the *in*

vitro chromosomal aberrations assay, the *in vitro* sister chromatid exchange assay, and the *in vivo* rat micronucleus assay. Positive results were reported in only one of two replicates of the *in vitro* mouse lymphoma assay.

Impairment of Fertility

No adverse effects on fertility or early embryonic development were noted when phenylephrine hydrochloride was administered at doses of 50 mcg, 100 mcg, or 200 mcg/kg/day (up to 0.2 times HDD of 10 mg/60 kg/day based on body surface area) via single daily bolus injection for 28 days prior to mating to male rats or for 14 days prior to mating through Gestation Day 7 to female rats.

14 CLINICAL STUDIES

Increases in systolic and mean blood pressure following administration of phenylephrine were observed in 42 literature-based studies in the perioperative setting, including 26 studies where phenylephrine was used in low-risk (ASA 1 and 2) pregnant women undergoing neuraxial anesthesia during cesarean delivery, 3 studies in non-obstetric surgery under neuraxial anesthesia, and 13 studies in patients undergoing surgery under general anesthesia. Mean arterial blood pressure increases were also observed in two double-blind, active-controlled studies in patients with septic shock.

16 HOW SUPPLIED/STORAGE AND HANDLING

Phenylephrine Hydrochloride Injection, USP, 10 mg per mL, is supplied as follows:

NDC	Phenylephrine Hydrochloride Injection Package Factor (10 mg per mL)
7 1 2 8 8 - 807-10 mg per 1 mL Single-Dose Vial 02	25 vials per carton

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature]. **Protect from light.** Keep covered in carton until time of use.

The 1 mL vials are for single-dose only. The diluted solution should not be held for more than 4 hours at room temperature or for more than 24 hours under refrigerated conditions (2° to 8°C). Discard any unused portion.

Sterile, Nonpyrogenic.

The container closure is not made with natural rubber latex.

17 PATIENT COUNSELING INFORMATION

Inform patients, families, or caregivers that the primary side effect of phenylephrine hydrochloride injection is hypertension and, rarely, hypertensive crisis. Patients may experience bradycardia (slow heart rate), which in some cases may produce heart block or other cardiac arrhythmias, extra ventricular beats, myocardial ischemia in patients with underlying cardiac disease, and pulmonary edema (fluid in the lungs) or rales. Common, less serious symptoms include the following:

- chest pain
- skin or tissue damage if the drug leaks out of the venous catheter into the surrounding tissue
- headache, nervousness, tremor, numbness/tingling (paresthesias) in hands or feet
- nausea, vomiting
- excitability, dizziness, sweating, flushing

meitheal[®]

Mfd. for Meitheal Pharmaceuticals

Chicago, IL 60631 (USA)

©2023 Meitheal Pharmaceuticals Inc.

Mfd. by Nanjing King-Friend Biochemical Pharmaceutical Co., Ltd.

Nanjing, China 210061

Revised: May 2023

8H7EAM9-01

PRINCIPAL DISPLAY PANEL - VIAL LABEL

Rx Only

Phenylephrine HCl Injection, USP

10 mg per mL

For Intravenous Use

Dilute Before Use

1 mL Single-Dose Vial

DISCARD UNUSED PORTION



PRINCIPAL DISPLAY PANEL - OUTER PACKAGE

NDC 71872-7343-1

Phenylephrine HCl Injection, USP

10 mg per mL

For Intravenous Use

Dilute Before Use

1 x 1 mL Single-Dose Vials

DISCARD UNUSED PORTION

Rx Only



Rxonly



(01) 0 0371872 73431 8
(17) 000101
(10) XXXX-XXXXXX
(21) 7287A000001

Phenylephrine HCl Injection, USP
10 mg per mL

Qty: 1 vial Lot # XXXX-XXXXXX Exp: 01/01/1900

For Intravenous Use Dilute Before Use.DISCARD UNUSED PORTION.

LIGHT SENSITIVE: Keep covered in carton until time of use.

Each mL contains: Phenylephrine Hydrochloride 10 mg; Sodium Chloride 3.5 mg; Sodium Citrate Dihydrate 4 mg; Citric Acid Monohydrate 1 mg; Sodium Metabisulfite 2 mg; Water for Injection q.s. pH adjusted with Sodium Hydroxide and/or Hydrochloric Acid if necessary. pH 3.0 to 6.5.

USUAL DOSAGE: See Package Insert

PROTECT FROM LIGHT. Keep covered in carton until time of use.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature]. Sterile, Nonpyrogenic. The container closure is not made with natural rubber latex.

Manufactured by: Melitreal Pharmaceuticals Chicago, IL 60631 (USA)
Manufacturer NDC: 71288-0807-02

Repackaged
& Distributed By:



Medical Purchasing Solutions
Scottsdale, AZ 85260
www.medicalpurchasingsolutions.com

PHENYLEPHRINE HYDROCHLORIDE

phenylephrine hydrochloride injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71872-7343(NDC:71288-807)
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 1 mL
---	-----------------------------	------------------

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71872-7343-1	1 in 1 BAG	01/20/2025	
1		1 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210334	04/27/2018	

Labeler - Medical Purchasing Solutions, LLC (601458529)

Establishment

Name	Address	ID/FEI	Business Operations
Medical Purchasing Solutions, LLC		601458529	repack(71872-7343)